Bounds on the Picard rank of toric Fano varieties with minimal curve constraints

Roya Beheshti, Ben Wormleighton

arXiv:2202.01852

http://arxiv.org/abs/2202.01852v4

Minimal curves constrain the fan

Let $X = X_\Sigma$ be a smooth toric Fano variety of dimension n. Write $G(\Sigma)$ for the primitive ray generators. Then

$$\rho(X) = |G(\Sigma)| - n.$$

A component $\mathcal{C}$ of rational curves is *minimal* when it covers X and the subfamily through a general point is complete. Its degree is $-K_X \cdot C$, and its codegree is $q = n + 1 - \deg(\mathcal{C})$. In the toric setting such components correspond to primitive collections $\{v_1, \ldots, v_{n+1-q}\}$ satisfying $v_1 + \cdots + v_{n+1-q} = 0$.

Main theorem (Beheshti–Wormleighton, Theorems 3.2 and 4.1)**.** *For every $q \geq 0$ there is a constant A_q, depending only on q, such that every smooth toric Fano variety carrying a minimal component of codegree q satisfies*

$$\rho(X) \leq A_q.$$

For $q = 2$ one may take $A_2 = 5$, and this bound is sharp in every dimension.

Thus a family whose anticanonical degree stays within a fixed amount of the maximum forces a Picard bound independent of dimension. The general result implies the Chen–Fu–Hwang inequality in sufficiently high dimension for each fixed codegree. In codegree at most two, the paper proves both that inequality and its stronger proposed bound.

Ingredient 1: only finitely many primitive extensions

A primitive collection is a minimal set of rays which fails to span a cone. Proposition 3.1 says that, for each $m \geq 1$, there is a dimension-independent constant C_m such that a fixed cone $\langle v_1, \ldots, v_m \rangle$ belongs to at most C_m primitive collections of the form $\{v_1, \ldots, v_m, w\}$.

The proof inducts on m and compares the primitive relations attached to two putative extensions. If too many extensions existed, one could choose one whose new ray forms cones with the relevant proper subsets on the other side of a relation. Subtracting the two relations would then produce an effective curve class of nonpositive anticanonical degree. This contradicts the ampleness of $-K_X$. The Fano condition has thereby converted a local fan configuration into a uniform counting statement.

Ingredient 2: cover all positive combinations by few cones

Proposition 3.4 gives a second dimension-free statement. For any q rays, their nonnegative span is covered by at most M_q cones of Σ, each of dimension at most $q2^q$. Starting from the effective relation obtained by placing the sum of the rays in its minimal cone, the proof inducts on anticanonical degree. Reid's theorem on an extremal primitive relation supplies the covering cones in degree one. In the inductive step one omits a left-hand generator and adjoins the generators of that minimal cone; if the resulting set is not a cone, its new effective relation has strictly smaller anticanonical degree.

Lemma 3.3 is then applied recursively to the long zero-sum primitive collection representing $\mathcal{C}$. It produces auxiliary blocks whose total excess is exactly q. Every remaining ray $w \in G(\Sigma)$ can be paired, modulo the span of the original collection, with a nonnegative combination of those auxiliary rays. Proposition 3.4 places that combination in one of finitely many bounded-dimensional cones. The ray w cannot join the vertices of that cone to form a cone of Σ; Proposition 3.1 therefore bounds the number of possible w solely in terms of q. Hence $|G(\Sigma)| \leq n + A_q$, which is precisely $\rho(X) \leq A_q$.

Ingredient 3: the sharp two-dimensional quotient

For codegree two, the proof keeps the geometry rather than the enormous recursive constants. After fixing the appropriate part of the zero-sum primitive collection, the fan is projected to a two-dimensional quotient. Suppose three projected rays surround the origin and each of the three pairs spans a cone of Σ. Their triple is then a primitive collection, whose relation has only three possible shapes: its sum is zero, is one ray, or is a sum of two rays from the fixed set.

Reid's cone theorem propagates each extremal relation to neighbouring cones. Repeatedly, any proposed extra ray would combine with these cones to give an effective class of anticanonical degree zero, impossible on a Fano variety. Lemma 4.2 separately handles an opposite pair $\{x, -x\}$ and already forces $\rho(X) \leq 5$. If no pairwise-compatible triple as above exists, the proof chooses a pair spanning a maximal projected cone. Every projected ray outside it must fail to span a cone with one of the chosen rays; uniqueness of the possible non-neighbour, together with Lemma 4.2's exclusion of opposite pairs, covers the quotient plane by the resulting cones and bounds all remaining rays. Dimensions four and five are checked from classified toric Fano lists; the fan argument treats dimensions at least six.

Sharp model, hard point, and scope

Let dP_6 be the toric del Pezzo surface of degree six. For

$$X = \mathbf{P}^{n-2} \times dP_6,$$

a line in the first factor has anticanonical degree $n-1$, hence codegree two, while $\rho(X) = 1 + \rho(dP_6) = 5$. This model explains both the dimension-free nature and the sharp value of the special bound.

The hardest part of the general theorem is not a classification: it is the two-layer uniformity that bounds primitive extensions and then covers every positive combination by finitely many fan cones, all independently of n. The resulting A_q is not claimed to be sharp or usefully explicit. The sharp argument is special to $q = 2$, smoothness and the Fano positivity are essential throughout, and the paper leaves the proposed linear bound $\rho(X) \leq 2q + 2$ open for general codegree and also asks about singular toric Fano analogues.