

K-stability of birationally superrigid Fano varieties

Charlie Stibitz, Ziquan Zhuang

arXiv:1802.08381

<http://arxiv.org/abs/1802.08381v2>

Two stability theories meet

Work over $\mathbf{C}$. A $\mathbf{Q}$ -Fano variety X is normal, klt, and has $-K_X$ ample. Its alpha invariant is

$$\alpha(X) = \sup\{t > 0 : (X, tD) \text{ is log canonical for every effective } \mathbf{Q}\text{-divisor } D \sim_{\mathbf{Q}} -K_X\}.$$

For a Picard-rank-one Fano variety, birational superrigidity is equivalent to $\mathbf{Q}$ -factorial terminal singularities together with canonicity of (X, M) for every movable boundary $M \sim_{\mathbf{Q}} -K_X$ (Theorem 2.3). This resembles K-stability: both conditions control singularities of anticanonical data, but one uses movable systems and the other divisorial valuations.

Main theorem (Stibitz–Zhuang, Theorem 1.2). *Let X be a Picard-rank-one $\mathbf{Q}$ -Fano variety. Assume that (X, M) is log canonical for every movable boundary $M \sim_{\mathbf{Q}} -K_X$ (in particular this holds if X is birationally superrigid). If $\alpha(X) \geq \frac{1}{2}$, then X is K-semistable; if $\alpha(X) > \frac{1}{2}$, then X is K-stable.*

The equality case is sharper for the original superrigid hypothesis: Corollary 3.1 shows that a birationally superrigid X with $\alpha(X) \geq \frac{1}{2}$ is already K-stable.

Ingredient I: translate K-stability into one integral

For a dreamy prime divisor F over X , choose a projective birational model $\pi : Y \rightarrow X$ on which F is a prime divisor. Write $A = A_X(F)$, let $\tau = \tau(F)$ be the pseudo-effective threshold, put $V = (-K_X)^n$, and set

$$b = \frac{1}{V} \int_0^\tau \text{vol}_X(-K_X - xF) dx. \quad (1)$$

The Fujita–Li valuative criterion says $\beta(F) = V(A - b)$; hence K-semistability is exactly $b \leq A$ for every dreamy F (Theorem 2.2). Assume for contradiction that $b > A$.

The derivative of volume is encoded by the restricted volume $w(x) = \text{vol}_{Y|F}(-\pi^*K_X - xF)$. Integration by parts gives the barycenter identity

$$\int_0^\tau (x - b)w(x) dx = 0. \quad (2)$$

Thus b is the center of mass of the nonnegative density w .

Ingredient II: movable canonicity isolates one fixed divisor

Because $\rho(X) = 1$, every divisor class is a multiple of $-K_X$. If two distinct irreducible anticanonical divisors had F -order greater than A , their pencil would yield a movable anticanonical boundary that is not log canonical. Therefore at most one irreducible $D \sim_{\mathbf{Q}} -K_X$ can satisfy $\text{ord}_F(D) > A$. Under the assumption $b > A$, this unique fixed divisor occurs with $\text{ord}_F(D) = \tau$.

For $x > A$, it is forced into the stable base locus with multiplicity at least $(x - A)/(\tau - A)$, giving the exact tail profile

$$w(x) = \left(\frac{\tau - x}{\tau - A} \right)^{n-1} w(A). \quad (3)$$

For $0 \leq x \leq A$, log-concavity of restricted volume gives the opposite one-sided estimate

$$w(x) \geq \left(\frac{x}{A} \right)^{n-1} w(A). \quad (4)$$

Ingredient III: the half-threshold contradiction

Insert (3)–(4) into (2). Since $x - b < 0$ on $[0, A]$, replacing the left part by its lower bound and using the exact right tail yields

$$0 \leq \int_0^A (x - b) \left(\frac{x}{A}\right)^{n-1} dx + \int_A^\tau (x - b) \left(\frac{\tau - x}{\tau - A}\right)^{n-1} dx.$$

The elementary integrals reduce this to

$$\frac{\tau - 2A}{n(n+1)} + \frac{A - b}{n} \geq 0. \quad (5)$$

Because $b > A$, (5) forces $\tau > 2A$. The divisor D above then satisfies

$$\text{lct}(X; D) \leq \frac{A_X(F)}{\text{ord}_F(D)} = \frac{A}{\tau} < \frac{1}{2},$$

contradicting $\alpha(X) \geq \frac{1}{2}$. Hence $b \leq A$ for every dreamy prime divisor, proving K-semistability. With the strict alpha inequality, the same argument rules out equality and proves K-stability.

The toy picture is literal center of mass: equations (3)–(4) prevent too much restricted-volume mass from lying to the right of the discrepancy A . If its barycenter b nevertheless crosses A , the tail must extend beyond $2A$, and that extension is exactly a divisor witnessing an alpha invariant below one half. The technically hard step is not the final integration but the base-locus argument making the tail profile exact.

Why equality is stable for a superrigid variety

If X were strictly K-semistable, choose dreamy F with $\beta(F) = 0$, so $b = A$. Let η be the largest order obtained from an anticanonical divisor not containing the fixed D . The alpha bound gives $\tau \leq 2A$, while movable-boundary log canonicity gives $\eta \leq A$. Repeating the integral estimate with η gives $(\tau - 2A) + (n - 1)(\eta - A) \geq 0$, so both upper bounds are equalities: $\eta = A$ and $\tau = 2A$. Consequently F is a log-canonical place both for $(X, \frac{1}{2}D)$ and for a movable boundary $M \sim_{\mathbf{Q}} -K_X$ (Corollary 3.1). Birational superrigidity makes every such movable pair canonical, excluding this exceptional log-canonical place; strict semistability is impossible.

Applications, hardest point, and limits

The theorem recovers K-stability of every smooth hypersurface of degree $n \geq 4$ in $\mathbf{P}^n$ (Corollary 1.3). For a Picard-rank-one X with the same movable-boundary log-canonicity hypothesis (in particular, if X is birationally superrigid), if also $\dim X = n \geq 3$, $-K_X$ generates $\text{Cl}(X)$, and $|-K_X|$ is base-point free, Theorem 1.6 proves $\alpha(X) \geq 1/(n+1)$ by Nadel vanishing and a movable-system contradiction. The paper also proves separatedness for families whose two central fibers are birationally superrigid (Theorem 1.7).

The main K-stability result retains the Picard-number-one hypothesis and requires the movable-boundary condition as well as the alpha threshold; it does not prove that every birationally superrigid Fano variety has $\alpha(X) > \frac{1}{2}$. That assertion is posed as Question 1.5.