

On positivity of the CM line bundle on K-moduli spaces

Chenyang Xu, Ziquan Zhuang

arXiv:1912.12961

<http://arxiv.org/abs/1912.12961v3>

A polarization for K-moduli

Work over an algebraically closed field of characteristic 0, and fix dimension n and anticanonical volume v . The good moduli space $M_{n,v}^{\text{Kps}}$ parametrizes n -dimensional K-polystable Fano varieties of volume v and carries the CM $\mathbf{Q}$ -line bundle Λ_{CM} . Nefness was known on proper K-moduli families, but ampleness is subtler when the fibres have positive-dimensional automorphism groups: ordinary uniform K-stability is too rigid, and a degeneration may be harmless after correction by a one-parameter subgroup of the automorphism torus.

For a maximal torus $T \subset \text{Aut}(X)$, reduced uniform K-stability measures a test configuration only after minimizing its norm over all such torus twists. Equivalently, there is $\eta > 0$ with $D^{\text{NA}} \geq \eta J_T^{\text{NA}}$, where D^{NA} is the non-Archimedean Ding invariant and J_T^{NA} is the reduced norm.

Main theorem (Theorem 1.1). *Let $M \subset M_{n,v}^{\text{Kps}}$ be a proper algebraic subspace. If every geometric point of M parametrizes a reduced uniformly K-stable Fano variety, then $\Lambda_{\text{CM}}|_M$ is ample.*

Over $\mathbf{C}$, smoothable K-polystable Fanos admit Kähler–Einstein metrics and hence satisfy the stability hypothesis; the proper smoothable component is therefore projective (Corollary 1.2).

Filtrations detect stability and CM degree

For a log Fano pair (X, Δ) choose $r > 0$ with $L = -r(K_X + \Delta)$ Cartier and set $R = \bigoplus_m H^0(X, mL)$. A linearly bounded multiplicative filtration $\mathcal{F}$ has an expected weight $S(\mathcal{F})$. From the log canonical thresholds of its graded base ideals the authors define a δ -log-canonical slope $\mu_\delta(\mathcal{F})$ and

$$\beta_\delta(\mathcal{F}) = \frac{\mu_\delta(\mathcal{F}) - S(\mathcal{F})}{r}.$$

Theorem 1.4 says that nonnegativity of these invariants for all filtrations characterizes K-semistability ($\delta = 1$), uniform K-stability (one fixed $\delta > 1$), and reduced uniform K-stability (after a twist $\mathcal{F} \mapsto \mathcal{F}_\xi$ for some $\xi \in \text{Hom}(\mathbf{G}_m, T)_{\mathbf{R}}$). The key comparison is $\beta(\mathcal{F}) \geq D^{\text{NA}}(\mathcal{F})$.

Now let $f: (X, \Delta) \rightarrow C$ be a generic log Fano family over a smooth projective curve. Fix $t \in C$ for which (X_t, Δ_t) is log Fano and retain the assumption that $-r(K_X + \Delta)$ is Cartier. If $\mathcal{F}_{\text{HN}}$ is the Harder–Narasimhan filtration of the direct-image section ring, Proposition 4.6 converts stability into geometry on the base:

$$\deg \lambda_{f,\Delta} \geq (n+1)(-K_{X_t} - \Delta_t)^n \beta_{X_t, \Delta_t}(\mathcal{F}_{\text{HN}}).$$

Thus K-semistability gives nefness. More strongly, if $\beta_\delta(\mathcal{F}_{\text{HN}}) \geq 0$ for some $\delta > 1$, Proposition 4.9 gives the uniform nef threshold

$$-(K_{X/C} + \Delta) + \frac{\delta}{(n+1)v(\delta-1)} f^* \lambda_{f,\Delta} \quad \text{nef}, \quad v = (-K_{X_t} - \Delta_t)^n.$$

Turning a torus correction into a family

Reduced uniform stability supplies the required nonnegative β_δ only after twisting by a vector ξ . Theorem 4.18 makes $\delta > 1$ uniform in terms of a stability slope, dimension, and an alpha-invariant bound. For an HN filtration, Proposition 5.6 moreover chooses ξ rational. After a finite base change whose degree clears its denominators, replace it by the resulting integral cocharacter; for a Cartier divisor A on the base, Section 5 realizes that integral twist geometrically by

$$X_\xi = \text{Proj}_C \bigoplus_{m, \chi} R_{m, \chi} \otimes \mathcal{O}_C(\langle \chi, \xi \rangle A).$$

Locally this is the original family, its HN filtration is $\mathcal{F}_{\text{HN}, \xi}$, and its CM class is unchanged because the Futaki character vanishes for K-semistable fibres. A concrete model is the passage from $\mathbf{P}^1 \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ to a Hirzebruch surface $\mathbb{F}_e$: the twist changes the global gluing while preserving the fibre. Here the same operation removes the automorphism direction that obstructs a useful nef threshold.

From curves to bigness, then ampleness

Over a normal projective base S , take an arbitrary covering family of curves and make the rational twist above. Write $g: (X, \Delta) \rightarrow U$ for the resulting birational pullback family, C for the geometric generic curve of the covering family, and $q: C \rightarrow S$ for the induced map. The strengthened ampleness lemma and product trick compare intersection numbers of a nef auxiliary divisor L_Y with an ample divisor H on S . Perturbing the boundary gives the opposite comparison with CM degree. Together they prove the uniform estimate of Theorem 7.2,

$$\deg(\lambda_{g, \Delta}|_C) \geq c_0 \deg(q^* H),$$

where $c_0 > 0$ is independent of the chosen covering curve and birational pullback. Consequently $\lambda_{f, \Delta} - c_0 H$ is pseudo-effective, so the CM class is big for every maximal-variation family of reduced uniformly K-stable pairs.

To apply this on the good moduli space, pass to a generically finite cover of a proper subspace, lift its generic point to the K-moduli stack, spread out the resulting family, and split a maximal automorphism torus after another finite cover. The curve estimate then proves bigness of the pulled-back CM class (Theorem 7.8). Since the CM class is already nef, applying this bigness statement to every subspace and using Nakai–Moishezon yields ampleness. The technically hardest part is uniformity in the curve and the twist; it is precisely why the birational ampleness lemma of Section 6 is needed. This note omits its determinant construction and the log-boundary perturbation estimates.