

On upper bounds of arithmetic degrees

Yohsuke Matsuzawa

arXiv:1606.00598

<http://arxiv.org/abs/1606.00598v3>

A geometric ceiling for arithmetic growth

Let X be a smooth projective variety over $\overline{\mathbf{Q}}$ and $f : X \dashrightarrow X$ a dominant rational map. Write I_f for its indeterminacy locus and

$$X_f(\overline{\mathbf{Q}}) = \{P : f^n(P) \notin I_f \text{ for every } n \geq 0\}.$$

For an ample height h_X , put $h_X^+ = \max\{h_X, 1\}$. The first dynamical degree and upper arithmetic degree are

$$\delta_f = \lim_{n \rightarrow \infty} \rho((f^n)^* |N^1(X)_{\mathbf{R}}|^{1/n}, \quad \overline{\alpha}_f(P) = \limsup_{n \rightarrow \infty} h_X^+(f^n(P))^{1/n}.$$

Main theorem (Matsuzawa, Theorem 1.4). *For every $\varepsilon > 0$ there is $C > 0$ such that, simultaneously for all $n \geq 0$ and $P \in X_f(\overline{\mathbf{Q}})$,*

$$h_X^+(f^n(P)) \leq C(\delta_f + \varepsilon)^n h_X^+(P). \quad (1)$$

Consequently $\overline{\alpha}_f(P) \leq \delta_f$ for every such P .

This supplies the universal “easy inequality” in the arithmetic-degree conjecture: geometry bounds arithmetic orbit growth even when pullback is not functorial under iteration. The ε is essential to the argument; the paper corrects an earlier proof in which a supposedly uniform constant actually depended on the chosen iterate block.

Ingredient I: resolve the map and keep the sign

Choose very ample divisors $D_1, \dots, D_r$ forming a basis of $N^1(X)_{\mathbf{R}}$ and an ample divisor $H \equiv \sum_i c_i D_i$ with $c_i \geq 0$ and $H \pm D_i$ ample. Resolve indeterminacy by $p : Y \rightarrow X$, with $g = f \circ p : Y \rightarrow X$. The negativity lemma gives effective exceptional divisors

$$Z_i = p^* p_* g^* D_i - g^* D_i. \quad (2)$$

One also chooses an ample H' on Y for which $p^* H - H'$ is effective and exceptional. Along a well-defined orbit, $p^{-1} f^j(P)$ avoids these exceptional supports. Heights of (2) may therefore be normalized to be nonnegative there, and they enter the upper estimate with a minus sign. This is the geometric reason indeterminacy does not create an uncontrolled positive error.

Ingredient II: turn numerical errors into a recurrence

Complete $g^* D_1, \dots, g^* D_r$ to a basis of $N^1(Y)_{\mathbf{R}}$. If A and B are the change-of-basis matrices used for pullback and pushforward, then $BA = M(f)$ is the matrix of f^* on $N^1(X)_{\mathbf{R}}$. The discrepancies left after these numerical identities are divisors numerically equivalent to zero. Put $\mathbf{c} = (c_1, \dots, c_r)$ and let $\|\cdot\|$ be the maximum absolute coordinate norm, for vectors and matrices. Choose associated heights with

$$h_H \geq 1, \quad h_{D_i} \geq |h_{D_i}| \ (1 \leq i \leq r), \quad h_{H'} \geq 1.$$

The numerically trivial error heights then satisfy the square-root estimate

$$|h_E(Q)| \ll \sqrt{h_{H'}(Q)}. \quad (3)$$

Combining (2), (3), and the ample comparison on Y yields, for

$$R = \max\{1, r^2 \|\mathbf{c}\| \|M(f)\|\}, \quad a_n = \frac{h_H(f^n(P))}{R^n},$$

a uniform recurrence of the shape

$$a_n \leq a_{n-1} + C_1 \sqrt{a_{n-1}} + C_2. \quad (4)$$

Lemma A.1 proves from (4) that $a_n \ll n^2 a_0$, hence

$$h_H(f^n(P)) \leq K n^2 R^n h_H(P). \quad (5)$$

The square-root error is the delicate arithmetic input: bounded errors would be easy, while a linear error could change the exponential rate. These normalizations make $a_n > 0$ and $a_0 \geq 1$, exactly as Lemma A.1 requires.

Ingredient III: choose one iterate, then fill the gaps

For rational maps one cannot replace $(f^k)^*$ by $(f^*)^k$. Instead use the definition of δ_f to choose k so large that

$$\frac{r^2 \|\mathbf{c}\| \|M(f^k)\|}{(\delta_f + \varepsilon)^k} < 1. \quad (6)$$

Apply (5) to the single map f^k . The ratio in (6) absorbs the factor n^2 and gives

$$h_H(f^{nk}(P)) \ll (\delta_f + \varepsilon)^{nk} h_H(P).$$

Lemma 3.3 supplies a coarse one-step bound $h_X^+(f^t(Q)) \leq C_0^t h_X^+(Q)$. Writing $m = qk + t$ with $0 \leq t < k$ therefore fills the finitely many residue classes and proves (1). Taking n th roots and then letting $\varepsilon \downarrow 0$ gives the arithmetic degree inequality.

Toy model, hardest point, and scope

For a degree- $d > 1$ morphism of $\mathbf{P}^1$, the familiar relation $h(f(P)) = d h(P) + O(1)$ telescopes to $h(f^n(P)) = d^n h(P) + O(d^n)$, so $\bar{\alpha}_f(P) \leq d = \delta_f$. The general proof preserves this mechanism after replacing a scalar d by pullback matrices and replacing bounded errors by (3). Its hardest point is uniformity: the basis, ample height, and coefficient vector are fixed before f is replaced by f^k , so the multiplier in (6) is genuinely independent of the block length.

The theorem does not prove that the arithmetic-degree limit exists, nor that it equals δ_f for a Zariski-dense orbit. The paper additionally proves that in Picard rank one, if f is algebraically stable and $\delta_f > 1$, then $h_X(f^n(P))/\delta_f^n$ converges (Proposition 1.11); the proof uses Theorem 1.4 through the more general Proposition 4.3. Theorem 1.10 gives sharper rank-one estimates. The paper also treats surjective morphisms separately and explains the positive-characteristic modification, neither of which is needed for the rational-map mechanism above.