

On semistable degenerations of Fano varieties

Konstantin Loginov

arXiv:1909.08319

<http://arxiv.org/abs/1909.08319v2>

How complicated can a Fano limit be?

Work over $\mathbf{C}$. Let $\pi : \mathcal{X} \rightarrow B \ni o$ be a projective family over a curve germ, with $\mathcal{X}$ smooth, $-K_{\mathcal{X}}$ relatively ample, and special fiber

$$F = \pi^{-1}(o) = F_1 + \cdots + F_r$$

reduced with simple normal crossings. If the fibers have dimension n , the dual complex $\mathcal{D}(F)$ has a vertex for each F_i and an m -cell for each irreducible component of an $(m+1)$ -fold intersection.

Main theorem (Loginov, Theorem A). *The dual complex is a simplex Δ^k with $k \leq n$. If $k = n$ and $n \leq 3$, the degeneration is unique and has trivial monodromy: it is obtained by successively blowing up, in one fiber of a smooth $\mathbf{P}^n$ -family, a flag*

$$\{\text{pt}\} = \mathbf{P}^0 \subset \mathbf{P}^1 \subset \cdots \subset \mathbf{P}^{n-1}. \quad (1)$$

Each of its $n+1$ components is $\mathbf{P}^n$ blown up along the shorter flag $\mathbf{P}^0 \subset \cdots \subset \mathbf{P}^{n-2}$.

Hu's construction, recalled as Proposition 3.5, shows separately that every admissible $k \leq n$ occurs for a degeneration of $\mathbf{P}^n$. Thus a Fano degeneration has no missing faces: once its components are known, every subset of them meets in exactly one stratum, until dimension forbids it. This is substantially stronger than mere contractibility of the dual complex.

Ingredient I: adjunction makes every stratum log Fano

Because F is a fiber, $F \sim_B 0$. Adjunction on a component gives

$$-K_{\mathcal{X}}|_{F_i} = -K_{F_i} - \sum_{j \neq i} F_j|_{F_i}, \quad (2)$$

which is ample. Hence each component, with the other components as boundary, is log Fano; the same argument applies to every stratum. Shokurov–Kollár connectedness applied first on $\mathcal{X}$ shows that any two F_i, F_j meet, and applied on F_i shows that their intersection is connected, hence irreducible by the snc assumption.

Theorem 3.3 iterates this idea. Suppose every m chosen components meet in a unique stratum Z_m . Connectedness for a suitable log Fano boundary on Z_{m-1} makes $Z_m \cap F_{i_{m+1}}$ nonempty; connectedness on Z_m makes it irreducible. Induction proves that every set of vertices spans exactly one face. The snc condition bounds its dimension by n , proving the simplex statement.

The elementary model is worth keeping in mind: two components give an edge, three give a filled triangle rather than a three-cycle with missing interior, and four give a filled tetrahedron. Blowing up the flag (1) realizes precisely these models in all allowed dimensions.

Ingredient II: surfaces are controlled by threefold MMP

For a family of del Pezzo surfaces, choose an ample curve $C \subset F_i$. Since $F \sim_B 0$ and F_i meets every other component,

$$C \cdot F_i = - \sum_{j \neq i} C \cdot F_j < 0. \quad (3)$$

An extremal ray in a decomposition of $[C]$ is therefore contained in F_i . There are no flipping contractions on the smooth threefold $\mathcal{X}$, so it gives a divisorial Mori contraction with exceptional divisor F_i .

Mori's list leaves types $(E1), (E2), (E3), (E5)$. Their possible gluings are then constrained by Friedman's d -semistability condition. For $D = F_i \cap F_j$ it reads

$$N_{D/F_i} \otimes N_{D/F_j} \otimes \bigotimes_{k \neq i,j} \mathcal{O}_D(F_k|_D) \simeq \mathcal{O}_D. \quad (4)$$

For surfaces this is the triple-point formula $D_{F_i}^2 + D_{F_j}^2 + n_D = 0$, where n_D is the number of triple points of F along D . Together with the simplex theorem it reduces the possibilities to Fujita's six configurations (Theorem 4.2), and all six occur. In every case, a sequence of component contractions reaches a smooth family, so all six have trivial monodromy. The maximal configuration has three components, all $\mathbb{F}_1$, each with boundary the negative section and a fiber.

Ingredient III: the maximal threefold case

Here maximality gives four components and a tetrahedral dual complex. Maeda's classification of smooth log Fano threefolds with integral boundary supplies a finite list of possible contractions for each pair $(F_i, \sum_{j \neq i} F_j|_{F_i})$. Formula (4), imposed simultaneously on all six pairwise intersections, eliminates the Picard-rank-one, $\mathbf{P}^2$ -bundle, and incompatible $\mathbf{P}^1$ -bundle cases. In the remaining case the normal bundle equations leave precisely the $\mathbb{F}_1$ flag-blow-up case below.

Consequently Proposition 6.1 identifies every component with

$$(\mathbf{P}_{\mathbb{F}_1}(\mathcal{O}_{\mathbb{F}_1}(-s-f) \oplus \mathcal{O}_{\mathbb{F}_1}), H \cup M_s \cup M_f), \quad (5)$$

Here s is the negative section and f a ruling fiber of $\mathbb{F}_1$; for the bundle projection p , H is the tautological divisor and $M_s = p^{-1}(s)$, $M_f = p^{-1}(f)$. The three boundary surfaces are $H \simeq \mathbb{F}_1$, $M_s \simeq \mathbf{P}^1 \times \mathbf{P}^1$, and $M_f \simeq \mathbb{F}_1$. Equivalently, each F_i is the blow-up of $\mathbf{P}^3$ along a flag $\{\text{pt}\} \subset \mathbf{P}^1$; the entire degeneration is toric. This is exactly the flag construction (1), which proves uniqueness and trivial monodromy for $n = 3$.

Proof roadmap and hardest point

The logical chain is

$$\text{relative Fano} \xrightarrow{\text{adjunction}} \text{log Fano strata} \xrightarrow{\text{connectedness}} \mathcal{D}(F) = \Delta^k \xrightarrow[\text{(4)}]{\text{low-dimensional MMP}} \text{unique maximal flag model}. \quad (6)$$

The simplex portion is conceptual and dimension-free. The difficult part is uniqueness: one must coordinate the classification on different components, because a locally permissible log Fano pair may have a normal bundle that makes (4) fail on a neighboring stratum. In dimension three this becomes the case analysis leading to (5).

Scope and open edge

The uniqueness assertion is proved only for maximal degenerations in dimensions at most three; outside the complete del Pezzo list, no general monodromy theorem is claimed. The paper asks whether maximal degenerations remain unique in every dimension and whether an arbitrary semistable Fano family becomes smooth after a suitable relative MMP. An explicit threefold example shows why the second question is subtle: divisorial contractions alone need not suffice, although a flip followed by contraction can reach a smooth family. The general classification of semistable Fano threefold degenerations also remains open.