Unobstructedness of deformations of Calabi-Yau varieties with a line bundle

Shizhang Li, Xuanyu Pan

arXiv:1310.7162

http://arxiv.org/abs/1310.7162v5

The deformation problem

Let X be a smooth proper variety over $\mathbf{C}$ and L a line bundle. For a local Artinian $\mathbf{C}$-algebra A, a deformation of (X, L) is a flat deformation X_A/A together with a line bundle L_A restricting to L. The pair is *unobstructed* if every such pair lifts across every small extension $A' \twoheadrightarrow A$.

The first Chern class defines the Atiyah extension

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{E}_L \longrightarrow T_X \longrightarrow 0.$$

An obstruction to lifting (X_A, L_A) lies in $H^2(X, \mathcal{E}_L)$, and its image in $H^2(X, T_X)$ is the ordinary obstruction to deforming X. Thus the problem is not solved merely by Tian–Todorov: the line bundle contributes an additional obstruction.

Main theorem (Theorem 1.2)**.** *Let X be a smooth proper complex variety whose canonical bundle is torsion, and let L be any line bundle on X. Then the deformation functor of the pair (X, L) is unobstructed.*

No ampleness is assumed. In particular, the theorem includes line bundles on abelian factors for which the classical polarized argument does not apply.

Two structural reductions

1. Finite étale covers detect the obstruction. If $f : Y \to X$ is finite étale, then $f^*\mathcal{E}_L \cong \mathcal{E}_{f^*L}$ and pullback on the relevant cohomology is injective, using the trace splitting. Proposition 4.4 shows more precisely that the obstruction class pulls back to the obstruction for $(Y_A, f_A^* L_A)$. Hence unobstructedness of the pulled back pair implies unobstructedness downstairs.

2. Products can be separated. Under the vanishings used for the Beauville–Bogomolov factors, deformations of $X \times Y$ preserve the product structure. Restricting a line bundle to the two factors sends its lifting obstruction in $H^2(\mathcal{O}_{X\times Y})$ to those on X and Y; Künneth makes this map an isomorphism in the situation of Proposition 5.2. Therefore unobstructed factor pairs give an unobstructed product pair.

The hypotheses here matter: Lemma 5.1 assumes trivial canonical bundles together with $H^0(Y, T_Y) = H^1(Y, \mathcal{O}_Y) = 0$. They identify first-order product deformations and, since the factors are already unobstructed, upgrade this to the full product deformation functor. The line-bundle obstruction then separates because the cross term in $H^2(\mathcal{O}_{X\times Y})$ vanishes.

These results reduce the theorem, after a finite étale cover, to abelian varieties, irreducible holomorphic symplectic varieties, and the paper's Calabi–Yau factors of dimension at least three.

The hard factor: an abelian variety with any line bundle

Let A be an abelian variety of dimension n and put $V = H^1_{\mathrm{dR}}(A/\mathbf{C})$ and $F = H^0(A, \Omega^1_A)$. A nearby complex structure is a point $[W \subset V]$ in the period domain inside $\mathrm{Gr}(n, V)$. For $c = c_1(L) \in \bigwedge^2 V$, the condition that c remain of type $(1,1)$ cuts out the Hodge locus

$$G_c = \{[W] : c \mapsto 0 \text{ in } \bigwedge^2 (V/W)\}.$$

3. Formal Hodge condition equals extension of the line bundle. Theorem 6.9 treats a proper formally smooth formal scheme over $\mathbf{C}[[x_1, \ldots, x_r]]$. It proves that a special-fiber line bundle extends if and only if the horizontal lift of its first Chern class lies in $F^1 H^2_{\mathrm{dR}}$. The proof identifies the Picard obstruction through logarithms on $1 + \mathfrak{m}\mathcal{O}$ and compares it with the Hodge-filtration quotient. This turns the image of $\mathrm{Def}(A, L) \to \mathrm{Def}(A)$ into the formal Hodge locus.

Concretely, the logarithm identifies $1 + \mathfrak{m}\mathcal{O}$ with $\mathfrak{m}\mathcal{O}$. Its connecting map sends L to a class $\Theta(L) \in H^2_{\mathrm{dR}}$ which is horizontal and reduces to $c_1(L)$. Projecting $\Theta(L)$ to $H^2(\mathcal{O})$ recovers the actual Picard obstruction, while degeneration of the Hodge-to-de Rham spectral sequence identifies its kernel with $F^1 H^2_{\mathrm{dR}}$. This is the two-way implication: the Hodge condition is not merely a necessary infinitesimal constraint.

4. That Hodge locus is smooth. At a point W of the Hodge locus inside the period domain, the real $(1,1)$ class c has off-diagonal block form. Write $W^\perp = (V/W)^* \subset V^*$ for the annihilator of W. If $c' : W^\perp \to W$ is one block, then

$$T_{G_c, W} = \{\phi : W \to V/W : \phi \circ c' = 0\} \cong \mathrm{Hom}(W/c'(W^\perp), V/W).$$

The rank of c' is $\frac{1}{2}\operatorname{rank}(c)$ throughout this Hodge locus, so the tangent dimension is the constant $n(n - \operatorname{rank}(c)/2)$. The Hodge locus is therefore smooth. Theorem 6.9 supplies a section of the forgetful functor over it, while the formally smooth dual abelian scheme supplies the Pic^0 directions. The resulting formal space is isomorphic to $\mathrm{Def}(A, L)$ (Theorem 7.8), proving the abelian case.

The final identification uses more than smoothness of the base. Over the formal Hodge locus there is a universal formal abelian scheme $\mathcal{X}$ carrying an extension $\mathcal{L}$ of L. Completing its dual abelian scheme along the identity adds every degree-zero twist; the Poincaré bundle on the resulting fiber product produces a formal family of pairs. The universal property of the dual abelian scheme then identifies this formally smooth family with the whole deformation functor of (A, L).

For an elliptic curve this mechanism is visible without the linear algebra: a line bundle's degree remains an integral $(1, 1)$ class under deformation, while tensoring by degree-zero bundles accounts for the Picard directions. In higher dimension, the Hodge locus may be a proper subspace; the constant-rank calculation is what prevents it from developing singular obstruction directions.

The remaining factors and the final assembly

For a Calabi–Yau factor Y of dimension at least three in the paper's sense, $H^2(Y, \mathcal{O}_Y) = 0$, so the extra line-bundle obstruction vanishes. For an irreducible holomorphic symplectic Y, a line bundle with $c_1(L) = 0$ is trivial and remains so. If $c_1(L) \neq 0$, then $H^2(Y, \mathcal{O}_Y) = \mathbf{C}$, and the cup product

$$H^1(Y, T_Y) \xrightarrow{\cup c_1(L)} H^2(Y, \mathcal{O}_Y)$$

is nonzero, hence surjective: the Beauville–Bogomolov form lets one choose a Kähler class pairing nontrivially with $c_1(L)$. Lemma 3.4 then kills the Atiyah obstruction.

Indeed, for $\dim Y = 2m$, contraction with the holomorphic volume form σ^m identifies T_Y with $\bigwedge^{2m-1} \Omega^1_Y$. The relevant composite sends σ^{m-1} to $\sigma^{m-1} c_1(L)[\omega]$. After Serre-pairing with $\bar{\sigma}^{m-1}$, the resulting top-degree integral is a nonzero multiple of $q_Y([\omega], c_1(L))$. Nondegeneracy of the Beauville–Bogomolov form therefore supplies the required nonzero cup product rather than an abstract dimension count.

Finally, Beauville–Bogomolov gives a finite étale cover of X by a product of precisely these factors. Product unobstructedness assembles the pulled-back pair, and injectivity under the finite cover descends the vanishing of every obstruction to (X, L).

The proof is over $\mathbf{C}$ and uses Hodge theory and Beauville–Bogomolov decomposition; it is not a mixed-characteristic statement. It proves formal unobstructedness of the pair, not that every deformation of X carries L: allowed deformations lie on the smooth Hodge locus. Several structural inputs are cited rather than reproved, while the paper establishes the crucial line-bundle Hodge criterion in full.