Canonical heights and the arithmetic complexity of morphisms on projective space

Shu Kawaguchi, Joseph H. Silverman

Shu Kawaguchi and Joseph H. Silverman, arXiv:0706.2166v1 (2007), 16 pages

http://arxiv.org/abs/0706.2166v1

Canonical heights as coordinates on a space of maps

Let h be the absolute logarithmic Weil height on $\mathbf{P}^N(\overline{\mathbb{Q}})$. For a morphism $\varphi : \mathbf{P}^N \to \mathbf{P}^N$ of degree $d \geq 2$, its canonical height satisfies

$$\widehat{h}_\varphi = h + O(1), \qquad \widehat{h}_\varphi \circ \varphi = d\widehat{h}_\varphi.$$

The paper measures the arithmetic separation of two maps by

$$\widehat{\delta}(\varphi, \psi) = \sup_{P \in \mathbf{P}^N(\overline{\mathbb{Q}})} \left|\widehat{h}_\varphi(P) - \widehat{h}_\psi(P)\right|.$$

Writing Rat_d^N as the coefficient projective space, let $h(\varphi)$ be the Weil height of the coefficient point of φ.

Main theorem (Kawaguchi–Silverman, Theorems 1 and 15)**.** *If $\varphi, \psi : \mathbf{P}^N \to \mathbf{P}^N$ are morphisms of degrees at least two over $\overline{\mathbb{Q}}$, then*

$$\widehat{\delta}(\varphi, \psi) - h(\psi) \ll h(\varphi) \ll \widehat{\delta}(\varphi, \psi) + h(\psi),$$

where the constants depend only on N and the two degrees. More explicitly, if $d = \deg \varphi$, the upper bound has leading coefficient $(d+1)\binom{N+d}{N}$ in front of $\widehat{\delta}(\varphi, \psi)$, up to a term controlled by N, d, and $h(\psi)$.

Thus a dynamical invariant defined by evaluating heights at every algebraic point is quantitatively comparable with the ordinary height of one point in the parameter space of maps.

Ingredient 1: replace an asymptotic height by one iterate

For $d = \deg \varphi$, introduce

$$\widehat{\Delta}_\psi(\varphi) = \sup_P \left|\frac{1}{d}\widehat{h}_\psi(\varphi(P)) - \widehat{h}_\psi(P)\right|.$$

Proposition 6 is the elementary but essential bridge

$$\frac{d}{d+1}\widehat{\Delta}_\psi(\varphi) \leq \widehat{\delta}(\varphi, \psi) \leq \frac{d}{d-1}\widehat{\Delta}_\psi(\varphi).$$

It follows by inserting $\widehat{h}_\varphi$ and using $\widehat{h}_\varphi(\varphi(P)) = d\widehat{h}_\varphi(P)$. The triangle inequality for $\widehat{\delta}$ then reduces comparison with an arbitrary ψ to comparison with the power map, whose canonical height is h.

Ingredient 2: recover every coefficient from generic values

Put $K = \binom{N+d}{N}$, the number of degree-d monomials in $N+1$ variables. Away from a fixed determinantal locus, the values of φ at K points determine all its coefficients: the evaluation matrix has rows consisting of those monomials, and its adjugate gives a multihomogeneous reconstruction formula. Proposition 11 converts that formula into

$$h(\varphi) \le d(K-1)\sum_{j=1}^{K} h(P_j) + \sum_{j=1}^{K} h(\varphi(P_j)) + O_{N,d}(1).$$

Height-zero points are Zariski dense in $\mathbf{P}^N$: one may take coordinates that are zero or roots of unity. Hence a K-tuple of such points can be chosen off the determinantal locus. The first sum vanishes, while the definition of $\widehat{\Delta}$ bounds each remaining value, giving the explicit estimate

$$h(\varphi) \le d\binom{N+d}{N}\widehat{\Delta}(\varphi) + O_{N,d}(1).$$

This interpolation step is the hard quantitative core of the proof.

Ingredient 3: the reverse inequality and the global comparison

For the reverse estimate, the universal degree-d morphism over the open parameter space of morphisms is inserted into the Call–Silverman variation theorem for canonical heights. Uniformly in the parameter it gives $\widehat{\Delta}(\varphi) \ll_{N,d} h(\varphi) + 1$. Combining the two estimates for $\widehat{\Delta}$, Proposition 6, and the triangle inequality through the power map proves the main theorem.

For the power map λ, whose canonical height is h, the unsubscripted notation $\widehat{\Delta}(\varphi)$ means $\widehat{\Delta}_\lambda(\varphi)$. As a toy family, take $\varphi_A(z) = z^d + Az^{d-1}$ with $A \in \mathbb{Z}\setminus\{0\}$ and let $|A| \to \infty$. Its coefficient height grows like $\log|A|$, while the one-step height discrepancy grows on the same scale. The example suppresses the multivariate interpolation issue but retains the mechanism by which coefficient size is detected dynamically.

Northcott consequence and scope

Fix ψ and $d \ge 2$. A bound on $\widehat{\delta}(\varphi,\psi)$ gives a bounded-height subset of Rat_d^N. Northcott's theorem therefore leaves only finitely many such φ whose fields of definition have bounded degree. The field-degree hypothesis is indispensable: bounded height alone is not finite over $\overline{\mathbb{Q}}$.

The result concerns morphisms, not rational maps with indeterminacy, and its constants depend on the ambient dimension and degrees. The final section proposes a conjugacy-invariant infimum on the dynamical moduli space, but whether that infimum is always attained is left open.