Obstructions to algebraizing topological vector bundles

Aravind Asok, Jean Fasel, Michael J. Hopkins

arXiv:1506.08111

http://arxiv.org/abs/1506.08111v2

When algebraic Chern classes are not enough

Let X be a smooth complex affine variety and X^{an} its analytic space. A topological rank-two bundle has *algebraic Chern classes* when its classes lie in the images of the cycle maps $\mathrm{cl} : \mathrm{CH}^i(X) \to H^{2i}(X^{\mathrm{an}}, \mathbb{Z})$. In dimensions at most three this condition is sufficient for algebraization. The paper identifies the first obstruction in dimension four.

Main theorem (Theorems 2 and 4)**.** *Let X be a smooth complex affine fourfold and E^{an} a rank-two complex analytic bundle with algebraic topological Chern classes c_i^{top}. Then E^{an} is algebraizable if and only if there are lifts $c_i \in \mathrm{CH}^i(X)$, with $\mathrm{cl}(c_i) = c_i^{\mathrm{top}}$, such that*

$$\mathrm{Sq}^2(c_2) + c_1 c_2 = 0 \quad \textit{in } \mathrm{CH}^3(X)/2.$$

Here Sq^2 is the integral Steenrod square on Chow groups.

Moreover, there is a smooth hypersurface $Z \subset \mathbb{P}^1 \times \mathbb{P}^3$ of bidegree $(3, 4)$ such that, for $X = (\mathbb{P}^1 \times \mathbb{P}^3) \setminus Z$, the cycle map is injective in degrees $i \leq 2$, yet X^{an} carries a rank-two topological bundle with algebraic Chern classes that is not algebraizable.

The existential choice of the Chow lifts is essential: even when $c_1^{\mathrm{top}} = 0$, a lift c_1 need not vanish if the degree-one cycle map has a kernel. Products with affine space propagate the counterexample to every dimension at least four.

Ingredient 1: topological bundles are classified by two classes

The analytic space of a smooth affine fourfold has the homotopy type of a CW complex of dimension at most four. The universal map

$$(c_1^{\mathrm{top}}, c_2^{\mathrm{top}}) : B\mathrm{GL}_2(\mathbb{C}) \longrightarrow K(\mathbb{Z}, 2) \times K(\mathbb{Z}, 4)$$

is a 4-equivalence. Ordinary obstruction theory therefore gives a bijection from rank-two topological bundles to $H^2(X^{\mathrm{an}}, \mathbb{Z}) \times H^4(X^{\mathrm{an}}, \mathbb{Z})$. Consequently a pair of topological Chern classes exists uniquely; the real question is whether it lifts through algebraic geometry.

Ingredient 2: the motivic primary obstruction

Morel's representability theorem identifies algebraic rank-two bundles on smooth affine X with $[X, B\mathrm{GL}_2]_{\mathbb{A}^1}$. The paper studies the homotopy fibre F_2 of the motivic Chern map

$$B\mathrm{GL}_2 \longrightarrow K(\mathbf{K}_1^M, 1) \times K(\mathbf{K}_2^M, 2).$$

Its relevant first homotopy sheaf is $\pi_2^{\mathbb{A}^1}(F_2) \simeq \mathbf{I}^3$, so fixing the determinant line bundle L places the primary lifting obstruction in $H^3(X, \mathbf{I}^3(L))$.

The filtration $0 \to \mathbf{I}^{j+1}(L) \to \mathbf{I}^j(L) \to \mathbf{K}_j^M/2 \to 0$, together with the Milnor conjecture and Totaro's calculation, sends its visible quotient to

$$c_2 \longmapsto \mathrm{Sq}^2(c_2) + c_1(L)c_2.$$

For a smooth affine fourfold over an algebraically closed field, $H^3(X, \mathbf{I}^4(L))$ and $H^4(X, \mathbf{I}^4(L))$ vanish. Hence this quotient is the entire primary obstruction, not merely a necessary shadow of it.

Ingredient 3: no hidden higher obstruction remains

Once the primary class vanishes, the secondary obstruction lies in degree four. The proof resolves the relevant homotopy sheaf into Grothendieck–Witt, Milnor K-theory, and powers of the fundamental ideal. Its degree-four cohomology vanishes: the Gersten term for the Grothendieck–Witt part is zero, $\mathrm{CH}^4(X)$ is divisible so its quotient modulo 12 vanishes, and $\mathbf{I}^5$ restricts trivially. Nisnevich cohomological dimension four excludes any later obstruction. This is the hardest abstract step: it upgrades one computable class into a necessary *and sufficient* criterion, after which $\mathbb{A}^1$-representability turns the lift into an algebraic bundle.

The concrete $(3, 4)$ counterexample

Let ξ, μ be the hyperplane classes of $\mathbb{P}^1 \times \mathbb{P}^3$. The paper chooses a smooth $(3, 4)$-hypersurface Z, defined over $\mathbb{Q}$ with a specified singular specialization, and sets $X = (\mathbb{P}^1 \times \mathbb{P}^3) \setminus Z$. Lefschetz and Gysin arguments make the cycle maps injective for $i \leq 2$. If $\psi \in \mathrm{CH}^2(X)$ is the restriction of $\xi\mu$, the Cartan formula gives

$$\mathrm{Sq}^2(\psi) = \xi\mu^2\big|_X .$$

The coprimality of 3 and 4 keeps ψ nonzero. Totaro's specialization argument says every curve in Z has even degree over $\mathbb{P}^1$; hence $\xi\mu^2$ is not in the Gysin image modulo 2, so the displayed square is nonzero in $\mathrm{CH}^3(X)/2$.

Topologically there is therefore a unique rank-two bundle with Chern classes $(0, \mathrm{cl}(\psi))$. Both classes are algebraic, and injectivity makes ψ the unique second lift, but its Steenrod obstruction is nonzero. This produces the promised non-algebraizable bundle. The example is tied to failure of the integral Hodge conjecture for Z; the paper does not claim that every fourfold or every algebraic pair of Chern classes is obstructed.