Landen transforms as families of (commuting) rational self-maps of projective space

Michael Joyce, Shu Kawaguchi, Joseph H. Silverman

arXiv:1308.5355

http://arxiv.org/abs/1308.5355v1

From a coefficient filter to projective dynamics

Fix integers $m \geq 1$ and $0 \leq k < m$. If K is a field in which $m \neq 0$ and ζ_m is a primitive mth root of unity, the (m,k)-Landen transform of $\varphi \in K(z)$ is characterized by

$$\mathcal{F}_{m,k}(\varphi)(w^m) = \frac{1}{mw^k} \sum_{t=0}^{m-1} \zeta_m^{-kt} \varphi(\zeta_m^t w).$$

The right side is invariant under $w \mapsto \zeta_m w$, descends to $K(w^m)$, and is independent of the chosen primitive root. On a Laurent series $\sum_n c_n z^n$ it simply retains the congruence class $n \equiv k \pmod m$ and sends $c_{mj+k} z^{mj+k}$ to $c_{mj+k} z^j$.

Write a degree-d rational function as

$$\varphi_{a,b}(z) = \frac{F_a(z)}{F_b(z)}, \qquad [a,b] = [a_0, \ldots, a_d, b_0, \ldots, b_d] \in \mathbf{P}^{2d+1}.$$

Thus Rat_d is the complement of the resultant hypersurface in $\mathbf{P}^{2d+1}$.

Main theorem (Theorem 2 and Corollary 4)**.** *For every $d \geq 1$ there is a unique rational map over $\mathbf{Z}$*

$$\mathcal{R}_{d,m,k} : \mathbf{P}^{2d+1}_{\mathbf{Z}} \dashrightarrow \mathbf{P}^{2d+1}_{\mathbf{Z}}$$

which agrees with $\mathcal{F}_{m,k}$ wherever the latter preserves degree d. Its indeterminacy locus is exactly $\{[a,b] : b = 0\} \cong \mathbf{P}^d$, and it maps the complement of this linear space morphically to itself. The map $\mathcal{R}_{d,m,0}$ is dominant of algebraic degree m. If $1 \leq k < m$, its image is the hyperplane $\{a_0 = 0\} \cong \mathbf{P}^{2d}$; for every k, the induced rational self-map of that hyperplane is dominant of algebraic degree m.

For fixed d, the maps $\{\mathcal{R}_{d,m,0} : m \geq 1\}$ commute. After dehomogenizing by $b_0 \neq 0$, they give commuting dominant polynomial self-maps of $\mathbf{A}^{2d+1}$, and likewise of the invariant $\mathbf{A}^{2d}$ cut out by $a_0 = 0$.

The integral-coordinate mechanism

1. Clear all twisted denominators at once. The decisive construction is Proposition 16. It produces

$$G_{a,b,m,k}(z) \in \mathbf{Z}[a,b,z], \qquad H_{b,m}(z) \in \mathbf{Z}[b,z]$$

such that

$$\mathcal{F}_{m,k}(\varphi_{a,b})(z) = \frac{G_{a,b,m,k}(z)}{H_{b,m}(z)}, \quad H_{b,m}(w^m) = \prod_{t=0}^{m-1} F_b(\zeta_m^t w).$$

The numerator is the analogous weighted sum after multiplying by all but one denominator. Root-of-unity invariance and Galois invariance put both polynomials over $\mathbf{Z}$; a Laurent-series calculation supplies the otherwise non-obvious divisibility by m. Their z-coefficients are homogeneous of total degree m, with G linear in a and of degree $m-1$ in b.

If $F_b(z) = b_0 \prod_i (z - \beta_i)$, then, up to the recorded sign,

$$H_{b,m}(z) = b_0^m \prod_i (z - \beta_i^m).$$

The resultant formula compares $\mathrm{Res}(G,H)$ with $\mathrm{Res}(F_a, F_b)$ and the discriminant quotient $\mathrm{Disc}(H)/\mathrm{Disc}(F_b)$. In the homogeneous degree-d convention, it is

$$\mathrm{Res}(G,H) = (-1)^{d(m-k+1)} b_0^{m-1} b_d^{m-1-k} \mathrm{Res}(F_a, F_b) \frac{\mathrm{Disc}(H)}{\mathrm{Disc}(F_b)}.$$

The quotient on the right is itself in $\mathbf{Z}[b]$; its root formula shows exactly when two denominator roots become identified after taking mth powers. This is the arithmetic input that prevents the coefficient construction from being valid only over a splitting field. Consequently degree is preserved on a dense open set: it suffices that the denominator roots are nonzero and distinct roots do not acquire the same mth power.

2. Read the compactified map from coefficients. Taking the $d+1$ coefficients of G followed by those of H defines $\mathcal{R}_{d,m,k}$. All coordinates vanish precisely when $b=0$: if H vanishes identically, one factor $F_b(\zeta_m^t w)$ does, hence $b=0$. Since this base locus has dimension d, the coordinate forms cannot have a common divisor; therefore the projective map really has algebraic degree m. The leading numerator coefficient is proportional to $a_0 b_0^{m-1}$ for $k=0$ and is zero for $k>0$, which explains the hyperplane image.

3. Prove dominance without losing bad characteristics. Because G is linear in a and H is independent of a, the Jacobian is block triangular. A weight argument isolates a coefficient monomial with coefficient ± 1 in its diagonal block. This makes the Jacobian nonzero when m is invertible. If a prime p divides m, the maps are first factored using the composition law and reduced to $m=p$. Modulo p, the denominator coordinates become $(b_0^p,\dots,b_d^p)$; taking pth roots of a generic target and inverting the surviving a-linear block gives a preimage. Thus the reduction is generically surjective even though it is inseparable.

More explicitly, for $k=0$ the two diagonal blocks satisfy $D_{d,m} = mA_{d,m,0}$. A filtration by the ideals $(b_0,\dots,b_{j-1})$ makes $A_{d,m,0}$ triangular to leading order and exhibits the monomial $(b_0\cdots b_d)^{m-1}$ with unit coefficient in its determinant. This is why the same calculation both proves generic separability away from primes dividing m and survives modulo those primes after the differential itself degenerates. For $k>0$ one deletes the first row and column, restricts to $a_0=0$, and uses the parallel unit-coefficient monomial of Lemma 25.

Commutation, toy case, and scope

The Laurent-series filter also gives Proposition 14,

$$\mathcal{F}_{m,k}\circ\mathcal{F}_{n,\ell} = \mathcal{F}_{mn,kn+\ell}.$$

Since the coefficient maps agree on a dense open set, the same identity holds for the $\mathcal{R}$'s. Setting $k=\ell=0$ yields commutation.

For $d=m=2$, the theorem becomes concrete:

$$\mathcal{R}_{2,2,0} = [b_0a_0,\ b_2a_0 - b_1a_1 + b_0a_2,\ b_2a_2, b_0^2,\ 2b_2b_0 - b_1^2,\ b_2^2].$$

This toy case displays all structural features: quadratic coordinates, the base plane $b=0$, and the denominator roots being squared. What it hides is the uniform integral argument and the inseparable fibers.

There is also a clean factor of the construction which forgets the numerator. If

$$h_m : \mathbf{P}^d \longrightarrow \mathbf{P}^d, \qquad b \longmapsto [H_{m,0}(b),\dots,H_{m,d}(b)],$$

then the same common-zero argument shows that h_m is a morphism. On the chart $b_0=1$, let $\bar\sigma_d$ identify the symmetric quotient of ordered roots with coefficient space and let $\pi_m(u_1,\dots,u_d) = (u_1^m,\dots,u_d^m)$. Proposition 28 gives

$$h_m = \bar\sigma_d \circ \pi_m \circ \bar\sigma_d^{-1}.$$

Thus the denominator dynamics is coordinatewise powering after passing to unordered roots. The same description applies on each boundary stratum obtained by letting the initial coefficients of F_b vanish; this explains why no additional base point appears at infinity.

The result concerns maps on a parameter space, not iteration of the original φ. The extension over $\mathbf{Z}$ is a rational map, not a morphism on all projective space. For $k>0$ the full image drops to a hyperplane, and the paper only outlines the parallel Jacobian calculation for its dominance (including an omitted technical lemma). The denominator map itself is especially transparent: on the affine root quotient it is conjugate to coordinatewise mth power. In characteristic zero the generic separable degree of $\mathcal{R}_{d,m,0}$ is m^d; in characteristic p, writing $m=p^e n$ with $p\nmid n$, the paper records separable degree n^d and inseparable degree p^{ed}.