Non-recurrent rational maps with disconnected Julia set

Yan Gao, Lele Xu, Luxian Yang

Yan Gao, Lele Xu, and Luxian Yang, arXiv:2509.06645v1 (2025), 13 pages

http://arxiv.org/abs/2509.06645v1

Critical recurrence controls disconnected Julia sets

Let $f : \widehat{\mathbf{C}} \to \widehat{\mathbf{C}}$ be a rational map of degree at least two with disconnected Julia set J_f. A Julia component is *wandering* if its forward images never repeat. A compact connected set is of full, annular, or complex type according as its complement has one, two, or at least three components. A Julia component is *exposed* if it meets the boundary of a Fatou component. Finally, a compact connected set E is non-recurrent if some neighborhood D of E satisfies $f^n(E) \cap D = \varnothing$ for every $n \geq 1$.

Main theorem (Gao–Xu–Yang, Theorems 1.1 and 1.2)**.** *Suppose every wandering exposed Julia component containing a critical point is non-recurrent. Then every wandering exposed Julia component is a singleton.*

Under the stronger hypothesis that every wandering Julia component containing a critical value is non-recurrent, J_f has only finitely many periodic components of complex type.

The first conclusion extends the Branner–Hubbard shrinking phenomenon from polynomials to a class of geometrically infinite rational maps. The second rules out infinitely many topologically complicated periodic pieces. Notice the deliberate asymmetry: Theorem 1.1 concerns critical *points* on exposed wandering components, whereas Theorem 1.2 assumes non-recurrence for components containing critical *values*.

Ingredient 1: boundary components and exceptional transitions

After passing to an iterate, an exposed wandering component can be studied on the boundary of a fixed multiply connected Fatou component U. Put $\infty \in U$, let $\mathcal{E}$ be the components of ∂U, and let $\widehat{E}$ denote E with all bounded complementary components filled in. Iteration induces $\sigma_f(E) = f(E)$ on $\mathcal{E}$. Call E essential when $\widehat{E} \cap f^{-1}(U) \neq \varnothing$, and call it critical when its filling $\widehat{E}$ contains a critical point.

Lemma 2.1 supplies the finite combinatorial skeleton. There are at most $\deg(f) - 1$ essential elements; each is non-full and critical; and every wandering $E \in \mathcal{E}$ is eventually full. The authors collect the preperiodic critical elements (and the parabolic boundary element, when present) in a finite invariant set $\mathcal{E}_1$. A second set $\mathcal{E}_2$ consists of the finite initial segments of non-full wandering critical orbits, up to their first full image. Every essential element lies in $\mathcal{E}_1 \cup \mathcal{E}_2$. Thus all failures of clean pullback occur along explicitly isolated boundary orbits.

Ingredient 2: bounded-degree puzzle pullbacks shrink

Choose a small Jordan domain $U_0 \Subset U$ around the attracting fixed point, or the analogous parabolic petal, and let U_N be a sufficiently deep pullback. The components of $\widehat{\mathbf{C}} \setminus U_{N+n}$ are depth-n puzzle pieces P_n; for $E \in \mathcal{E}$,

$$\bigcap_{n \geq 0} P_n(E) = \widehat{E}.$$

Away from essential elements, $P_{n+1}(E)$ is an entire component of the pullback of $P_n(\sigma_f E)$; away from critical elements the map between them is a homeomorphism (Propositions 2.1–2.2).

Ingredient 2 continued: degree control and shrinking

Here is the decisive use of non-recurrence. Along suitable escape times n_k, the pieces $f^j(P_{n_k}(E))$, $0 \le j \le n_k$, meet each wandering critical boundary element at most once and no preperiodic critical one. Consequently

$$\deg\Big(f^{n_k} : P_{n_k}(E) \longrightarrow P_0(\sigma_f^{n_k} E)\Big) \le d^{m_0},$$

where $d = \deg(f)$ and m_0 is the number of wandering critical elements of $\mathcal{E}$. The Shrinking Lemma for inverse images of a Jordan disk with uniformly bounded degree then forces $\operatorname{diam} P_{n_k}(E) \to 0$. This is the analytic engine: bounded ramification converts combinatorial escape into a singleton intersection.

Proof roadmap for the singleton theorem

Lemma 2.5 constructs the escape times unless the orbit accumulates on $\mathcal{E}_2$. If $\mathcal{E}_2$ were nonempty, critical non-recurrence lets one choose an element of it whose orbit accumulates on no element of $\mathcal{E}_2$; the escape-time lemma and shrinking would make that element a singleton, contradicting its defining non-full initial segment. Hence $\mathcal{E}_2 = \varnothing$. Every wandering boundary component therefore has the required escape times and is a singleton; the original exposed Julia component lies in its filling and is the same singleton.

A useful toy picture is a nested sequence of disks pulled back univalently except at finitely many critical encounters. If no critical orbit returns to the same small puzzle, the total pullback degree stays bounded, so the disks shrink. The rational-map proof must additionally handle multiply connected Fatou domains, essential boundary components, and the parabolic construction.

Ingredient 3: infinitely many periodic pieces force recurrence

For Theorem 1.2, conjugate so that $f(\infty) = \infty$, let V_f be the critical-value set, and let $\mathcal{C}$ be the periodic complex-type components. Those meeting $V_f \cup \{\infty\}$, or separating at least two bounded regions that meet V_f, form a finite set $\mathcal{S}$. Outside their orbits, the remaining components split into classes $\mathcal{A}$ and $\mathcal{D}$ according as exactly one, or no, bounded complementary component meets the critical values. A first-return map on $\mathcal{A}$ associates to K a distinguished pullback region $U(K)$. The subset

$$\mathcal{B} = \{K \in \mathcal{A} : U(K) \neq V(K)\}$$

records a genuine change of the critical-value side $V(K)$.

If $\mathcal{C}$ were infinite, its periods would be unbounded: for each fixed period, McMullen's quasiconformal model places a periodic point in every such component, and f has only finitely many periodic points of that period. Large-period components must enter $\mathcal{A}$, then $\mathcal{B}$; hence $\mathcal{B}$ is infinite (Lemmas 3.3–3.5 and Proposition 3.1).

Two elements of $\mathcal{B}$ are equivalent when their regions contain the same subset of the finite critical-value set. There are therefore finitely many equivalence classes, each totally ordered by nesting. An infinite class $\mathcal{B}_i$ yields

$$E_i = \bigcap_{K \in \mathcal{B}_i} V(K),$$

a wandering Julia component containing critical values. The first-return map sends an infinite nested subsequence to another infinite class and shows that the orbit of E_i accumulates on the corresponding E_j. Iterating this relation among finitely many classes creates a directed cycle, so some E_i is recurrent—the required contradiction.

The conclusions concern exposed wandering components and periodic complex-type components only; they neither classify all disconnected Julia sets nor remove the stated non-recurrence hypotheses. The main difficulty is precisely the rational-map feature absent for polynomials: a puzzle piece need not map onto a puzzle piece until the essential transitions have been isolated.