

Arithmetic Degrees are Cohomological Lyapunov Multipliers

Jiarui Song, Junyi Xie, She Yang

arXiv:2507.17643

<http://arxiv.org/abs/2507.17643v1>

Discrete growth rates for dense orbits

Let X be a geometrically connected normal projective variety over a characteristic-zero field, let $f : X \rightarrow X$ be a surjective endomorphism, and let x have Zariski-dense forward orbit. After spreading the data to a finitely generated field $K/\mathbb{Q}$, an ample line bundle L and a Moriwaki height define

$$\alpha_f(x) = \lim_{n \rightarrow \infty} h_L(f^n(x))^{1/n}.$$

(Over a number field this is the usual Weil-height definition.) If $d = \dim X$, put $\lambda_0(f) = 1$ and let $\lambda_i(f)$ be the i th dynamical degree. The ratios

$$\mu_i(f) = \frac{\lambda_i(f)}{\lambda_{i-1}(f)}, \quad 1 \leq i \leq d, \quad (1)$$

are the *cohomological Lyapunov multipliers*; log-concavity gives $\mu_1 \geq \dots \geq \mu_d > 0$.

Main theorem (Song–Xie–Yang, Theorem 1.2). *Under the hypotheses above,*

$$\alpha_f(x) \in \{\mu_1(f), \dots, \mu_d(f)\} \cap \mathbb{R}_{\geq 1}. \quad (2)$$

The arithmetic degree was already known to be the modulus of an eigenvalue of f^* on $N^1(X)_{\mathbb{R}}$. Formula (2) replaces that generally larger spectral list by d intrinsic positive numbers. In particular, if $\lambda_1(f) > \lambda_2(f)$, then $\mu_1 = \lambda_1$ while every later $\mu_i < 1$; hence (2) proves the Kawaguchi–Silverman prediction $\alpha_f(x) = \lambda_1(f)$ in this case.

The Albanese threshold and equivariant lifting

Let $A = \text{Alb}(X)$ and let $g : A \rightarrow A$ be induced by f . Density of the orbit makes the Albanese morphism $\pi : X \rightarrow A$ surjective and the orbit of $\pi(x)$ dense. Arithmetic degrees on abelian varieties give

$$\alpha_f(x) \geq \alpha_g(\pi(x)) = \lambda_1(g). \quad (3)$$

Here, as in the paper, $\lambda_1(g)$ is defined to be 0 when A is trivial. This convention also applies below. If equality holds, the relative dynamical-degree formula already identifies $\lambda_1(g)$ with one of the multipliers of f .

The harder case is $\alpha_f(x) > \lambda_1(g)$. The exact sequence

$$0 \longrightarrow \text{Pic}^0(X)_{\mathbb{R}} \longrightarrow \text{Pic}(X)_{\mathbb{R}} \longrightarrow N^1(X)_{\mathbb{R}} \longrightarrow 0 \quad (4)$$

contains the obstruction to turning a numerical eigenclass into a genuine line-bundle class. Since $\text{Pic}^0(X) \simeq A^{\vee}$, Proposition 2.5 bounds the spectral radius of f^* on $\text{Pic}^0(X)$ by $\sqrt{\lambda_1(g)}$. Lemma 3.1 therefore lifts every generalized μ -eigenspace with $\mu > \sqrt{\lambda_1(g)}$ through (4), equivariantly for f^* . Concretely, a numerical Jordan chain is corrected by a convergent series whose error decays like a polynomial times $(\rho(f^*| \text{Pic}^0)/\mu)^n$.

A big divisor detects a forbidden gap

Set $\mu_{d+1}(f) = 0$. Assume for contradiction that $\mu_{\ell} > \alpha_f(x) > \mu_{\ell+1}$ for some $\ell \in \{1, \dots, d\}$. Theorem 2.3 supplies a big numerical class $B = M_1 + M_2$, where M_1 lies in the first ℓ generalized eigenspaces and M_2 in the remaining ones; when $\ell = d$, the latter sum is empty and $M_2 = 0$. The lifted high part has a vector-valued canonical height. In Jordan coordinates its value along the orbit is an exponential polynomial

$$\widehat{h}_{M_1}(f^n(x)) = \sum_{1 \leq i \leq \ell} \sum_k c_{i,k} n^k \mu_i^n. \quad (5)$$

For an arbitrary lift of M_2 , Corollary 3.3 bounds its height by a polynomial times ρ_1^n , with

$$\rho_1 \leq \max\{\mu_{\ell+1}, \sqrt{\lambda_1(g)}\} < \alpha_f(x). \quad (6)$$

Lemma 2.8 is the bridge back to the orbit: for a big divisor lifting B , there is a subsequence n_j on which its height is unbounded and its n_j th-root growth is exactly $\alpha_f(x)$. If all coefficients in (5) vanish, (6) makes this growth at most $\max\{\rho_1, 1\}$, contradicting either $\rho_1 < \alpha_f(x)$ or the required unboundedness when $\alpha_f(x) = 1$. If some coefficient survives, the dominant term along that subsequence forces growth at least one of $\mu_1, \dots, \mu_{\ell}$, strictly larger than $\alpha_f(x)$. Thus no gap between consecutive multipliers can contain the arithmetic degree, proving (2).

Product model, application, and scope

For $X = \mathbb{P}^1 \times \mathbb{P}^1$ and $f = (z \mapsto z^a, w \mapsto w^b)$ with $a \geq b \geq 2$, one has $(\lambda_0, \lambda_1, \lambda_2) = (1, a, ab)$ and hence $(\mu_1, \mu_2) = (a, b)$. The two possible exponential scales in (5) are visible directly. A point whose first coordinate is preperiodic may grow at rate b , but its product orbit is not dense; a dense orbit has rate a . This illustrates both the spectral quantization and the role of density.

Theorem 1.2 does not by itself prove $\alpha_f(x) = \lambda_1(f)$ in all cases. It also assumes an everywhere-defined surjective endomorphism, not a general dominant rational map. Corollary 1.5 considers projective varieties X, Y over $\mathbb{C}$, surjective endomorphisms f, g , and points x, y whose individual orbits are Zariski dense. If the two multiplier spectra have no common element in $\mathbb{R}_{\geq 1}$, and if a positive-dimensional irreducible $V \subset X \times Y$ has both projections generically finite onto their images, then V cannot contain a dense set of synchronous orbit points $(f^n(x), g^n(y))$. Indeed, Theorem 1.2 puts the two projected height growth rates in the disjoint spectra, making them incompatible on V .