

Algebraizability of Vector Bundles over Real Algebraic Varieties

Hanqi Wang

arXiv:2607.07856

<http://arxiv.org/abs/2607.07856v1>

When is a topological bundle algebraic?

Let X be a smooth affine real algebraic variety and write $X(\mathbb{R})$ for its Euclidean real locus. Following the paper's convention, “real algebraic” permits localizing the structure sheaf at functions nonvanishing on $X(\mathbb{R})$, so a bundle need only be defined on an algebraic neighbourhood of that locus. A topological real vector bundle V on $X(\mathbb{R})$ is *algebraizable* if it is topologically isomorphic to the real locus of such an algebraic bundle.

The cycle-class map

$$\mathrm{cl}: \mathrm{CH}^i(X) \longrightarrow H^i(X(\mathbb{R}); \mathbb{Z}/2)$$

defines the subgroup H_{alg}^i . Algebraic bundles necessarily have algebraic Stiefel–Whitney classes; the paper proves a sharp converse in low dimension.

Main theorem (Wang, Theorem 3.2.4). *If X has dimension at most 3, then a topological real vector bundle V on $X(\mathbb{R})$ is algebraizable if and only if*

$$w_1(V) \in H_{\mathrm{alg}}^1(X; \mathbb{Z}/2), \quad w_2(V) \in H_{\mathrm{alg}}^2(X; \mathbb{Z}/2).$$

No compactness hypothesis on $X(\mathbb{R})$ occurs here. The theorem turns an existence problem for algebraic transition functions into the detection of only two topological classes.

Ingredient I: replace bundles by motivic maps

For smooth affine X , representability identifies algebraic rank- n bundles with $\mathbb{A}^1$ -homotopy classes

$$\mathrm{Vect}_n(X) \simeq [X, B\mathrm{GL}_n]_{\mathbb{A}^1}.$$

The Nisnevich cohomological dimension of X is at most $\dim X$. Consequently a Postnikov tower for $B\mathrm{GL}_n$ has no obstruction above degree 3 when $\dim X \leq 3$. The proof therefore asks whether algebraic lifts of the first low homotopy invariants can be chosen to realize the topological characteristic classes of V .

Ingredient II: rank two and the twisted Euler class

On a 3-complex, a rank-two real bundle is classified by w_1 and its Euler class

$$e_{w_1}(V) \in H^2(X(\mathbb{R}); \mathbb{Z}_{w_1}),$$

with coefficients twisted by the orientation local system. The second Postnikov stage of $B\mathrm{GL}_2$ is governed by Milnor–Witt K -theory, and Proposition 3.1.2 says that V algebraizes precisely when w_1 and this twisted Euler class lift algebraically.

Theorem 3.1.3 is the decisive comparison: for an algebraic line bundle L , the image of the twisted integral cycle-class map is exactly the inverse image of $H_{\mathrm{alg}}^2(X; \mathbb{Z}/2)$ under reduction mod 2. Since $e_{w_1}(V) \bmod 2 = w_2(V)$, algebraicity of w_1, w_2 supplies the required motivic lift and proves the rank-two case.

Ingredient III: higher rank collapses to rank three

A rank- $r \geq 3$ bundle on a space of dimension at most 3 splits off a trivial summand of rank $r - 3$. Rank-three bundles are topologically classified by (w_1, w_2) ; Wu's formula determines w_3 from them. Motivically, the map

$$(c_1, c_2): BGL_3 \longrightarrow K(\mathbf{K}_1^M, 1) \times K(\mathbf{K}_2^M, 2)$$

is an isomorphism on the first two $\mathbb{A}^1$ -homotopy sheaves and a surjection on the third. Cohomological dimension then makes the induced map on $[X, -]_{\mathbb{A}^1}$ surjective. Algebraic preimages of w_1, w_2 therefore lift to an algebraic rank-three bundle, which completes the proof.

Why dimension four is different

Assume now that $\dim X = 4$ and $X(\mathbb{R})$ is compact. Stable orientable real bundles are classified topologically by (w_2, w_4, p_1) , but their algebraic lifts have one remaining fourth-degree Postnikov obstruction. For the homotopy fibre F of (c_2, c_4, p_1) , the paper computes

$$H^4(X, \pi_3^{\mathbb{A}^1} F) \simeq H^4(X(\mathbb{R}); \mathbb{Z}) \oplus \operatorname{coker} \left(\operatorname{CH}^3(X) \xrightarrow{(-) \smile w_1(X)} \operatorname{CH}^4(X) \right). \quad (1)$$

The difficult step (Theorem 4.1.6) identifies the boundary map in (1) with cup product by $w_1(X)$ using contracted homotopy sheaves and Rost–Schmid complexes.

Theorem 4.1.8 packages the obstruction componentwise. There are maps $k_i: \operatorname{CH}^2(X) \rightarrow H^4(M_i; \mathbb{Z})$ on orientable components and $k'_j: \operatorname{CH}^2(X) \rightarrow H^4(N_j; \mathbb{Z}/2)$ on nonorientable ones. A bundle algebraizes exactly when some $c_1 \in \operatorname{CH}^1(X)$, $c_2 \in \operatorname{CH}^2(X)$, and $\alpha \in H_{\text{alg}}^3(X; \mathbb{Z}/2)$ satisfy

$$\begin{aligned} \operatorname{cl}(c_1) &= w_1(V), & \operatorname{cl}(c_2) &= w_2(V), \\ p_1(V|_{M_i}) &= k_i(c_2 + c_1^2), & (w_4 + w_1 w_3)|_{N_j} &= \alpha|_{N_j} \smile w_1(N_j) + k'_j(c_2 + c_1^2). \end{aligned} \quad (2)$$

Thus algebraic characteristic classes cease to be sufficient in dimension four: their possible lifts must also satisfy (2).

A boundary model and scope

For $X = Y \times S^1$ with Y a smooth affine real threefold whose real locus is compact and orientable, the Chow-group calculation in Lemma 4.2.1 kills the integral map k . Theorem 4.2.2 reduces (2) to the transparent criterion

$$V \text{ algebraizable} \iff w_*(V) \in H_{\text{alg}}^*(X; \mathbb{Z}/2) \text{ and } p_1(V) = 0.$$

This model isolates exactly what first appears in dimension four: a genuine Pontryagin obstruction beyond mod-2 algebraicity.

The other principal application computes the reduced algebraic group $\tilde{K}_0(X)$. With

$$G = 1 \oplus H_{\text{alg}}^1(X; \mathbb{Z}/2) \oplus \operatorname{CH}^2(X), \quad H = w_1(X) \smile H_{\text{alg}}^3(X; \mathbb{Z}/2),$$

Proposition 4.2.3 gives $0 \rightarrow H \rightarrow \tilde{K}_0(X) \rightarrow G \rightarrow 0$. If $X(\mathbb{R})$ is orientable, write $\operatorname{CH}^2(X) = \mathbb{Z}^t \oplus (\mathbb{Z}/2)^s$, set $b_1 = \dim_{\mathbb{Z}/2} H_{\text{alg}}^1$ and $\delta = \dim_{\mathbb{Z}/2} \{v \in H_{\text{alg}}^1 : v^2 = 0\}$; Theorem 4.2.5 yields

$$\tilde{K}_0(X) \simeq \mathbb{Z}^t \oplus (\mathbb{Z}/2)^{s-b_1+2\delta} \oplus (\mathbb{Z}/4)^{b_1-\delta}. \quad (3)$$

The compactness and stable-algebraization arguments belong to the fourfold theory, not to the dimension- ≤ 3 theorem; the paper treats irreducible X and bundles of constant rank under its localized real-algebraic convention.