A stacky approach to identifying the semistable locus of bundles

Dario Weissmann, Xucheng Zhang

arXiv:2302.09245; DOI: 10.14231/AG-2025-008

http://arxiv.org/abs/2302.09245v3

Can the moduli property detect semistability?

Let C be a smooth projective connected curve over an algebraically closed field k, let G be a connected reductive group, and write Bun_G for the stack of principal G-bundles on C. Classical semistability is sufficient for constructing a moduli space. The paper asks whether the existence and geometry of that space conversely recognize the semistable locus.

Main theorem (Weissmann–Zhang, Theorems 2.1, 4.1, and 5.5)**.** *The semistable locus* $\mathrm{Bun}_G^{\mathrm{ss}}$ *is the unique maximal open substack of* Bun_G *admitting a* schematic adequate *moduli space.*

If $g(C) > 1$*, the semistable locus* $\mathrm{Bun}_2^{d,\mathrm{ss}}$ *is likewise the unique maximal open substack of rank-*2*, degree-d bundles admitting a* separated adequate *moduli space. By contrast, if* $g(C) > 2$*, there is an open dense* $U \subset \mathrm{Bun}_3^2$*, not contained in* $\mathrm{Bun}_3^{2,\mathrm{ss}}$*, which admits a separated non-proper good moduli space.*

In characteristic zero, adequate and good moduli spaces agree. Thus the first statement intrinsically recovers semistability from schematicity, the second does so from separatedness in rank two, and the third proves that separatedness alone fails in higher rank.

Ingredient 1: line bundles turn schematicity into stability

If an open substack V of a smooth algebraic stack with affine diagonal has a quasi-projective adequate moduli space, Theorem 2.3 produces a line bundle $\mathcal{L}$ on the ambient stack such that

$$V \subseteq (\mathrm{Bun}_G)_{\mathcal{L}}^{\text{a-ss}} \subseteq (\mathrm{Bun}_G)_{\mathcal{L}}^{\Theta\text{-ss}}.$$

For G almost simple and simply connected, $\mathrm{Pic}(\mathrm{Bun}_G) \cong \mathbf{Z}$, generated (up to sign) by the determinant-of-cohomology line bundle $\mathcal{L}_{\det}$. Its positive powers have classical semistable locus as their Θ-semistable locus; exponent zero has empty adequate-semistable locus, and a negative exponent has empty Θ-semistable locus. Therefore $V \subseteq \mathrm{Bun}_G^{\mathrm{ss}}$.

For general reductive G, a central isogeny with multiplicative-type kernel μ reduces to products of simply connected almost-simple factors and a torus. Adequate affineness passes through the resulting Bun_μ-torsor. The Picard-group decomposition forces positive determinant-line exponents on every semisimple factor, while the torus factor descends to an algebraic space and contributes no Θ-weight. This closes the same containment and proves the first theorem.

Ingredient 2: two valuative tests for separated moduli

The existence theorem used in Section 3 says that a quasi-compact algebraic stack, locally of finite type over a field, with affine diagonal and locally reductive, has a separated adequate moduli space exactly when it is both Θ-reductive and S-complete. The authors translate these conditions into bundle data.

Θ-reductivity says that a filtration of the generic fiber of a DVR family extends without its special associated graded leaving the chosen open substack (Proposition 3.6). S-completeness says that whenever two bundles E, E' admit *opposite filtrations*—their successive graded pieces match in reverse order—their common associated graded must remain in the open substack (Proposition 3.17). Thus two abstract test stacks become concrete statements about degenerations and filtrations of bundles.

Ingredient 3: rank two creates a contradiction

Suppose an open $V \subset \mathrm{Bun}_2^d$ contains an unstable bundle. Its Harder–Narasimhan sequence has line-bundle factors of unequal degrees. By varying the factors in $\mathrm{Pic}^0(C)$, Proposition 4.6 flips this extension and produces two bundles in V with opposite filtrations. If V is S-complete, their associated graded—a direct sum $L_1 \oplus L_2$ with $\deg L_1 \neq \deg L_2$—must also lie in V.

But an elementary modification of such a direct sum supplies a Θ-testing family whose generic and special extensions lie in V while the limiting associated graded does not. Proposition 4.2 therefore says that a Θ-reductive open cannot contain this unequal-degree sum. The two valuative requirements contradict one another, so every open with separated adequate moduli lies inside the semistable locus. This rank-two flip is the clean toy model for the paper's stacky method.

Ingredient 4: the rank-three escape

For rank 3, degree 2, let λ be the Harder–Narasimhan polygon with vertices $(0,0), (1,1), (3,2)$. Inside the open locus of polygons below λ, remove every bundle fitting into

$$0 \to L \to E \to F \to 0 \quad \text{or} \quad 0 \to F \to E \to L \to 0,$$

where $\operatorname{rk} L = 1$, $\deg L = 1$, $\operatorname{rk} F = 2$, $\deg F = 1$, and F is not $(1,0)$-stable. Here a rank-2, degree-1 bundle is $(1,0)$-stable exactly when every line subbundle has negative degree. The complement U still contains unstable sums $L \oplus F$ when F is $(1,0)$-stable. The polygon bound and the removal rule make U locally reductive, Θ-reductive, and S-complete, so it has a separated good moduli space.

It is not proper. A family can have generic rank-two factor F_K $(1,0)$-stable but special factor F_κ stable and not $(1,0)$-stable. A suitable extension remains a stable rank-three family; its generic fiber lies in U and its special fiber does not. Langton uniqueness prevents replacing the special factor by an allowed one.

Hard point and remaining question

The conceptual difficulty is making the two valuative conditions modular enough to control all degenerations, while the technical heart of the schematic theorem is the central-isogeny and Picard-group reduction for a general reductive group. The paper does not settle whether the semistable locus is the unique maximal open admitting a *proper* good moduli space in higher rank. Its rank-three example shows exactly why "separated" cannot replace "proper" in that question.