

Common preperiodic points for quadratic polynomials

Laura DeMarco, Holly Krieger, Hexi Ye

arXiv:1911.02458

<http://arxiv.org/abs/1911.02458v2>

A finite configuration determines the parameter

For $c \in \mathbf{C}$ let $f_c(z) = z^2 + c$ and write $\text{Preper}(f_c) \subset \mathbf{P}^1(\mathbf{C})$ for its preperiodic points. Equality of the full preperiodic sets already forces equality of the parameters. The paper proves the much stronger, quantitative statement below.

Main theorem (DeMarco–Krieger–Ye, Theorems 1.1 and 11.1). *For every pair $c_1 \neq c_2$ in $\mathbf{C}$,*

$$|\text{Preper}(f_{c_1}) \cap \text{Preper}(f_{c_2})| \leq 10^{103}. \quad (1)$$

In particular, the bound is independent of the parameters.

The size of the displayed constant is not the point: the known examples are tiny by comparison. The achievement is uniformity even as the two parameters approach one another and their Julia sets become difficult to distinguish.

The global measuring device

Suppose first that $c_1, c_2 \in \overline{\mathbf{Q}}$ and choose a number field K containing them. At a place v let $\lambda_{c,v}$ be the escape-rate function and $\mu_{c,v}$ the equilibrium measure on the Berkovich affine line. With $r_v = [K_v : \mathbf{Q}_v]/[K : \mathbf{Q}]$, the adelic energy

$$\langle f_{c_1}, f_{c_2} \rangle = \sum_{v \in M_K} r_v \int \lambda_{c_1,v} d\mu_{c_2,v} \quad (2)$$

is symmetric and nonnegative, and vanishes exactly when $c_1 = c_2$. It simultaneously records how far apart the two canonical measures are at every archimedean and nonarchimedean place. Write $E_v = \int \lambda_{c_1,v} d\mu_{c_2,v}$ for the local summand and h for the logarithmic Weil height on $\mathbf{A}^2(\overline{\mathbf{Q}})$.

The proof constructs two inequalities pulling (2) in opposite directions. For absolute constants, Theorems 1.6–1.8 give

$$\langle f_{c_1}, f_{c_2} \rangle \geq \delta > 0, \quad (3)$$

$$\frac{1}{192}h(c_1, c_2) - \frac{3}{17} \leq \langle f_{c_1}, f_{c_2} \rangle \leq \frac{1}{2}h(c_1, c_2) + \frac{7}{3}, \quad (4)$$

$$\langle f_{c_1}, f_{c_2} \rangle \leq \left(\varepsilon + \frac{C(\varepsilon)}{N-1} \right) (h(c_1, c_2) + 1), \quad (5)$$

where N is the number of common preperiodic points, $0 < \varepsilon < 1$, and one may take $C(\varepsilon) = 40 \log(25/\varepsilon)$. The authors make (3) effective with $\delta = 10^{-96}$.

Ingredient I: local separation and helpful places

At the complex places, Koebe distortion and explicit control of the first few preimages of 0 bound the energy between the two Julia measures. At a nonarchimedean place away from 2, the Julia tree gives

$$\frac{1}{4} \log^+ |c_1 - c_2|_v \leq E_v \leq \frac{1}{2} \log^+ \max\{|c_1|_v, |c_2|_v\}; \quad (6)$$

the prime above 2 requires a weaker lower bound and an additive correction.

For fixed $L \geq 1$, define

$$\ell_v = \begin{cases} \log \max\{|c_1|_v, |c_2|_v, L\}, & v \mid \infty, \\ \log \max\{|c_1|_v, |c_2|_v, 16\}, & v \mid 2, \\ \log \max\{|c_1|_v, |c_2|_v, 1\}, & \text{otherwise,} \end{cases} \quad h_L(c_1, c_2) = \sum_v r_v \ell_v. \quad (7)$$

To obtain (4), the proof calls a large place *helpful* when c_1 and c_2 are not too v -adically close. At every helpful place the local estimates imply $E_v \geq \ell_v/64$. The product formula for $c_1 - c_2$ prevents almost all of the height from hiding at close places; quantitatively,

$$\sum_{v \text{ helpful}} r_v \ell_v \geq \frac{1}{3} h_L(c_1, c_2) - O(1). \quad (8)$$

Summing (8), with $L = 1000$, gives the strong height lower bound (4). A related archimedean dichotomy proves (3): if the total energy tended to zero, many complex embeddings would force $c_1 - c_2$ to be extremely small; the product formula would then make its Weil height large, contradicting the weaker global lower bound in terms of $h(c_1 - c_2)$.

Ingredient II: many common points force small energy

Remove the always-common fixed point ∞ and let F be the remaining Galois-invariant set of common preperiodic points, so $|F| = N - 1$. Replace its atomic measure by small circles of adelic radii η_v . The energy metric satisfies the triangle inequality

$$\langle f_{c_1}, f_{c_2} \rangle^{1/2} \leq d(\mu_{c_1}, [F]_\eta) + d(\mu_{c_2}, [F]_\eta). \quad (9)$$

The radii cannot be arbitrarily small, because the self-energy of an atom then blows up. The decisive choice is $\eta_v = \rho^{3\ell_v}$. Archimedean distortion and its Berkovich analogues show that regularization changes the potential by at most $\rho\ell_v$, while the self-energy costs $3\ell_v \log(1/\rho)/(2|F|)$. After summing over all places, (9) becomes

$$\langle f_{c_1}, f_{c_2} \rangle \leq 4 \left(\rho + \frac{3 \log(1/\rho)}{2(N-1)} \right) h_L(c_1, c_2), \quad (10)$$

which yields (5). This is the proof's central tension: a large common finite set tries to approximate both canonical measures, hence tries to make two distinct dynamical systems adelically indistinguishable.

Closing the argument

For parameters of large height, compare (4) and (5) with $\varepsilon = 1/384$; the height terms then force a uniform bound on N . For bounded height, compare (5) with the height-independent gap (3). The explicit constants give $N < 10^{103}$ over $\overline{\mathbf{Q}}$.

Finally suppose the parameters are complex. If $B + 1$ distinct common points existed, their two eventual-periodicity relations would place the parameters and points in a function field of a curve over a number field. Away from finitely many parameters one may specialize to algebraic values while preserving distinctness, the two parameters, and all orbit relations. This would produce $B + 1$ algebraic common preperiodic points, a contradiction.

A toy picture and the genuine difficulty

At a good-reduction place both canonical measures sit at the Gauss point, so that place contributes no separation. At a helpful bad place, the two Julia trees branch at measurably different scales and contribute a definite fraction of ℓ_v . The product formula says these local discrepancies cannot all be suppressed at once. This picture captures (8), but not the hardest analytic step: the complex Julia sets require uniform distortion estimates as $|c| \rightarrow \infty$ and as $c_1 - c_2 \rightarrow 0$. Nor does the paper establish the analogous uniformity conjectured for all rational maps of a fixed degree; (1) is specific to the monic quadratic family.