

Fano varieties with torsion in the third cohomology group

John Christian Ottem, Jorgen Vold Rennemo

arXiv:2309.10793

<http://arxiv.org/abs/2309.10793v2>

A Fano analogue of the Artin–Mumford phenomenon

For a smooth complex projective variety, torsion in $H^3(X, \mathbb{Z})$ is a stable birational invariant. It vanishes for all Fano varieties of dimension at most three, prompting the question whether Fano geometry forces torsion-freeness in every dimension.

Main theorem (Ottem–Rennemo, Theorems 1.1 and 1.2). *For every even $d \geq 4$ there is a smooth d -dimensional Fano variety X of Picard rank one with*

$$H^3(X, \mathbb{Z}) \simeq \mathbb{Z}/2. \quad (1)$$

For $d \geq 6$, its coniveau and strong-coniveau filtrations differ:

$$0 = \tilde{N}^1 H^3(X, \mathbb{Z}) \neq N^1 H^3(X, \mathbb{Z}) = H^3(X, \mathbb{Z}) \simeq \mathbb{Z}/2. \quad (2)$$

These give the first Fano varieties with torsion in H^3 and hence Fano varieties that are rationally connected but not stably rational. Formula (2) also supplies higher-dimensional rationally connected examples where ordinary support on a divisor cannot be upgraded to pushforward from a smooth divisor.

The construction from rank-four quadrics

Let $V = \mathbb{C}^n$ and let $\overline{Z}_{r,n} \subset \mathbb{P}(\mathrm{Sym}^2 V^\vee)$ be the locus of quadrics of rank at most r . For even r , a rank- r quadric has two connected families of maximal isotropic linear spaces. Their Stein factorization defines a double cover

$$\sigma : W_{r,n} \longrightarrow \overline{Z}_{r,n}, \quad (3)$$

ramified over $\overline{Z}_{r-1,n}$. Write $H = \sigma^* \mathcal{O}(1)$. The paper concentrates on $r = 4$, where

$$\dim W_{4,n} = 4n - 7, \quad K_{W_{4,n}} = -2nH, \quad \mathrm{Sing} W_{4,n} = \sigma^{-1}(\overline{Z}_{2,n}). \quad (4)$$

For general $H_i \in |H|$, set

$$X_{n,c} = W_{4,n} \cap H_1 \cap \cdots \cap H_c. \quad (5)$$

The desired variety is $X = X_{n,2n-1}$ for $n \geq 5$; its dimension is $2n - 6$. Since the singular locus in (4) has dimension $2n - 2$, a general codimension- $(2n - 1)$ section misses it. Bertini therefore makes X smooth, while adjunction gives $K_X = -H$. Thus X is Fano, and the low-degree comparison below gives $H^2(X, \mathbb{Z}) = \mathbb{Z}$; together with Fano vanishing, this gives Picard rank one. Putting $n = d/2 + 3$ realizes every even $d \geq 4$.

Ingredient I: a classifying-space source of 2-torsion

Choose a four-dimensional quadratic space S . Orthogonal invariant theory identifies the double cover with a connected-similitude quotient

$$W_{4,n} \simeq \mathrm{Hom}(V, S)^{\mathrm{ss}} // \mathrm{GO}(S)^\circ. \quad (6)$$

On the complement of the preimage of the rank-two locus the action is free, so the smooth part of $W_{4,n}$ receives low-degree cohomology from $B\mathrm{GO}(4)^\circ$.

The starting computation is

$$H^*(B\mathrm{SO}(4), \mathbb{Z}) = \mathbb{Z}[\nu, e, p]/(2\nu), \quad |\nu| = 3, \quad |e| = |p| = 4, \quad (7)$$

and the mod-2 reduction of ν is w_3 , so $\bar{\nu}^2 = w_3^2$ is nonzero. The exact sequence $1 \rightarrow \mathrm{SO}(4) \rightarrow \mathrm{GO}(4)^\circ \rightarrow \mathbb{C}^* \rightarrow 1$ and its Gysin sequence then yield (Proposition 3.1)

$$H^1(B\mathrm{GO}(4)^\circ, \mathbb{Z}) = 0, \quad H^2(B\mathrm{GO}(4)^\circ, \mathbb{Z}) = \mathbb{Z}, \quad H^3(B\mathrm{GO}(4)^\circ, \mathbb{Z}) = \mathbb{Z}/2, \quad (8)$$

with the square of the nonzero degree-three class still nonzero modulo 2.

Ingredient II: two Lefschetz comparisons preserve the class

The excluded subset of $\mathrm{Hom}(V, S)$ has codimension $n - r + 2 = n - 2$. Equivariant cohomology therefore agrees with that of a point in degrees below $2n - 5$ (Lemmas 3.2–3.3). A generalized Lefschetz theorem of Goresky–MacPherson compares the smooth quotient with a general linear section. In the range needed for $X = X_{n,2n-1}$, the two comparisons give

$$H^3(B\mathrm{GO}(4)^\circ, \mathbb{Z}) \xrightarrow{\sim} H^3(X, \mathbb{Z}) \quad (n \geq 5), \quad H^6(B\mathrm{GO}(4)^\circ, \mathbb{Z}/2) \hookrightarrow H^6(X, \mathbb{Z}/2) \quad (n \geq 6). \quad (9)$$

The geometric range is controlled by $\dim X = 2n - 6$, while the equivariant range is $j < 2n - 5$. Thus, when $n \geq 5$, (9) transports (8) to $H^3(X, \mathbb{Z}) = \mathbb{Z}/2$, proving Theorem 4.1 and hence (1). When $n \geq 6$, the comparison is also injective in degree six; Corollary 3.6 therefore preserves the nonvanishing

$$\bar{\alpha}^2 \neq 0 \quad \text{in } H^6(X, \mathbb{Z}/2). \quad (10)$$

where α generates $H^3(X, \mathbb{Z})$.

Why the square detects strong coniveau

Since X is Fano, it is rationally connected, and Proposition 5.1 gives $N^1 H^l(X, \mathbb{Z}) = H^l(X, \mathbb{Z})$ for every $l > 0$. On the other hand, the topological obstruction recalled as Proposition 5.2 says that any degree-three class of strong coniveau at least one must have square zero after reduction modulo 2. Equation (10) excludes the unique nonzero class α from $\tilde{N}^1 H^3$. Because the whole group has order two, the strong-coniveau subgroup is zero, proving (2).

Geometrically, when the section is rationally connected, the torsion generator corresponds under $\mathrm{Br}(X) \simeq \mathrm{Tors} H^3(X, \mathbb{Z})$ to the Brauer–Severi family obtained by restricting the maximal-isotropic-space fibration over $W_{4,n}$. This is the promised higher-dimensional echo of the Artin–Mumford construction.

The fourfold model and the limit of the obstruction

The smallest example is $X_{5,9}$, a Fano fourfold with

$$\mathrm{Pic}(X) = \mathbb{Z}H, \quad -K_X = H, \quad H^4 = 10, \quad H^3(X, \mathbb{Z}) = \mathbb{Z}/2.$$

It already answers the torsion question. However, its generator has square zero modulo 2, so the paper does not determine whether it has strong coniveau one. This explains the $d \geq 6$ threshold in (2): the construction exists in dimension four, but the degree-six injectivity needed to carry the square obstruction through the linear section begins only at $n = 6$.

The smooth Fano examples of Theorem 1.1 occur only in even dimensions. The paper also constructs odd-dimensional smooth unirational, non-stably rational varieties by resolving different linear sections, but they are not the Fano examples above. It remains open whether rationally connected fourfolds or fivefolds can have unequal coniveau filtrations by some other mechanism.