

On automorphisms of moduli spaces of parabolic vector bundles

Carolina Araujo, Thiago Fassarella, Inder Kaur, Alex Massarenti

arXiv:1902.04136

<http://arxiv.org/abs/1902.04136v1>

The central parabolic-bundle model

Fix $n \geq 5$ general points $p_1, \dots, p_n \in \mathbf{P}^1$ and set $S = p_1 + \dots + p_n$. A rank-two quasi-parabolic bundle is a degree-zero bundle E with a line $V_i \subset E_{p_i}$ at each marked point. For a weight vector $\mathcal{A} = (a_1, \dots, a_n)$, stability compares the parabolic slopes

$$\mu_{\mathcal{A}}(E) = \frac{1}{2} \sum_i a_i, \quad \mu_{\mathcal{A}}(L, E) = \deg L + \sum_{L_{p_i} = V_i} a_i \quad (1)$$

for every line subbundle $L \subset E$. Let $\mathcal{M}_{\mathcal{A}}$ be the resulting projective moduli space. At the central weight $\mathcal{A}_F = (1/2, \dots, 1/2)$, the variety $\mathcal{M}_{\mathcal{A}_F}$ is Fano of dimension $n - 3$; it is smooth for n odd and has finitely many singular points for n even.

For an even subset $R \subset \{1, \dots, n\}$, the elementary transform el_R is defined by

$$0 \longrightarrow E' \longrightarrow E \longrightarrow \bigoplus_{i \in R} (E_{p_i}/V_i) \otimes \mathcal{O}_{p_i} \longrightarrow 0, \quad E' \mapsto E' \otimes \mathcal{O}_{\mathbf{P}^1}(|R|/2), \quad (2)$$

with the induced parabolic directions. These involutions form the group $\text{El} \simeq (\mathbf{Z}/2\mathbf{Z})^{n-1}$ and preserve the central weight.

Main theorem (Araujo–Fassarella–Kaur–Massarenti, Theorem 1.2). *For $n \geq 5$ general marked points,*

$$\text{Aut}(\mathcal{M}_{\mathcal{A}_F}) = \text{El} \simeq (\mathbf{Z}/2\mathbf{Z})^{n-1}.$$

Thus every automorphism has the modular interpretation of an elementary transformation at an even collection of parabolic points.

Ingredient I: the nilpotent cone remembers the flags

For a parabolic bundle $(E, \mathbf{v})$ on a curve and a line bundle L , consider the rank-three bundle of traceless strongly parabolic endomorphisms $\text{SPEnd}_0(E) \otimes L$. The closure of the fiberwise nilpotent endomorphisms is a quadratic cone bundle N_E . Near a marked point, choose a parameter t and a basis with $V_i = \langle (1, 0) \rangle$. A local section has matrix

$$\begin{pmatrix} t\alpha & \beta \\ t\gamma & -t\alpha \end{pmatrix}, \quad \det = -t(t\alpha^2 + \beta\gamma),$$

so in fiber coordinates (a, b, c) the cone is

$$ta^2 + bc = 0. \quad (3)$$

Its projectivization has a reducible conic over $t = 0$: one component maps to the fiber of $\mathbf{P}(E)$ and the other contracts to $\mathbf{P}(V_i)$. Gluing this description proves Proposition 3.2:

$$\mathbf{P}(N_E) \simeq \text{Bl}_{\{\mathbf{P}(V_1), \dots, \mathbf{P}(V_n)\}} \mathbf{P}(E) \quad \text{over } \mathbf{P}^1. \quad (4)$$

Consequently the nilpotent cone recovers precisely the ruled surface with its parabolic markings; isomorphism of these blowups is equivalent to an elementary transformation.

Ingredient II: the Hitchin system is intrinsic

On the stable locus, parabolic Serre duality identifies

$$T_{(E, \mathbf{v})}^* \mathcal{M}_{\mathcal{A}_F}^s \simeq H^0(\mathbf{P}^1, \text{SPEnd}_0(E) \otimes \omega_{\mathbf{P}^1}(S)). \quad (5)$$

The determinant defines the Hitchin map

$$h : T^* \mathcal{M}_{\mathcal{A}_F}^s \longrightarrow H^0(\mathbf{P}^1, \omega_{\mathbf{P}^1}^{\otimes 2}(S)). \quad (6)$$

For general a , the spectral curve $C_a \subset \text{Tot}(\omega_{\mathbf{P}^1}(S))$ is the double cover cut out by $s^2 + \pi^* a = 0$; it is smooth of genus $n - 3$, and the completed Hitchin fiber is the polarized abelian variety $\text{Pic}^{n-2}(C_a)$ (Proposition 2.7). Crucially, (6) is the affinization of the cotangent bundle, so an automorphism φ of the moduli space intrinsically induces a map f on the Hitchin base.

Ingredient III: Torelli fixes the Hitchin base

Fiber dilations make $d\varphi$ equivariant for the action $a \mapsto t^2 a$, forcing f to be linear. For general a , the induced isomorphism of Hitchin fibers extends across the codimension-two complement to an isomorphism of polarized Picard varieties

$$\mathrm{Pic}^{n-2}(C_a) \simeq \mathrm{Pic}^{n-2}(C_{f(a)}). \quad (7)$$

It commutes with the involutions induced by the two sheets. Torelli then recovers an isomorphism of spectral curves descending to an automorphism of $\mathbf{P}^1$ that carries the branch locus of C_a to that of $C_{f(a)}$. Those branch loci contain the fixed general points $p_1, \dots, p_n$, so the descended automorphism is the identity. Hence $f(a)$ is a scalar multiple of a ; rescaling reduces to $f = \mathrm{id}$.

Ingredient IV: recover the parabolic bundle

For a general $(E, \mathbf{v})$ and $n \geq 6$, the underlying degree-zero bundle is free, and Lemma 3.3 proves that $\mathrm{SPEnd}_0(E) \otimes \omega_{\mathbf{P}^1}(S)$ is globally generated. Indeed, parabolic Serre duality reduces this to

$$H^0(\mathbf{P}^1, \mathrm{PEnd}_0(E) \otimes \mathcal{O}(1)) = 0 : \quad (8)$$

traceless 2×2 matrices of linear forms constitute a six-dimensional space, while the conditions imposed by the n general parabolic directions have rank six (already six are independent). Evaluation therefore gives a quotient

$$T_E^* \mathcal{M}_{\mathcal{A}_F}^s \times \mathbf{P}^1 \twoheadrightarrow \mathrm{SPEnd}_0(E) \otimes \omega_{\mathbf{P}^1}(S). \quad (9)$$

The inverse image of the fiberwise nilpotent cone is characterized solely by the restricted Hitchin map: over p , it consists of Higgs fields whose determinant vanishes at p . Since $d\varphi$ fixes the Hitchin base, it preserves this cone and its vertex, hence descends through (9) to an isomorphism $N_E \simeq N_{E'}$ for $E' = \varphi(E)$. Formula (4) now says that $(E, \mathbf{v})$ and $(E', \mathbf{v}')$ differ by an elementary transformation. Equality on a general moduli point proves the group identity. The case $n = 5$ is supplied by the previously known degree-four del Pezzo calculation.

Toy model, extension in weight space, and limits

The local surface $ta^2 + bc = 0$ is the proof's toy model. Away from $t = 0$ its projectivized fibers are smooth conics; at the marked point the conic splits into two lines, and choosing which line to contract records the elementary transformation. What the toy model suppresses is the global task of recognizing this cone using only the intrinsic Hitchin geometry.

For weights in the interior of a specific truncated demi-hypercube Π , the model $\mathcal{M}_{\mathcal{A}}$ is a small modification of the central Fano model. Corollary 1.4 shows $\mathrm{Aut}(\mathcal{M}_{\mathcal{A}}) = \mathrm{El}_{\mathcal{A}}$, the subgroup whose reflected weight remains in the same stability chamber: an automorphism induces a pseudo-automorphism of the central Fano model, which is an automorphism and hence elementary. Outside this region the equality need not hold (some models are projective spaces). The theorem also relies on general marked points; it does not classify extra symmetries arising from special configurations. Section 5 also derives an involutorial-bundle and Kummer-variety application from the central model; that secondary application is omitted here.