

Uncritical polynomial maps with rational multipliers

Valentin Huguin

arXiv:2009.02422

<http://arxiv.org/abs/2009.02422v1>

Can the cycle multipliers determine the map?

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a polynomial of degree $d \geq 2$. If z has exact period n , its cycle multiplier is

$$\lambda_f(z) = (f^n)'(z) = \prod_{j=0}^{n-1} f'(f^j(z)); \quad (1)$$

it is constant along the cycle and invariant under affine conjugacy. The map is *unicritical* when it has only one finite critical point. A power map is conjugate to z^d , while a Chebyshev map is conjugate to $\pm T_d$, where $T_d(z + z^{-1}) = z^d + z^{-d}$.

Main theorem (Huguin, Theorem 6). *If a unicritical polynomial map $f : \mathbb{C} \rightarrow \mathbb{C}$ has rational multiplier at every periodic cycle, then f is a power map or a Chebyshev map.*

Since T_d has $d - 1$ critical points, the Chebyshev alternative is unicritical only in degree 2. Thus the conclusion in normalized coordinates is especially rigid: the parameter below is $c = 0$, except that $c = -2$ is also allowed when $d = 2$. This is evidence for Milnor's broader conjectural classification of rational maps with integral multipliers.

The algebraic encoder: multiplier polynomials

Every unicritical degree- d polynomial is affinely conjugate to

$$f_c(z) = z^d + c, \quad (2)$$

with c unique up to multiplication by a $(d - 1)$ st root of unity. The dynatomic polynomials separate formal periods by

$$f^n(z) - z = \prod_{k|n} \Phi_k^f(z).$$

Propositions 30 and 38 attach to a monic f a monic multiplier polynomial M_n^f through

$$(M_n^f(\lambda))^n = \text{Res}_z(\Phi_n^f(z), \lambda - (f^n)'(z)). \quad (3)$$

The n points in one exact n -cycle contribute the same factor to the resultant, explaining the n th root. Corollary 42 is the decisive bridge: for a subring $R \subset \mathbb{C}$, all exact-period- n multipliers lie in R exactly when M_n^f splits into linear factors over R . Formal-period points of smaller period can add only the controlled root $\lambda = 1$ (Propositions 34 and 41).

For (2), write $M_n(c, \lambda) = M_n^{f_c}(\lambda)$ and $\Delta_n(c) = \text{disc}_\lambda M_n(c, \lambda)$. Rational splitting forces the coefficients to be rational and each discriminant to be a rational square. The proof now needs only a few small periods.

Three degree regimes

Quadratic case. For $d = 2$, the period-1 through period-4 multiplier polynomials suffice (Proposition 17). Their splitting first forces $c \in \mathbb{Q}$, and the explicit fourth discriminant is

$$\Delta_4(c) = -2^{24}(64c^3 + 144c^2 + 108c + 135)(c + 2)^2c^6. \quad (4)$$

Outside $c = 0, -2$, squarehood of (4) would give an $r \in \mathbb{Q}$ with

$$-(64c^3 + 144c^2 + 108c + 135) = r^2.$$

The rational substitution in the proof turns this into $a^3 + b^3 = 4$. Clearing denominators produces integers x, y, z with $z \neq 0$ and $x^3 + y^3 = 4z^3$, contradicting Lemma 56. Consequently f_c is z^2 or $T_2(z) = z^2 - 2$.

Cubic case. Rational multipliers are real. Splitting of the period-1 and period-2 polynomials makes c^2 real and their discriminants satisfy

$$\Delta_1 = -3^6(27c^2 - 4)c^2 \geq 0, \quad \Delta_2 = -3^{12}(27c^2 + 32)c^6 \geq 0. \quad (5)$$

The first inequality puts c^2 in $[0, 4/27]$, the second in $[-32/27, 0]$; hence $c = 0$ (Proposition 23).

Degree at least four. Fixed points already suffice. Example 24 computes

$$M_1(c, \lambda) = \lambda(\lambda - d)^{d-1} + (-d)^d c^{d-1}. \quad (6)$$

If its roots are real, so are those of

$$L(\lambda) = \lambda^d M_1(c, \lambda^{-1} + d) = (-d)^d c^{d-1} \lambda^d + d\lambda + 1.$$

Rolle's theorem then makes every root of $L'(\lambda) = (-1)^d d^{d+1} c^{d-1} \lambda^{d-1} + d$ real. For $c \neq 0$ these roots form an orbit under the $(d - 1)$ st roots of unity, which cannot be wholly real when $d - 1 \geq 3$. Thus $c = 0$ (Proposition 25).

The hard arithmetic step

The obstruction used after (4) is proved rather than delegated to a database of rational points. In the Eisenstein integers $A = \mathbb{Z}[j]$, with $j = e^{2\pi i/3}$ and $\lambda = 1 - j$, cubes modulo λ^3 have only the residue classes $-1, 0, 1$. Lemma 59 assumes a primitive nonzero solution of

$$x^3 + uy^3 = 4vz^3, \quad u, v \in A^\times,$$

with minimal λ -adic valuation of z . The congruence forces $\lambda \mid z$ and $u = \pm 1$. The common factor must first be removed:

$$\begin{aligned} x + uy &= \lambda a, & jx + j^2uy &= \lambda b, & j^2x + juy &= \lambda c, \\ z &= \lambda d, & a + b + c &= 0, & abc &= 4vd^3. \end{aligned} \tag{7}$$

The normalized factors a, b, c are pairwise coprime. Unique factorization writes two as unit times cubes and the third as four times a unit times a cube. Their relation $a + b + c = 0$ produces a new solution with

$$\text{ord}_\lambda(Z) = \text{ord}_\lambda(z) - 1,$$

contradicting minimality. This infinite descent is the genuinely arithmetic heart of the quadratic case.

Toy model, proof roadmap, and scope

For $f(z) = z^d$, every nonzero point fixed by f^n satisfies $z^{d^n-1} = 1$, so (1) gives $(f^n)'(z) = d^n$; the fixed critical point has multiplier 0. The Chebyshev examples come from quotienting this power map by the involution $z \leftrightarrow z^{-1}$. This explains why both model families have integral multipliers, but not why no other unicritical map does.

The full proof is therefore

$$\text{cycles} \Rightarrow M_n \Rightarrow \text{rational splitting} \Rightarrow \begin{cases} \text{Eisenstein descent,} & d = 2, \\ \text{two sign constraints,} & d = 3, \\ \text{real-root geometry,} & d \geq 4. \end{cases} \tag{8}$$

The theorem assumes rationality for every cycle, although the proof shows that periods at most 4 suffice for $d = 2$, periods 1, 2 suffice for $d = 3$, and fixed points suffice for $d \geq 4$. It does not classify general polynomials or rational maps. The paper's separate cubic-with-symmetries theorem assumes *integer*, not merely rational, multipliers; its rational analogue is left open.