Rational Periodic Points of x-to-the-d Plus c and Fermat–Catalan Equations

Chatchawan Panraksa

arXiv:2105.03715

http://arxiv.org/abs/2105.03715v6

Cycles become high-power Diophantine equations

For $d > 2$ and $c \in \mathbf{Q}$, write

$$f_{d,c}(x) = x^d + c.$$

A rational point is preperiodic when its forward orbit is finite, and is periodic of exact period n when n is the first positive iterate returning it to itself. The paper asks whether the size of $\mathrm{PrePer}(f_{d,c}, \mathbf{P}^1(\mathbf{Q}))$ can be bounded independently of the degree inside this family. Its principal large-degree result is conditional, but notably uniform in the parameter c.

Main theorem (Panraksa, Theorem 3)**.** *Assume the abc-conjecture. For all sufficiently large d,*

(i) *$f_{d,c}$ has no rational periodic point of exact period greater than one unless $c = -1$;*
(ii) $\#\,\mathrm{PrePer}(f_{d,c}, \mathbf{P}^1(\mathbf{Q})) \le 4$.

If the paper's explicit abc-conjecture is assumed, both conclusions hold for every $d \ge 7$.

The exceptional map for even d really has a nontrivial cycle: $0 \mapsto -1 \mapsto 0$ under $x^d - 1$; also $1 \mapsto 0$, while ∞ is fixed. Thus four is the natural bound retained by the theorem.

Ingredient 1: one denominator for an entire orbit

A valuation lemma says that if X/Z is a rational preperiodic point, written in lowest terms, and $c = C/N$ is also reduced, then $N = Z^d$. Hence every point of a rational cycle can be written $x_i = X_i/Z$ with the same denominator and $c = C/Z^d$. The product identity

$$\prod_i \frac{X_i^d - X_{i+1}^d}{X_i - X_{i+1}} = Z^{n(d-1)}$$

also yields $\gcd(X_i, X_{i+1}) = 1$ (Corollary 1 and Theorem 8). This is the arithmetic compression step: all iterates disappear into a single denominator and coprimality condition.

Choose the cyclic indexing so that $|x_2|$ is maximal. Since $f(x_2) - f(x_1) = x_3 - x_2$, clearing denominators gives

$$X_2^d - X_1^d = (X_3 - X_2)Z^{d-1}, \qquad \gcd(X_1, X_2) = 1,$$

with $|X_2| = \max_i |X_i|$. Unless $|X_2|$ and $|Z|$ are both one, this is a nonzero coprime *abc* equation whose three terms have degree d but whose radical involves only $X_1, X_2, X_3 - X_2$, and Z.

Ingredient 2: the radical cannot keep pace

Let $W = \max(|X_1|, |X_2|, |X_3|, |Z|) > 1$. Lemma 2 applies *abc* to the last equation. Because $|X_1|, |X_3| \le |X_2|$ and $|X_3 - X_2| \le 2|X_2|$, the radical is bounded by a constant times W^4. Separating the cases $|X_2| \le |Z|$ and $|X_2| > |Z|$ gives, for every $\varepsilon > 0$,

$$d < \log_b(2^{1+\varepsilon} K_\varepsilon) + 5 + 4\varepsilon \quad (1 < b \le W),$$

where K_ε is the *abc* constant. Taking $b = 2$ contradicts the existence of such an orbit once d is large. If $W = 1$, then c is integral; the elementary integral-orbit argument leaves only fixed points and the cycle for $c = -1$.

The explicit form supplies numerical radical bounds. It rules out the displayed system for $d \ge 10$; odd $d = 7, 9$ are already covered by the cited odd-degree theorem, while the paper handles $d = 8$ using the resulting height cutoff together with finite searches (and Hutz's table for periods above two). These computations are part of the proof of the stated threshold.

Ingredient 3: from cycles to all finite orbits

Excluding cycles is not yet a bound on preperiodic tails. For even d and $c \neq 0$, Lemma 4 uses the same *abc* equation (and the correspondence between two fixed points and a two-cycle for a related parameter) to leave at most one rational finite fixed point in large degree. If x_0 is that fixed point, evenness makes $-x_0$ the only possible distinct rational point mapping to it. If an alleged point r lay two steps before x_0, write $r = R/Z$ and $x_0 = X_0/Z$ in lowest terms, so $\gcd(R, Z) = \gcd(X_0, Z) = 1$. Lemma 5 applies *abc* once more to the resulting equation

$$X_0^{d-1} - \frac{R^d}{X_0} = 2Z^{d-1},$$

and excludes such a point. The omitted parameter $c = 0$ is elementary: for even d, its rational preperiodic set is contained in $\{\infty, 0, 1, -1\}$, with $-1 \mapsto 1$. The cited odd-degree result supplies the bound four for odd d. Together with the explicit $c = -1$ orbit, these statements prove the second half of Theorem 3.

Model, hard point, and scope

For a two-cycle $x_1 \leftrightarrow x_2$, subtracting the two orbit equations already gives

$$\frac{x_1^d - x_2^d}{x_1 - x_2} = -1.$$

For even $d = 2k$, the factorization and coprimality in Theorem 8 refine the common-denominator form to

$$X_1^k + X_2^k = \delta Z_1^{2k-1}, \qquad \delta \in \{1, 2\},$$

with $\delta = 1$ when k is odd. This toy case captures the full proof's mechanism: dynamics provides a high-power identity, while coprimality makes its radical too small.

The hard input is precisely the conjectural radical estimate; the claimed large-degree theorem is not unconditional. The explicit threshold also uses finite computations recorded or cited in the paper. The digest does not cover the paper's separate classifications for $d = 4, 6$ or its Fermat–Catalan obstruction to certain two-cycles, and the theorem does not classify rational preperiodic graphs in the remaining small even degrees.