

On the Hilbert Property and the Fundamental Group of Algebraic Varieties

Pietro Corvaja, Umberto Zannier

arXiv:1601.07066

<http://arxiv.org/abs/1601.07066v2>

Hilbert irreducibility meets topology

For an integral quasi-projective variety X over a number field k , the *Hilbert property* (HP) means that for every finite collection of covers $\pi_i : Y_i \dashrightarrow X$ of degree > 1 ,

$$X(k) \setminus \bigcup_i \pi_i(Y_i(k)) \quad (1)$$

is Zariski dense. It is a birational property, but it is much stronger than density of rational points.

Main theorem (Corvaja–Zannier, Theorems 1.4 and 1.6). *The Fermat quartic K3 surface*

$$F : x^4 + y^4 = z^4 + w^4 \subset \mathbf{P}^3 \quad (2)$$

is not unirational over $\mathbf{C}$ and has the Hilbert property over $\mathbf{Q}$.

Conversely, every projective variety X/k with the Hilbert property is algebraically simply connected: the normalization admits no unramified cover of degree > 1 . The fundamental group of every normal projective model of X has no subgroup of finite index > 1 .

The positive statement is striking because curves with HP are rational, whereas (2) is a non-unirational surface. The negative statement explains topologically why many varieties with dense rational points still fail HP.

Ingredient I: Chevalley–Weil is opposite to HP

If $Y \rightarrow X$ is a finite unramified cover, Chevalley–Weil lifts $X(k)$ to $Y(k')$ for one finite extension k'/k . Restriction of scalars and the diagonal $X \hookrightarrow \text{Res}_{k'/k} X$ replace this by finitely many degree- > 1 covers defined over k whose k -points cover $X(k)$. Thus an unramified cover makes (1) empty. Birational invariance and normalization give the second assertion of the theorem; Proposition 1.1 identifies these algebraic covers with finite-index subgroups of the topological fundamental group for a normal complex model.

Ingredient II: two elliptic pencils on the quartic

The surface F contains 48 lines, eight over $\mathbf{Q}$. Planes through $L = \{x = z, y = w\}$ cut F into L and a cubic E_λ , where

$$\lambda = \frac{w - y}{x - z}. \quad (3)$$

Two other rational lines give an origin and a non-torsion section on the generic cubic. Silverman specialization shows that $E_l(\mathbf{Q})$ has positive rank for all but finitely many $l \in \mathbf{Q}$, proving density. A second pencil E'_μ comes from $L' = \{x = w, y = -z\}$ and $\mu = (w - x)/(y + z)$. The second fibration is not cosmetic: one pencil alone cannot rule out finite unions of isogeny images on its elliptic fibres.

Suppose finitely many covers $Y_i \rightarrow F$ violate HP. Since the smooth K3 surface F is simply connected, every such finite cover has a branch curve. Restrict first to E_l . If a branch component is not contained in a λ -fibre, the restricted cover is generically ramified; its geometrically irreducible components have genus > 1 , so Faltings gives only finitely many rational lifts. If it is geometrically reducible, the same conclusion holds outside a thin set of parameters. Hence only covers whose branch curves are vertical for λ can remain relevant.

Ingredient III: isogeny cosets and the transverse pencil

For a remaining cover, $\pi^{-1}(E_l) \rightarrow E_l$ is generically an unramified cover of elliptic curves, hence a translate of an isogeny. The images of rational points are cosets of subgroups of the positive-rank group $E_l(\mathbf{Q})$. Lemma 3.2 says that if finitely many such cosets miss only finitely many points, then they miss none. Thus the assumed failure of HP would force every point of $E_l(\mathbf{Q})$ to lift, for every $l \in \mathbf{Q}$ outside one thin set T .

Now restrict the same covers to a general transverse fibre E'_m . Their vertical branch curves meet E'_m , so these restrictions are ramified and again have only finitely many rational lifts. The remaining issue is to find infinitely many $p \in E'_m(\mathbf{Q})$ with $\lambda(p) \notin T$. Lemma 3.3 proves that

$$\lambda|_{E'_m} : E'_m \longrightarrow \mathbf{P}^1 \quad (4)$$

has degree two and four ramification points whose branch values vary nontrivially with m . Choose m so these values avoid the branch loci of the finitely many covers of $\mathbf{P}^1$ defining T . Their fibre products with (4) then have genus at least two; Faltings makes $\lambda(E'_m(\mathbf{Q})) \cap T$ finite. The infinitely many remaining points contradict both the “every point lifts” conclusion on E_l and the finite-lifting conclusion on E'_m . This proves HP. Controlling this thin parameter set through moving ramification is the hardest step.

The Enriques counterpoint and limitations

The companion surface

$$x_0x_2^4 + x_1x_3^4 = x_0^2x_1^3 + x_0^3x_1^2 \quad (5)$$

has dense rational points and no nonconstant rational map to an abelian variety, yet fails HP (Theorem 1.3). A smooth model is Enriques and has an unramified double cover arising from adjoining $\sqrt{x_1/x_0}$; directly, the defining equation forces x_0x_1 at every rational point to be a square up to sign, so two double covers contain all rational points. This contrasts the simply connected K3 cover with its nonsimply-connected quotient.

The proof establishes HP for the specific quartic over $\mathbf{Q}$. The authors conjecture only the smooth-projective converse: a smooth algebraically simply connected projective variety with Zariski-dense rational points has HP; they explicitly warn that smoothness cannot be dropped. They also leave weak and weak-weak approximation for F open; HP alone does not settle either property.