Families of polynomials of every degree with no rational preperiodic points

Mohammad Sadek

arXiv:2010.09910

https://arxiv.org/abs/2010.09910

Eliminating every rational finite orbit

Let K be a number field with ring of integers $\mathcal{O}_K$, and fix an integer $d \geq 2$. A point of K is preperiodic for a polynomial $f \in K[x]$ when its forward orbit is finite; equivalently, one of its iterates is periodic. The Morton–Silverman conjecture predicts a uniform upper bound for the number of such points in terms of d and $[K : \mathbf{Q}]$. Sadek instead asks for entire one-parameter families having none at all.

For $P(T) \in \mathcal{O}_K[T]$ of degree N, put

$$Q(T, S) = S^N P(T/S).$$

Main theorem (Theorem 2.3)**.** *Assume that $d \mid N$, every root of P has multiplicity at most $d-1$, and $\mathrm{Gal}(P/K)$ contains an element that fixes no root of P. Then there is a nonzero $w \in \mathcal{O}_K$ such that, for every $t \in K$,*

$$f_t(x) = x^d + \frac{1}{wP(t)}$$

has no K-rational preperiodic point.

This removes the earlier restrictions that $K = \mathbf{Q}$ and that d be divisible by 2 or 3. It really yields infinitely many families: for fixed degree N, polynomials with Galois group S_N have density tending to one as their height grows (Remark 2.4); such a group contains a derangement, and the polynomial is separable.

Two inputs and one contradiction

The proof joins two results that at first look unrelated.

1. **An arithmetic shape forced by a cycle.** Proposition 2.1, due to Narkiewicz, says that if $x^d + c$ with $c \neq 0$ has a K-rational periodic point, then

$$c = \frac{a}{b^d} \quad (a, b \in \mathcal{O}_K), \qquad (a\mathcal{O}_K, b^d\mathcal{O}_K) = I^d$$

   for some ideal $I \subset \mathcal{O}_K$. Thus periodicity imposes a global d-th-power condition: the gcd ideal of the numerator and denominator is a d-th power.
2. **A twist with no rational point.** Legrand's theorem, applied using the root-multiplicity and Galois hypotheses, provides $w \neq 0$ for which the twisted superelliptic curve

$$y^d = wP(T),$$

   or homogeneously $y^d = wQ(T, S)$, has no nonzero K-rational homogeneous solution. The fixed-point-free Galois element is the obstruction that makes such a pointless twist available.

Now specialize $c = 1/(wP(t))$. If f_t had a periodic point, Proposition 2.1 would convert its required ideal-theoretic d-th-power presentation into a nontrivial K-point on $y^d = wQ(T, S)$. This contradicts the choice of w. Hence f_t has no periodic point of any period. A preperiodic point would eventually land on a periodic point, so there are no preperiodic points either. The same w works simultaneously for every $t \in K$.

What the reduction isolates

The argument can be viewed as a local-to-global denominator test. Iteration itself is compressed into Proposition 2.1: a finite orbit forces the constant term into a rigid ideal class. Homogenization then absorbs an arbitrary parameter $t = u/v$ into $Q(u, v)$ and turns that rigidity into precisely the rational-point equation forbidden by the twist. This is the useful toy model for the proof: rather than controlling every iterate, build a parameter space on which the mere existence of a cycle would be visible as one rational point, then choose a twist with no such point.

The hardest ingredient is therefore external to the elementary dynamical step: the existence theorem for pointless twists of superelliptic curves. Its hypotheses explain all three restrictions on P: $d \mid N$ makes $y^d = wQ(T, S)$ weighted-homogeneous of weighted degree N when y has weight N/d, root multiplicities below d rule out the relevant degeneracy, and the Galois derangement supplies the arithmetic obstruction.

The conclusion is deliberately negative and uniform in the parameter. It does not classify the dynamics over extensions of K, give a quantitative height bound, or construct the integer w explicitly in the proof. Nor does it prove the general uniform boundedness conjecture. What it supplies is a large source of degree-d families in which every K-rational finite orbit is excluded at once.

Codex mathematical paper digest, 2026-10-11. Source paper should be consulted for full statements and proofs.