

The Arakelov-Zhang pairing and Julia sets

Andrew Bridy, Matt Larson

arXiv:1906.02654

<http://arxiv.org/abs/1906.02654v2>

From a dynamical pairing to Julia geometry

Let K be a number field, M_K its places, and $r_v = [K_v : \mathbf{Q}_v]/[K : \mathbf{Q}]$. For a rational map $\phi \in K(x)$ of degree at least two, write h_ϕ for its canonical height and $\mu_{\phi,v}$ for its invariant probability measure on the Berkovich projective line over $\mathbf{C}_v$. The support of $\mu_{\phi,v}$ is the v -adic Julia set. The Arakelov–Zhang pairing $\langle \psi, \phi \rangle$ is a symmetric nonnegative adelic energy; for $\psi(x) = x^2$, its canonical height is the ordinary Weil height.

Main theorem (Bridy–Larson, Theorem 1.3). *For every such ϕ ,*

$$\langle x^2, \phi \rangle = h_\phi(0) - \sum_{v \in M_K} r_v \int_{|\alpha|_v < 1} \log |\alpha|_v d\mu_{\phi,v}(\alpha). \quad (1)$$

Only finitely many summands are nonzero. Consequently $\langle x^2, \phi \rangle \geq h_\phi(0)$, with equality exactly when the Julia set is disjoint from the Berkovich open unit disk at every place.

The correction in (1) is visibly nonnegative after the minus sign. It measures precisely how much canonical Julia mass lies inside the adelic unit disks, turning a global height pairing into local dynamical data.

Ingredient I: the local metric computation

The canonical metric of x^2 on $\mathcal{O}_{\mathbf{P}^1}(1)$ equals the standard metric at every place. In homogeneous coordinates choose $s = x_0$ and $t = x_0 + x_1$. For $a = [a : 1]$,

$$\|s(a)\|_{x^2,v} = \frac{|a|_v}{\max\{|a|_v, 1\}} = \min\{|a|_v, 1\}. \quad (2)$$

The defining local energy is therefore

$$\langle x^2, \phi \rangle_{s,t,v} = - \int_{|\alpha|_v < 1} \log |\alpha|_v d\mu_{\phi,v}(\alpha). \quad (3)$$

In the global pairing formula the two divisor terms are $h_\phi(\operatorname{div} s) = h_\phi(0)$ and $h_{x^2}(\operatorname{div} t) = h_{x^2}(-1) = 0$, which gives (1). At a good-reduction nonarchimedean place the canonical measure is the Dirac mass at the Gauss point, outside the open unit disk; this both explains finiteness and removes every such local term.

Ingredient II: why the equality criterion is exact

Because the Julia set is the support of $\mu_{\phi,v}$, disjointness from the unit disk makes the v -integral zero. Conversely, if the Julia set meets that disk, it meets a smaller open ball on which $\log |\alpha|_v \leq -\varepsilon$. Every neighborhood of a support point has positive measure, so the integral is strictly negative. Thus no cancellation between places is possible and the “if and only if” in the theorem is genuinely local place by place.

Ingredient III: the preimage-average proof

The paper also establishes, for nonexceptional β and $d = \deg \phi$,

$$\langle \psi, \phi \rangle = \lim_{n \rightarrow \infty} \frac{1}{d^n} \sum_{\phi^n(\alpha) = \beta} h_\psi(\alpha), \quad (4)$$

with multiplicities. Equidistribution of the preimages and the fact that μ_ϕ has no atoms show that no repeated preimage can carry a positive proportion of the average; the small-points theorem for the pairing then gives (4).

For a monic polynomial whose Julia set avoids 0 at every place, choose a finite set S containing the archimedean and bad-reduction places. Choose $\beta \in \mathcal{O}_{K,S}$ nonexceptional for ϕ and, for every $v \in S$, outside the v -adic closure of $\{\phi^n(0) : n \geq 0\}$. Factoring $\phi^n(x) - \beta$ converts the average Weil height in (4) into

$$\frac{h(\phi^n(0) - \beta)}{d^n} - \sum_{v \in S} r_v \frac{1}{d^n} \sum_{\substack{\phi^n(\alpha) = \beta \\ |\alpha|_v < 1}} \log |\alpha|_v + O(d^{-n}). \quad (5)$$

Here S -integrality removes every height contribution outside S , while the orbit-separation condition bounds $|\phi^n(0) - \beta|_v$ away from zero for $v \in S$ and justifies the $O(d^{-n})$ term. The first term tends to $h_\phi(0)$; preimage equidistribution turns the second into the integrals in (1). The technical restriction is essential for this proof: avoiding 0 makes $\log |\alpha|_v$ bounded and continuous on the Julia support. The adelic-metric proof above has no such restriction and applies to every rational map.

A concrete model and the rigidity consequence

If $\phi \in \mathbf{Z}[x]$ is monic, all nonarchimedean terms vanish, so

$$\langle x^2, \phi \rangle = h_\phi(0) - \int_{|z| < 1} \log |z| d\mu_\phi(z). \quad (6)$$

This is the clean toy model: the gap from the canonical height of 0 is exactly the logarithmic mass of the complex Julia set inside the unit disk. For the Chebyshev polynomial T_n the known arcsine measure on $[-2, 2]$ gives the degree-independent value

$$\langle x^2, T_n \rangle = \frac{3\sqrt{3}}{4\pi} L(2, \chi) \approx 0.3231, \quad (7)$$

where χ is the nontrivial character modulo 3.

The same mechanism proves Theorem 1.7. If $\phi = x^d + a_{d-1}x^{d-1} + \cdots + a_0 \in \mathcal{O}_K[x]$, the point 0 is preperiodic, and every archimedean Julia set avoids the open unit disk, then integrality gives good reduction at every nonarchimedean place. Hence $h_\phi(0) = 0$ and (1) yields $\langle x^2, \phi \rangle = 0$. Vanishing of the pairing identifies the canonical heights, and the Kawaguchi–Silverman classification of polynomials with equal canonical heights forces $\phi(x) = x^d$.

Scope

The formula also feeds explicit height-error estimates through a theorem of Petsche–Sziro–Tucker and is worked out for quadratic and linearly conjugate maps. Those computations are omitted here. Formula (4) is not claimed for an exceptional target, and the elementary proof of (1) is not claimed for general rational functions; the unrestricted result rests on the local adelic metric calculation.