

Unobstructed deformations for singular Calabi-Yau varieties

Robert Friedman

arXiv:2506.09857

<http://arxiv.org/abs/2506.09857v3>

A singular Bogomolov–Tian–Todorov theorem

In this paper a singular Calabi–Yau variety is a connected compact Gorenstein singular analytic space Y with $\omega_Y \simeq \mathcal{O}_Y$. The standing conventions further make Y reduced and normal, over $\mathbb{C}$, of dimension $n \geq 3$, with isolated singular locus Z . Let $\pi: \tilde{Y} \rightarrow Y$ be a good resolution with simple-normal-crossing exceptional divisor $E = \pi^{-1}(Z)$.

Main theorem (Friedman, Theorem 1.6). *Assume that*

- (i) *the $\partial\bar{\partial}$ -lemma holds on $\tilde{Y}$;*
- (ii) *every singularity of Y is isolated, local complete intersection (lci), and Du Bois;*
- (iii) *$H^1(Y, \mathcal{O}_Y) = 0$.*

Then the deformation functor Def_Y is unobstructed.

Thus the semiuniversal deformation base is smooth even though the usual global obstruction space T_Y^2 need not vanish. Compared with earlier results, Du Bois replaces stronger rational or weighted-homogeneous conditions, and the resolution need only satisfy the $\partial\bar{\partial}$ -lemma rather than be Kähler. Unobstructedness does not itself assert that Y admits a smoothing.

Ingredient I: relative Hodge theory supplies liftable classes

The complex

$$\Omega_{\tilde{Y}}^{\bullet}(\log E)(-E)$$

computes $H^*(\tilde{Y}, E; \mathbb{C}) \simeq H^*(Y, Z; \mathbb{C})$. The $\partial\bar{\partial}$ hypothesis gives E_1 -degeneration of its Hodge-to-de Rham spectral sequence. The Du Bois condition identifies the degree-zero term after pushforward, while lci singularities make Ω_Y^1 reflexive and give

$$\Omega_Y^1 \simeq \pi_* \Omega_{\tilde{Y}}^1(\log E)(-E). \quad (1)$$

Leray and Steenbrink vanishing then compare the degree-one pieces on Y and $\tilde{Y}$ (Lemmas 2.2–2.3).

For the two-term complex $C_Y^{\bullet} = \{\mathcal{O}_Y \xrightarrow{d} \Omega_Y^1\}$, these comparisons and $H^{n-1}(Y, \mathcal{O}_Y) = 0$ imply

$$H^{n-1}(Y; \mathbb{C}) \twoheadrightarrow \mathbb{H}^{n-1}(Y, C_Y^{\bullet}) \simeq H^{n-2}(Y, \Omega_Y^1). \quad (2)$$

Here the required H^{n-1} -vanishing follows from hypothesis (iii) by Serre duality and $\omega_Y \simeq \mathcal{O}_Y$. The delicate point is that (2) is obtained through relative cohomology on the resolution, so classes do not acquire uncontrolled contributions at the singular set.

Ingredient II: the same surjectivity persists in families

Let A be an Artin local $\mathbb{C}$ -algebra and $\mathcal{Y} \rightarrow \operatorname{Spec} A$ a deformation of Y . Replace the second term of $C_Y^\bullet$ by $\Omega_{\mathcal{Y}/A}^1$. Constant cohomology classes factor through this relative complex, so (2) gives Lemma 2.6:

$$H^{n-2}(\mathcal{Y}, \Omega_{\mathcal{Y}/A}^1) \rightarrow H^{n-2}(Y, \Omega_Y^1). \quad (3)$$

Flatness away from the isolated singularities and formal base change upgrade (3) to Theorem 2.1: the corresponding map remains surjective after every quotient $A \twoheadrightarrow A'$.

Ingredient III: surjectivity kills every lifting obstruction

Put $A_k = \mathbb{C}[t]/(t^{k+1})$. The T^1 -lifting criterion (Theorem 3.3) reduces unobstructedness to surjectivity of

$$T^1(Y_k/A_k) \longrightarrow T^1(Y_{k-1}/A_{k-1}) \quad (4)$$

for every infinitesimal deformation Y_k . Lemma 3.4 identifies these sets with $\operatorname{Ext}^1(\Omega_{Y_k/A_k}^1, \mathcal{O}_{Y_k})$. Du Bois singularities also ensure $\omega_{Y_k/A_k} \simeq \mathcal{O}_{Y_k}$ (Lemma 3.1), so relative duality applies throughout the tower.

Gross's duality diagram, recorded as Theorem 3.9, identifies failure of (4) with the boundary following

$$H^{n-2}(Y, \Omega_Y^1) \longrightarrow H^{n-1}(Y_{k-1}, \Omega_{Y_{k-1}/A_{k-1}}^1).$$

Equivalently, it is enough that

$$H^{n-2}(Y_k, \Omega_{Y_k/A_k}^1) \longrightarrow H^{n-2}(Y, \Omega_Y^1) \quad (5)$$

be surjective. Lemma 2.6 is exactly (5); hence every map (4) is surjective, and the T^1 -lifting criterion proves Theorem 1.6. The decisive bridge is therefore not vanishing of T_Y^2 , but persistence of one Hodge-theoretic quotient under every infinitesimal thickening.

A nodal model and the extensions

If Y is a Calabi–Yau threefold with only ordinary double points, the singularities are isolated rational lci and hence Du Bois. With a Kähler (therefore $\partial\bar{\partial}$) resolution and $H^1(Y, \mathcal{O}_Y) = 0$, the theorem recovers the classical nodal unobstructedness result. The model also shows the limitation: smoothing individual nodes is a separate question about the image of global first-order deformations in the local deformation spaces.

The paper goes further in three directions. Without lci, Theorem 3.10 proves only a *generalized* T^1 -lifting property, with the local-cohomology image $\operatorname{Im} H_Z^{n-2}(Y, \Omega_Y^1)$ divided out; together with prorepresentability it gives the structural conclusions of Theorem 3.8, not blanket unobstructedness. Theorem 3.15 gives the analogous unobstructedness statement for lci log Calabi–Yau pairs, assuming the $\partial\bar{\partial}$ -lemma also on all boundary strata. Finally, Theorem 3.17 proves unobstructedness when Y has isolated lci rational singularities and ω_Y^{-1} is nef and big, using a cyclic cover, logarithmic T^1 lifting, and vanishing to forget the boundary.