

# A Bogomolov property for the canonical height of maps with superattracting periodic points

Nicole R.Looper

arXiv:2106.13003

<http://arxiv.org/abs/2106.13003v2>

## A dynamical Bogomolov theorem

For a degree- $d$  polynomial  $f$  over a product-formula field  $K$ , write  $\widehat{h}_f$  for its canonical height and  $K^{\text{ab}}$  for the maximal abelian extension of  $K$ . A finite periodic point  $p_0$  is *superattracting* when  $(f^m)'(p_0) = 0$  for its period  $m$ . At a nonarchimedean place  $v$ , bad reduction is detected dynamically: the Berkovich filled Julia set  $\mathcal{K}_v$  is not a single disk.

**Main theorem** (Looper, Theorem 1.5). *Let  $K$  be a product-formula field and  $f \in K[z]$ . Suppose that  $f$  has a finite superattracting periodic point and a nonarchimedean place of bad reduction. Then some  $\varepsilon > 0$  satisfies*

$$\#\{P \in K^{\text{ab}} : \widehat{h}_f(P) < \varepsilon\} < \infty. \quad (1)$$

*In particular,  $f$  has only finitely many preperiodic points in  $K^{\text{ab}}$ .*

The hypotheses force  $d \geq 3$ . Thus (1) gives a genuine height gap over an infinite extension, where Northcott finiteness is unavailable; it is the polynomial-dynamical analogue of a Bogomolov property for canonical heights on abelian varieties.

## The local “splash” and its arithmetic consequence

After a finite extension and affine conjugacy, the proof makes  $f$  monic. The local construction is presented after passing to a suitable iterate so that one point  $p_0$  of the superattracting cycle is fixed. Let  $B$  be the largest Berkovich disk about  $p_0$  contained in  $\mathcal{K}_v$ . Proposition 3.1 constructs disks  $B_i \subset \mathcal{K}_v$  outside  $B$  whose distances to  $p_0$  decrease to the radius of  $B$ . These side branches are the nonarchimedean version of a “splash” from the immediate attracting basin.

Suppose, contrary to (1), that distinct  $P_i \in K^{\text{ab}}$  satisfy  $\widehat{h}_f(P_i) \rightarrow 0$ , and let  $L_i$  be the Galois closure of  $K(P_i)/K$ . Small-point equidistribution on  $\mathcal{K}_v$  sees every splash disk. Corollary 3.2 therefore gives a subsequence for which the local decomposition groups above  $v$  have orders tending to infinity. Corollary 3.3 strengthens this: for every disk  $B' \supsetneq B$ , a fixed positive proportion of the global conjugates lies in  $B'$ .

Here abelianity is decisive. All decomposition groups of  $L_i/K$  above  $v$  coincide. Choose a prime  $\mathfrak{Q}_i$  of  $L_i$  above  $v$ ; whenever  $P_i$  lies in  $B'$  at one completion, its whole local orbit

$$T_i = \{\sigma(P_i) : \sigma \in \text{Decomp}(\mathfrak{Q}_i \mid v)\} \quad (2)$$

lies in  $B'$ . Choose  $B'$  with logarithmic capacity  $\log \gamma_v(B') \leq \log \gamma_v(\mathcal{K}_v) - \eta$ . The product formula for the discrete transfinite diameters of  $T_i$  gives a global sum equal to zero. The positive proportion of completions controlled by (2) contributes at most  $-\eta$  times a fixed positive local weight, while equidistribution bounds every remaining finite or infinite-place error by  $O(\varepsilon')$ . Taking  $\varepsilon'$  sufficiently small makes the same sum strictly negative, the required contradiction. Synchronizing local clustering across all primes above  $v$  is the hardest step and explains exactly why the extension must be abelian.

## A second theorem cluster: uniformity in $\mathcal{S}_{d,m}$

Fix  $d \geq 3$  and  $m \geq 1$ , and let  $\mathcal{S}_{d,m}(K)$  be the degree- $d$  polynomials over  $K$  with a finite superattracting point of period  $m$ . With  $M_K$  the place set,  $M_K^0$  and  $M_K^\infty$  its nonarchimedean and archimedean parts,  $S_d$  the places above prime integers at most  $d$ ,  $r_v$  the normalized local weight, and  $\widehat{\lambda}_v$  the local escape rate, set

$$\lambda_{\text{crit},v}(f) = \max_{a \text{ finite critical}} \widehat{\lambda}_v(a), \quad h_{\text{crit}}(f) = \sum_{v \in M_K} r_v \lambda_{\text{crit},v}(f). \quad (3)$$

For  $K$  a number field or a one-dimensional function field of characteristic zero, Theorems 1.7–1.8 say:

- assuming the *abc* conjecture when  $K$  is a number field, and assuming non-isotriviality when  $K$  is a function field, there is  $B(d, m, K)$  bounding the number of  $K$ -rational preperiodic points of every  $f \in \mathcal{S}_{d,m}(K)$ ;
- assuming *abc* only in the number-field case, some  $\kappa(d, m, K) > 0$  satisfies

$$\widehat{h}_f(P) \geq \kappa \max\{1, h_{\text{crit}}(f)\} \quad (4)$$

for every non-preperiodic  $P \in K$ .

The maximum with 1 in (4) is essential: a postcritically finite map can still have bad reduction.

## How equidistribution meets *abc*

Theorem 2.4 is a uniform quantitative equidistribution statement. A large finite set  $T \subset K$  whose average canonical height is sufficiently small relative to  $h_{\text{crit}}(f)$  distributes correctly at a fixed depth around  $p_0$ . If  $E$  is the smallest disk containing  $\mathcal{K}_v$  and  $E_1 = f^{-1}(E)$ , it also distributes across every wing decomposition of  $E_1$ , on a  $\delta$ -slice  $S$  of the bad places. Here “slice” is weighted, not counted: for a set  $\Sigma \subset M_K^0 \setminus S_d$  of bad places,

$$\sum_{v \in S} r_v \lambda_{\text{crit},v}(f) \geq \delta \sum_{v \in \Sigma} r_v \lambda_{\text{crit},v}(f). \quad (5)$$

The constants depend only on the displayed family parameters, not on  $f$ .

Three reductions turn this analytic input into Diophantine leverage. First, nested pullback annuli about  $p_0$  have moduli decaying geometrically; Newton polygons show that the denominators of their radii grow uniformly (Proposition 3.4). Second, equidistribution forces a positive proportion of pairs  $(P_i, P_j) \in T^2$  to have a large aggregate separation over a set carrying positive critical-height weight (Lemma 3.5). Third, let  $S_{\text{good}}$  be the good-reduction places. Most differences  $\alpha = P_i - P_j$  are  $\varepsilon$ -good (Proposition 4.1), meaning

$$\sum_{v \in S_d \cup M_K^\infty \cup S_{\text{good}}} r_v \log |\alpha^{-1}|_v \leq \varepsilon h_{\text{crit}}(f); \quad (6)$$

this controls the small-prime, archimedean, and good-reduction contribution to the radical.

Choose three pairwise distinct points, translate one to 0, and pass after bounded field extension to a monic conjugate. Applying *abc* to

$$(-P_1, P_2, P_1 - P_2), \quad -P_1 + P_2 + (P_1 - P_2) = 0, \quad (7)$$

the annular denominators make the bad-place differences too “powerful”: height minus radical is a fixed positive proportion of  $h_{\text{crit}}(f)$ . This contradicts *abc* when the critical height is large. Bounded critical height is handled by the critical-height/moduli-height comparison and Northcott in the number-field case, and by the corresponding function-field finiteness and isotrivial alternatives. Passing to a suitable iterate reduces period  $m$  to the fixed-point case.

## Toy picture, roadmap, and scope

Imagine the nested disks around  $p_0$  as a trunk, with preimage disks accumulating along its boundary as side branches. A small-height Galois orbit must sample the branches. The abelian-extension theorem says that abelian local symmetry forces too much of the orbit into one disk of smaller capacity; the uniformity theorems instead compare many branch separations and feed them to *abc*. Symbolically,

$$\begin{aligned} &\text{superattracting cycle} + \text{bad reduction} \longrightarrow \text{nested annuli and splash disks} \longrightarrow \text{small-point equidistribution} \\ &\quad \left\{ \begin{array}{l} \text{abelian decomposition groups} \rightarrow \text{capacity contradiction,} \\ \text{many powerful differences} \rightarrow \text{abc contradiction.} \end{array} \right. \quad (8) \end{aligned}$$

Theorem 1.5 does not assert a uniform value of  $\varepsilon$  over a family. The bounds in Theorems 1.7–1.8 are conditional over number fields, and the paper treats polynomials with a finite superattracting cycle rather than arbitrary rational maps.