Moduli spaces of rational curves on Artin-Mumford double solids

Fumiya Okamura

Fumiya Okamura, arXiv:2501.09269v2 (2026), 20 pages

http://arxiv.org/abs/2501.09269v2

Four components in every degree

Work over an algebraically closed field k of characteristic 0. Let $W \simeq \mathbf{P}^3$ be a general 3-dimensional linear system of quadrics in $\mathbf{P}^3$, with ten-nodal quartic discriminant D_W. The branched double cover $f : X \to W$ is an *Artin–Mumford double solid*, a factorial terminal del Pezzo threefold of degree 2; let H generate $\mathrm{Pic}(X) \simeq \mathbf{Z}$.

Main theorem (Okamura, Theorem 1.1)**.** *The space* $\mathrm{Mor}(\mathbf{P}^1, X, 1)$ *of* H*-lines has two irreducible components* M_1^+ *and* M_1^-*, each of expected dimension* 5*. For every* $d \geq 2$*, the space* $\mathrm{Mor}(\mathbf{P}^1, X, d)$ *has exactly four irreducible components*

$$R_d^+, \quad R_d^-, \quad N_d^+, \quad N_d^-,$$

all of expected dimension $2d + 3$*. The component* $N_d^\pm$ *parametrizes* d*-sheeted covers of lines in* $M_1^\pm$*, whereas* $R_d^\pm$ *generically parametrizes embedded very free curves. The same classification holds for* $\overline{M}_{0,0}(X, d)$*, with every expected dimension lowered by* 3*.*

Precisely $R_d^\pm$ are Manin components; their number matches $|\mathrm{Br}_{\mathrm{nr}}(k(X)/k)| = 2$ (Corollary 4.12). To the author's knowledge, this is the first Fano example with multiple Manin components in every $\mathrm{Mor}(\mathbf{P}^1, X, d)$, $d \geq 2$.

Ingredient 1: Reye congruences split the lines

A line $L \subset W$ lifts to two H-lines exactly when it is bitangent to D_W. Call W *excellent* when D_W contains no lines; a general web is excellent. The bitangent surface is normalized by the Reye congruence $\mathrm{Rey}(W)$, an Enriques surface. Its incidence construction lifts the normalization to one component:

$$\mathrm{Rey}(W) \longrightarrow M_1^+ \longrightarrow \mathrm{Bit}(D_W).$$

The deck involution produces M_1^-; the generic two-point fiber shows there are no other line components (Theorem 4.3).

Ingredient 2: the conic base case detects two signs

Movable bend-and-break reduces a birational conic component to two free lines. The sign-type boundary loci $\Delta^{++}, \Delta^{+-}, \Delta^{--}$ are irreducible because so are the incidence fiber products $(M_1^\epsilon)' \times_X (M_1^{\epsilon'})'$.

On the blow-up $\widetilde{X}$, $\mathrm{Br}(\widetilde{X}) \simeq \mathbf{Z}/2\mathbf{Z}$ manifests as two algebraic-equivalence classes in one numerical class, separating the mixed-sign boundary from the same-sign ones (Lemma 3.6). A general H-section $S \in |H|$ is a degree-2 del Pezzo surface with 56 lines; Lemma 2.12 gives a conic bundle with reducible fibers of types $++$ and $--$, joining those strata. If $R^{\epsilon\epsilon'}$ denotes the unique birational component containing $\Delta^{\epsilon\epsilon'}$, then $R^{++} = R^{--} =: R_2^+$ while $R^{+-} =: R_2^-$, and the two double-cover loci $N_2^\pm$ complete the four conic components (Theorem 4.9).

Thus the Brauer class creates no new numerical degree but splits each degree's moving families into two deformation types.

Ingredient 3: bend-and-break and induction

General del Pezzo-threefold results make every stable-map component dominant of expected dimension and give a chain of free lines. Theorem 4.10 strengthens this: every birational component has free two-component degenerations displaying a line from each $M_1^\pm$. The conic relation changes a same-sign pair to the opposite sign, then the remaining chain is smoothed.

Inductively attach a marked $M_1^\pm$-line to an $R_{d-1}^\pm$-curve. Irreducible evaluation fibers make the four boundary loci irreducible; Theorem 4.10 pairs them into two components. Their strict transforms on $\widetilde{X}$ are not algebraically equivalent, so they cannot merge. Dimension separates $N_d^\pm$, and general members of $R_d^\pm$ are embedded and very free.

Only $R_d^\pm$ survive the Manin test; the line and multiple-cover loci are accumulating through families of H-lines. The hard point is to force every birational component to the free-line boundary while the Brauer sign prevents the last two components from coalescing.

Codex mathematical paper digest, 2026-10-19. Source paper should be consulted for full statements and proofs.