

Gorenstein spherical Fano varieties

Giuliano Gagliardi, Johannes Hofscheier

arXiv:1303.2994

<http://arxiv.org/abs/1303.2994v4>

Reflexive polytopes with colors

Fix a connected reductive complex group G , a Borel subgroup B , and a spherical homogeneous space G/H . Let M be the lattice of weights of B -semi-invariant rational functions, $N = \text{Hom}(M, \mathbb{Z})$, and $\mathcal{V} \subset N_{\mathbb{Q}}$ the valuation cone. The B -invariant prime divisors of G/H are its *colors* $D \in \mathcal{D}$; their valuations give $\rho(D) \in N$.

Brion's distinguished anticanonical section has divisor

$$\sum_{D \in \mathcal{D}} m_D D + \sum_j X_j, \quad (1)$$

where $m_D > 0$ depends only on G/H and the X_j are the G -invariant prime divisors of an embedding $G/H \hookrightarrow X$. For a complete spherical embedding set

$$Q_X = \text{conv}(\{\rho(D)/m_D : D \in \mathcal{D}\} \cup \{\nu_{X_j}\}_j) \subset N_{\mathbb{Q}}. \quad (2)$$

A polytope Q is G/H -reflexive when: it contains every $\rho(D)/m_D$; $0 \in \text{int } Q$; every vertex is either one of those color points or a lattice point in $\mathcal{V}$; and every *supported* vertex v of Q^* lies in M . Here supported means that the relative interior of the cone over the dual facet $\hat{v}$ meets $\mathcal{V}$.

Main theorem (Gagliardi–Hofscheier, Theorems 1.9 and 1.10). *The assignment $X \mapsto Q_X$ is a bijection between isomorphism classes of Gorenstein spherical Fano embeddings of G/H and G/H -reflexive polytopes. Moreover, if X is a d -dimensional $\mathbb{Q}$ -factorial Gorenstein spherical Fano variety, then*

$$\rho_X \leq 2d. \quad (3)$$

Equality holds exactly when d is even and $X \cong (S_3)^{d/2}$, where S_3 is the blowup of $\mathbb{P}^2$ at three non-collinear points.

The first statement extends the toric reflexive-polytope dictionary to all spherical varieties; the second shows that its combinatorics retains the sharp toric Picard bound.

Ingredient I: the anticanonical divisor supplies the vertices

Theorem 1.2 constructs the section in (1) with coefficient 1 on every G -invariant boundary divisor. Theorem 1.5 computes the color coefficients: $m_D = 1$ for colors of types a and $2a$, while for type b one has $m_D = \langle \alpha^\vee, \kappa_P \rangle \geq 2$. Thus (2) remembers the complete anticanonical divisor with the correct normalization. Omitting the factors m_D would destroy the Cartier/Fano criterion.

Ingredient II: reconstruct the embedding from faces

First ignore integrality on Q^* . A $\mathbb{Q}$ - G/H -reflexive polytope determines its supported colored face fan $\mathfrak{F}_{\text{supp}}(Q)$. On a maximal cone corresponding to $v \in Q^*$, define

$$\psi|_{C \cap \mathcal{V}} = -\langle \cdot, v \rangle.$$

This is the piecewise-linear function of $-K_X$. Convexity of Q makes ψ strictly convex, while inclusion of the color points supplies the strict color inequalities; the spherical ampleness criterion therefore makes $-K_X$ ample. Conversely, the support function of an anticanonical $\mathbb{Q}$ -Cartier Fano embedding recovers exactly the facets of Q_X . Propositions 7.2–7.4 show that these constructions are inverse.

Ingredient III: only supported dual vertices test Cartierness

Not every face cone of Q corresponds to a spherical orbit: only the cones meeting $\mathcal{V}$ in relative interior are supported. Proposition 8.3 detects this intrinsically by

$$v \text{ supported} \iff Q^* \cap (v + \text{cone}(\Sigma)) = \{v\}, \quad (4)$$

where Σ is the set of spherical roots. On the corresponding cone, $-K_X$ is represented by the linear form $-\langle \cdot, v \rangle$. Hence it is Cartier precisely when every supported v belongs to M (Theorem 8.5). This is the extra condition upgrading the $\mathbb{Q}$ -Gorenstein dictionary to the Gorenstein one.

Ingredient IV: count valuation vertices and colors

Write $r = \text{rank}(G/H) = \dim N_{\mathbb{Q}}$ and let S^p be the simple roots of the stabilizer of the open B -orbit. Under $\mathbb{Q}$ -factoriality, each supported facet has exactly r vertices of Q , and no such vertex can equal $\rho(D)$ for two colors (Proposition 12.1). These vertices need not all lie in $\mathcal{V}$. After enlarging Q without changing its already simplicial facets, adjacent-facet counting bounds by r the vertices in $\text{Vert}(Q) \cap \mathcal{V} \cap N$ pairing to 0 with a fixed supported dual vertex.

Lemma 12.4 puts 0 in the interior of the convex hull of the supported dual vertices together with $-\Sigma$. A positive integral dependence among them, summed over valuation vertices, then gives

$$|\text{Vert}(Q) \cap \mathcal{V} \cap N| \leq 3r. \quad (5)$$

If r' is the number of G -invariant prime divisors, the spherical Picard formula and the color/root bounds yield

$$\rho_X = r' + |\mathcal{D}| - r \leq 2r + |\mathcal{D}| \leq 2r + 2|S \setminus S^p| \leq 2d, \quad (6)$$

which proves (3). Equality forces every inequality to be sharp. The root system outside S^p becomes a product of A_1 's and Proposition 13.5 forces $\Sigma = \emptyset$; thus X is horospherical, where Pasquier's equality classification gives $(S_3)^{d/2}$.

Toric model and scope

For a torus there are no colors and $\mathcal{V} = N_{\mathbb{Q}}$, so every dual vertex is supported. The definition reduces to an ordinary reflexive lattice polytope, and $\mathbb{Q}$ -factoriality reduces to simpliciality. This toy case displays both modifications required in the spherical world: fixed color vertices and a Cartier test only on supported facets.

The classification is for embeddings of a fixed G/H over $\mathbb{C}$; the Picard bound additionally assumes Gorenstein and $\mathbb{Q}$ -factorial. Its technical bottleneck is controlling non-simplicial unsupported facets without losing the supported ones needed for Cartierness and the vertex count.