A dynamical pairing between two rational maps

Clayton Petsche, Lucien Szpiro, Thomas J. Tucker

arXiv:0911.1875v1 (2009), 22 pages

http://arxiv.org/abs/0911.1875v1

A distance between two arithmetic dynamical systems

Let K be a number field and let $\varphi, \psi : \mathbf{P}^1 \to \mathbf{P}^1$ be rational maps, both defined over K and of degree at least two. Write h_φ, h_ψ for their Call–Silverman canonical heights and $\mathrm{PrePer}(\varphi)$ for the φ-preperiodic points. The paper defines a symmetric global Arakelov–Zhang pairing $\langle \varphi, \psi \rangle$ by summing local energy pairings of the canonical adelic metrics on $\mathcal{O}(1)$.

Main theorem (Petsche–Szpiro–Tucker, Theorems 1–4)**.** *If $x_n \in \mathbf{P}^1(\overline{K})$ are distinct and $h_\psi(x_n) \to 0$, then*

$$h_\varphi(x_n) \longrightarrow \langle \varphi, \psi \rangle.$$

If $d = \deg \psi$, the same number is the periodic-point average

$$\langle \varphi, \psi \rangle = \lim_{n\to\infty} \frac{1}{d^n + 1} \sum_{\psi^n(x) = x} h_\varphi(x),$$

where fixed points are counted with multiplicity. Moreover the pairing is nonnegative, and the following are equivalent:

$$\langle \varphi, \psi \rangle = 0, \qquad h_\varphi = h_\psi, \qquad \mathrm{PrePer}(\varphi) = \mathrm{PrePer}(\psi),$$
$$\#\big(\mathrm{PrePer}(\varphi) \cap \mathrm{PrePer}(\psi)\big) = \infty, \qquad \liminf_x \big(h_\varphi(x) + h_\psi(x)\big) = 0.$$

Finally, for $\sigma(x) = x^2$ and every $x \in \mathbf{P}^1(\overline{K})$,

$$h_\psi(x) - h_{\mathrm{st}}(x) \le \langle \sigma, \psi \rangle + h_\psi(\infty) + \log 2.$$

Thus the pairing is the limiting φ-height seen by points dynamically small for ψ; symmetry makes the visibly asymmetric periodic-point limit symmetric after interchanging the maps.

Ingredient 1: local potentials assemble globally

At every place v, the canonical metric gives a continuous potential and probability measure $\mu_{\varphi,v}$ on the Berkovich projective line. The measure-valued Laplacian and integration by parts make the local pairing symmetric. Outside finitely many places the metrics are standard and the local terms vanish; the product formula makes the global sum independent of auxiliary sections and finite field extensions. The metric functional equation also makes $\langle \varphi, \varphi \rangle$ equal to $e\langle \varphi, \varphi \rangle$ for $e = \deg \varphi$, hence zero.

Ingredient 2: equidistribution turns energy into height

Let Z_n be finite Galois-invariant multisets that are h_ψ-small and equidistribute to $\mu_{\psi,v}$ at every place. Comparing the two canonical metrics gives continuous functions f_v, nonzero at only finitely many places. Equidistribution turns the average of $h_\varphi - h_\psi$ into $\sum_v r_v \int f_v \, d\mu_{\psi,v} = \langle \varphi, \psi \rangle$. Galois orbits of distinct ψ-small points supply Theorem 1; the multiset of solutions of $\psi^n(x) = x$ supplies Theorem 2. Nonnegativity then follows because $h_\varphi \ge 0$.

Ingredient 3: common small points force identical heights

An infinite common preperiodic set gives a sequence small for both maps. Its Galois orbits equidistribute to both local measures, forcing them to agree. The two potentials then differ by constants, so globally $h_\varphi = h_\psi + c$. Evaluation at a φ-preperiodic point and at a ψ-preperiodic point gives $c \le 0$ and $c \ge 0$; hence $c = 0$. This is the decisive implication in the five-way criterion.

Height comparison, toy model, and scope

For $\sigma(x) = x^2$, one has $h_\sigma = h_{\mathrm{st}}$. A generalized Mahler formula expresses $h_\psi(x) - h_\psi(\infty)$ through local averages of $\log|x-\alpha|_v$ over the finite points $\psi^n(\alpha) = \alpha$, counted with multiplicity. The bound $\log|x-\alpha|_v \le \log^+|x|_v + \log^+|\alpha|_v + \theta_v$, where $\theta_v = \log 2$ at archimedean places and 0 otherwise, followed by the periodic-point formula yields Theorem 4. The family $\psi_c(x) = x^2 + c$ is the concrete model: the paper proves $\langle \sigma, \psi_c \rangle = \frac{1}{2} h_{\mathrm{st}}(c) + O(1)$, so the pairing records the arithmetic size of the deformation away from squaring.

The hardest step is the compatible local-to-global normalization across all places. The results concern degree-at-least-two maps on $\mathbf{P}^1$ over a number field; the last estimate is one-sided, contains $h_\psi(\infty)$, and does not claim that $\log 2$ is optimal.

Codex mathematical paper digest, 2026-10-25. Source paper should be consulted for full statements and proofs.