

A rational map with infinitely many points of distinct arithmetic degrees

John Lesieutre, Matthew Satriano

arXiv:1809.00047

<http://arxiv.org/abs/1809.00047v1>

How finiteness of arithmetic degrees fails

Let $f : X \dashrightarrow X$ be a dominant rational self-map of a smooth projective variety over $\overline{\mathbf{Q}}$. Write $X_f(\overline{\mathbf{Q}})$ for the points whose forward orbits avoid the indeterminacy locus. For an ample height h_H , the arithmetic degree (when the limit exists) is

$$\alpha_f(P) = \lim_{m \rightarrow \infty} \max\{h_H(f^m(P)), 1\}^{1/m}.$$

It measures exponential height growth and is independent of H . Kawaguchi and Silverman had conjectured, among other things, that the set of values $\alpha_f(P)$ is finite. The paper isolates the failure of that finiteness assertion for rational maps.

Main theorem (Theorem 2). *The birational map $f : \mathbf{P}^4 \dashrightarrow \mathbf{P}^4$ defined by*

$$[X : Y : Z : A : B] \mapsto [XY + AX : YZ + BX : XZ : AX : BX]$$

admits a sequence $P_n \in X_f(\overline{\mathbf{Q}})$ such that every $\alpha_f(P_n)$ exists and the set $\{\alpha_f(P_n) : n \geq 10\}$ is infinite.

The construction deliberately preserves the parameters (a, b) . On the affine chart it is the family of plane birational maps

$$f_{a,b}(x, y) = (y + a, y/x + b).$$

Thus one ambient fourfold map packages infinitely many surface dynamics; special fibers will have different exponential growth rates.

A fiberwise transfer principle

Proposition 3 gives the conceptual bridge. Suppose a birational map over a parameter space has surface fibers $f_{t_n} : X_{t_n} \dashrightarrow X_{t_n}$ with three properties: each lifts on a birational model $\tilde{X}_{t_n}$ to an automorphism $\tilde{f}_{t_n}$; there is a point P_n in the common isomorphism locus with a well-defined, Zariski-dense orbit; and the first dynamical degrees $\lambda_1(f_{t_n})$ take infinitely many values. If $\tilde{P}_n$ denotes the unique lift of P_n under $\pi_n : \tilde{X}_{t_n} \rightarrow X_{t_n}$, then

$$\alpha_f(P_n) = \alpha_{f_{t_n}}(P_n) = \alpha_{\tilde{f}_{t_n}}(\tilde{P}_n) = \lambda_1(\tilde{f}_{t_n}) = \lambda_1(f_{t_n}).$$

Here restriction of an ambient ample height gives the first equality; arithmetic degree and dynamical degree are preserved through the indicated birational lift; and the middle equality is the known arithmetic-degree theorem for a surface automorphism at a point with dense orbit. Consequently, varying fiberwise dynamical degrees become varying arithmetic degrees for one map.

The special surface fibers

The Bedford–Kim and McMullen construction, assembled in Proposition 4, supplies rational parameters $t_n = (a_n, b_n)$ for every $n \geq 10$. Put

$$p_n(T) = T^{n-2}(T^3 - T - 1) + T^3 + T^2 - 1.$$

Then $\lambda_1(f_{t_n}) = \delta_n$, the largest real root of p_n , and the numbers δ_n increase strictly to the real root $\delta_* \simeq 1.32472$ of $T^3 - T - 1$. In particular they are infinitely many distinct values, all greater than one.

Each f_{t_n} preserves a cuspidal cubic $C_{t_n} \subset \mathbf{P}^2$. Blowing up n specified points in the smooth locus of this cubic produces $\pi_n : \tilde{X}_{t_n} \rightarrow \mathbf{P}^2$ on which f_{t_n} extends to an automorphism. The cusp q_n is not blown up and avoids the indeterminacy locus. In suitable local coordinates,

$$Df_{t_n}(q_n) = \begin{pmatrix} \delta_n^{-2} & 0 \\ 0 & \delta_n^{-3} \end{pmatrix},$$

so q_n is an attracting fixed point. These facts simultaneously provide the automorphism models, the infinite list of dynamical degrees, and a safe analytic region in which to build the required algebraic points.

Why dense algebraic orbits exist

Because $\delta_n > 1$, the lifted surface automorphism has positive dynamical degree and only finitely many periodic curves. Take a small complex analytic basin U_n around q_n such that $f_{t_n}(U_n) \subset U_n$, contraction is strict, U_n misses the indeterminacy locus, and π_n is an isomorphism above it. The cusp is the only periodic point in this basin.

Algebraic points are analytically dense, so one may choose $\tilde{P}_n \in \pi_n^{-1}(U_n)(\overline{\mathbf{Q}})$ outside the finite union of periodic curves. It is not periodic, and its orbit stays in the isomorphism locus. If its orbit closure on the surface were proper, its positive-dimensional part would yield a periodic curve; hence the orbit is Zariski dense. The image $P_n = \pi_n(\tilde{P}_n)$ therefore satisfies every hypothesis of the transfer principle, giving $\alpha_f(P_n) = \delta_n$.

A useful model is a family of linear-fractional recurrences whose spectral radius drops at special parameter values: a point with generic growth on each special fiber records that fiber's rate. The actual argument must do more. It replaces the recurrence by a birational surface automorphism, controls indeterminacy through blowups, and uses the attracting cusp to find algebraic points whose orbits are both globally dense and locally trapped.

Scope and hardest point

The example preserves the fibration by (a, b) , and the chosen points are not asserted to have bounded degree over $\mathbf{Q}$. Accordingly, the paper leaves open corrected finiteness statements excluding invariant fibrations on every invariant subvariety or imposing a uniform field-degree bound. It also does not refute existence of the arithmetic degree: existence is proved for every point used in the counterexample.

The hardest step is not writing the fourfold map but reconciling three requirements on each special fiber: an infinite and explicitly varying dynamical spectrum, an automorphism model to which the arithmetic-degree theorem applies, and an algebraic dense orbit that never meets indeterminacy. The invariant cubic and its attracting cusp are what make those requirements compatible.