

Quasi-projectivity of the moduli space of smooth Kähler-Einstein Fano manifolds

Chi Li, Xiaowei Wang, Chenyang Xu

arXiv:1502.06532v1

<http://arxiv.org/abs/1502.06532v1>

A positive CM line bundle across the boundary

Let $\mathcal{M}$ be the moduli space of smooth Kähler–Einstein Fano manifolds of fixed dimension, and let $\overline{\mathcal{M}}$ be the proper good moduli space of smoothable K-polystable $\mathbf{Q}$ -Fano varieties constructed in the authors’ earlier work. The CM line bundle is initially defined on a Hilbert parameter space. A fixed tensor power, absorbed below into the notation, is needed to descend through finite stabilizer component groups.

Main theorem (Li–Wang–Xu, Theorems 1.1 and 1.2). *The CM line bundle descends to a line bundle Λ_{CM} on $\overline{\mathcal{M}}$. It has a canonical continuous Hermitian metric h_{DP} whose curvature is a positive $(1,1)$ -current ω_{WP} extending the classical Weil–Petersson form on $\mathcal{M}$.*

Moreover Λ_{CM} is nef and big. If $n : \overline{\mathcal{M}}^n \rightarrow \overline{\mathcal{M}}$ is the normalization (an isomorphism over $\mathcal{M}$), then for $r \gg 1$ the rational map defined by $|n^(r\Lambda_{\text{CM}})|$ embeds $\mathcal{M}$. Hence $\mathcal{M}$ is quasi-projective.*

The point is not just positivity on the smooth moduli locus. The curvature must remain a well-defined positive current with controlled potentials at singular K-polystable limits, and the metrized bundle must descend to the coarse good moduli space.

Ingredient I: partial C^0 control of singular limits

For a local flat family $\pi : \mathcal{X} \rightarrow S$, choose a fixed pluri-anticanonical embedding and let $\tilde{\Omega}_t$ be the induced Fubini–Study volume form on the fibre X_t . Writing the Kähler–Einstein volume as

$$\omega_t^n = e^{-\tilde{u}_t} \tilde{\Omega}_t, \quad (1)$$

Moser iteration bounds the Bergman kernel from above, while the partial C^0 estimate bounds it from below; uniqueness of the Gromov–Hausdorff limit controls the embedding data. The authors obtain a continuous, uniformly bounded family of potentials $\tilde{u}_t$. The nonformal input is their Lemma 7.1:

$$\lim_{t \rightarrow 0} \int_{X_t} \tilde{\Omega}_t = \int_{X_0} \tilde{\Omega}_0. \quad (2)$$

This supplies uniform integrability near the singular set, where ordinary smooth convergence says nothing.

Ingredient II: Deligne pairing produces the metric

In additive notation the CM bundle is the dual Deligne pairing

$$\lambda_{\text{CM}} \simeq \langle -K_{\mathcal{X}/S}, \dots, -K_{\mathcal{X}/S} \rangle^{-1}. \quad (3)$$

The smooth reference metric from $\tilde{\Omega}$ induces a continuous Deligne metric $\tilde{h}_{\text{DP}}$. The change-of-metric formula corrects it by fibre integrals of $\tilde{u}$ against mixed products of the Fubini–Study and Kähler–Einstein forms:

$$h_{\text{DP}} = \tilde{h}_{\text{DP}} e^{-U}. \quad (4)$$

Continuity of $\tilde{u}$, convergence of mixed Monge–Ampère measures, and (2) make U continuous and uniformly bounded. On the smooth locus, the Deligne curvature formula and the change-of-metric identity give

$$-\frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} \log h_{\text{DP}} = \omega_{\text{WP}}^\circ. \quad (5)$$

Continuous local weights that are plurisubharmonic off the boundary extend uniquely as plurisubharmonic functions. Thus (5) extends as a positive current without trying to form an ill-defined singular fibre integral of an $(n+1)$ st power.

Ingredient III: descent through the good quotient

Near a closed K-polystable point, the compactification has an étale local GIT presentation by a slice modulo $\text{Aut}(X)$. The identity component acts trivially on the CM fibre because its character is the Futaki invariant, which vanishes for a K-polystable Fano variety. The remaining finite group has order $|\text{Aut}(X)/\text{Aut}(X)^0|$; boundedness of the universal family bounds these orders uniformly. One tensor power therefore kills all stabilizer actions, so Kempf–Alper descent gives Λ_{CM} . Functoriality of metrized Deligne pairings makes h_{DP} invariant, hence it descends too. Its continuity on the quotient again reduces to (2), and positivity is checked after pulling analytic curves to local slices.

Ingredient IV: turn curvature into quasi-projectivity

The paper proves a criterion for a normal proper algebraic space Y : if a line bundle L has nonnegative top intersection on every irreducible subspace, and strictly positive top intersection on each one meeting an open Y° , then a high power defines a rational map embedding Y° .

For $L = n^* \Lambda_{\text{CM}}$, positivity of the current with continuous bounded potentials gives nonnegative intersections. If an irreducible subspace meets $\mathcal{M}$, a local slice can be chosen so that the universal family is generically effective. Along effective Kodaira–Spencer directions the Weil–Petersson form is genuinely Kähler, and therefore

$$L^{\dim Z} \cdot Z = \int_Z \omega_{\text{WP}}^{\dim Z} > 0. \quad (6)$$

The nonnegative and positive intersections give nefness and bigness; applying the criterion to (6) gives the asserted embedding of $\mathcal{M}$.

Toy model, hard point, and limitations

For a smooth, effective one-parameter family entirely inside $\mathcal{M}$, the construction reduces to the familiar picture: the Deligne/Quillen metric is smooth and its curvature is the positive Weil–Petersson form. Boundary singularities introduce exactly the two missing tasks—integrability and descent.

The hardest analytic step is (2). After weak semistable (toroidal) reduction, inversion of adjunction gives discrepancy exponents $b_i, c_j < 1$. Hence the local densities $|y_i|^{-2b_i} |y_j|^{-2c_j}$ are integrable; dominated convergence preserves the mass on the strict transform of X_0 , while a logarithmic-coordinate estimate shows that every other vertical stratum contributes mass tending to zero.

The argument uses the prior construction of $\overline{\mathcal{M}}$, the Yau–Tian–Donaldson theory, and partial C^0 estimates. It proves quasi-projectivity of the smooth locus, not projectivity of the full compactification or ampleness along every boundary stratum. The paper obtains projectivity for the smoothable K-polystable del Pezzo moduli spaces as a separate orbifold-boundary consequence.