

The volume of singular Kähler-Einstein Fano varieties

Yuchen Liu

arXiv:1605.01034

<http://arxiv.org/abs/1605.01034v5>

Local singularities bound global Fano volume

An n -dimensional $\mathbf{Q}$ -Fano variety is normal, has klt singularities, and has $-K_X$ ample and $\mathbf{Q}$ -Cartier. Let $p \in X$ be closed and let Z be a zero-dimensional closed subscheme supported at p , with ideal I_Z . Write $\text{lct}(X; I_Z)$ for its log canonical threshold and $\text{mult}_Z X$ for its Hilbert–Samuel multiplicity.

Main theorem (Liu, Theorems 1–3). *If X admits a Kähler–Einstein metric, then*

$$(-K_X)^n \leq \left(1 + \frac{1}{n}\right)^n \text{lct}(X; I_Z)^n \text{mult}_Z X. \quad (1)$$

Equivalently at the level of local minimization, every real valuation v centered at p gives the companion bound

$$(-K_X)^n \leq \left(1 + \frac{1}{n}\right)^n \widehat{\text{vol}}_X(v), \quad \widehat{\text{vol}}_X(v) = A_X(v)^n \text{vol}(v). \quad (2)$$

If analytically $(X, p) \simeq (\mathbf{C}^n/G, 0)$ with $G \subset GL(n, \mathbf{C})$ acting freely in codimension one, then

$$(-K_X)^n \leq \frac{(n+1)^n}{|G|}. \quad (3)$$

Equality in (3) holds exactly when $|G \cap \mathbf{G}_m| = 1$ and $X \simeq \mathbf{P}^n/G$.

Thus stability turns the severity of one local singularity into a sharp global restriction. The proofs of (1) and (2) actually require only Ding-semistability; existence of the metric supplies it by Berman’s theorem.

Ingredient I: a blow-up volume profile

Let $\sigma: \widehat{X} \rightarrow X$ be the blow-up along Z and $\mathcal{O}_{\widehat{X}}(-F) = \sigma^{-1}I_Z \cdot \mathcal{O}_{\widehat{X}}$. For any ample $\mathbf{Q}$ -divisor L , Lemma 10 proves

$$\text{vol}_{\widehat{X}}(\sigma^*L - xF) \geq L^n - \text{mult}_Z X x^n \quad (4)$$

for $x \geq 0$. The polynomial on the right is the top self-intersection; the nontrivial direction is that the volume cannot fall below it. The proof uses sections on the thickened exceptional divisor, asymptotic Riemann–Roch, and vanishing. Lemma 10 also characterizes the Seshadri constant as the initial interval on which volume equals self-intersection.

Ingredient II: Ding stability integrates (4)

Fujita's theorem says that Ding-semistability makes the beta invariant of every proper subscheme nonnegative. For the present Z , with $V = (-K_X)^n$, this is exactly

$$\mathrm{lct}(X; I_Z)V \geq \int_0^\infty \mathrm{vol}_{\widehat{X}}(\sigma^*(-K_X) - xF) dx. \quad (5)$$

Use (4) until its right side vanishes, at $a = (V/\mathrm{mult}_Z X)^{1/n}$. Then

$$\int_0^\infty \mathrm{vol}(\sigma^*(-K_X) - xF) dx \geq \int_0^a (V - \mathrm{mult}_Z X x^n) dx = \frac{n}{n+1} Va. \quad (6)$$

Combining (5) and (6), cancelling V , and raising to the n th power gives (1). This short calculation is the decisive local-to-global bridge; all geometry is concentrated in the volume profile and the stability inequality.

Ingredient III: valuations and the same polynomial

For $L = -rK_X$ Cartier and a real valuation v , filter the section ring by

$$\mathcal{F}_v^x H^0(X, mL) = H^0(X, mL \cdot \mathfrak{a}_x(v)). \quad (7)$$

Lemma 19 shows that this filtration is saturated and, when $A_X(v) < \infty$, linearly bounded. Fujita's filtration theorem and a valuation on $X \times \mathbf{A}^1$ give Proposition 20, which bounds $A_X(v)$ below by an integral of the filtered volume. The exact sequence defined by (7) gives the analogue of (4),

$$\mathrm{vol}(\mathcal{F}_v S^{(x)}) \geq L^n - \mathrm{vol}(v)x^n. \quad (8)$$

Integrating (8) to its positive root produces (2). Theorem 27 then proves at every klt germ

$$\inf_I \mathrm{lct}(I)^n \mathrm{mult}(I) = \inf_v \widehat{\mathrm{vol}}(v), \quad (9)$$

with graded ideal sequences as the intermediary. This is why the ideal and valuation forms encode the same local obstruction.

Toy model, sharp application, and limitations

At a smooth point take $I_Z = \mathfrak{m}_p^q$. Then $\mathrm{lct}(I_Z) = n/q$ and $\mathrm{mult}(I_Z) = q^n$, so their product is n^n independently of the thickening. Formula (1) becomes Fujita's $(-K_X)^n \leq (n+1)^n$, with equality for $\mathbf{P}^n$. For a quotient germ, the pushed-forward order valuation has normalized volume $n^n/|G|$, and (2) gives (3). The hardest later step is equality: a finitely generated volume-minimizing valuation builds a zero-CM-weight test configuration, and K-polystability forces the projective-space quotient model.

The paper works over $\mathbf{C}$. It also uses (9) and normalized-volume minimization as follows. Let V be a Fano manifold of dimension $n-1$, and choose $r \in \mathbf{Q}_{>0}$ so that $H = -rK_V$ is an ample Cartier divisor. Then V is K-semistable exactly when every vertex-supported subscheme of the affine cone $C(V, H)$ satisfies $\mathrm{lct}^n \mathrm{mult} \geq r^{-1}(-K_V)^{n-1}$ (Theorem 7). The detailed equality analysis, existence questions for minimizers, and the classification consequences for log Del Pezzo surfaces are substantial further parts of the paper, omitted here.