

Variation of the canonical height for polynomials in several variables

Patrick Ingram

arXiv:1408.5416

<http://arxiv.org/abs/1408.5416v1>

Canonical height with bounded specialization error

Let K be a number field, X/K a projective curve, and $H \subset \mathbb{P}^N$ a hyperplane. A degree- $d \geq 2$ morphism $f: \mathbb{P}^N \rightarrow \mathbb{P}^N$ is *regular polynomial relative to H* when $f^*H = dH$ as divisors, not merely up to linear equivalence. A family f_t is *fibral* when the induced family on the invariant hyperplane $H \simeq \mathbb{P}^{N-1}$ is isotrivial (its moduli point is constant). Write η for the generic point of X and $\hat{h}_f$ for the canonical height.

Main theorem (Ingram, Theorem 1). *Let $f_t: \mathbb{P}^N \rightarrow \mathbb{P}^N$ be a fibral family of regular polynomial endomorphisms relative to H , defined over K , and let $P: X \rightarrow \mathbb{P}^N$ be a section. There is a divisor $A = A(f, P) \in \text{Div}(X) \otimes \mathbb{Q}$ of degree one such that, wherever the specialization is defined,*

$$\hat{h}_{f_t}(P_t) = \hat{h}_{f_\eta}(P_\eta) h_{X,A}(t) + O(1).$$

The constant is independent of t .

Call-Silverman gives only an $o(h(t))$ error for a general family. The bounded error is therefore the sharp natural form, since a Weil height is itself defined only up to $O(1)$. Fibrality is automatic for $N = 1$ but is a genuine restriction in higher dimension.

Ingredient I: uniform Green functions at infinity

Choose coordinates with $H = \{x_N = 0\}$ and $f_N = x_N^d$, and put

$$\| [x_0 : \cdots : x_N] \|_v = \frac{\max_{i < N} |x_i|_v}{|x_N|_v}, \quad G_{f,v}(Q) = \lim_{n \rightarrow \infty} d^{-n} \log^+ \|f^n(Q)\|_v.$$

The quantity $B_v(f)$ measures the lower-degree coefficients, hence how far one must go toward H before the top homogeneous part dominates. Lemma 3 proves that, once $\log \|Q\|_v > B_v(f) + \gamma_v$,

$$G_{f,v}(Q) = \log \|Q\|_v + O_v(1). \quad (1)$$

The proof compares f with its restriction to H , uses a homogeneous Nullstellensatz estimate, and telescopes the error under iteration. Both γ_v and the error in (1) depend only on that restriction. This is exactly where fibrality supplies uniformity as t varies.

Ingredient II: escape rates form the base divisor

After a finite extension over which a closed point $\beta \in X(\overline{K})$ is rational, regard β as a place of the corresponding function field and define

$$D_0(f, P) = \sum_{\beta \in X} G_{f_\eta, \beta}(P_\eta)[\beta]. \quad (2)$$

After replacing P by an iterate, every nonzero coefficient in (2) is in the range of (1), which also shows it is rational.

Theorem 4 is the local heart of the paper: on an affine open where f and P are regular, for every number-field place v ,

$$G_{f_t,v}(P_t) = \lambda_{X,D_0,v}(t) + O_v(1), \quad (3)$$

with an M_K -bounded error—it vanishes for all but finitely many v .

Ingredient III: patch every parameter regime

Fix w_β with its only pole at β . Lemmas 5–7 compare the pole order of any rational function with $\log^+ |w_\beta(t)|_v$. When the coefficient of $[\beta]$ in (2) is positive, (1) consequently identifies both sides of (3) with

$$\frac{G_{f_\eta,\beta}(P_\eta)}{\deg(w_\beta)} \log^+ |w_\beta(t)|_v + O_v(1). \quad (4)$$

The zero-coefficient case is subtler: a coefficient of f or coordinate of P may still have a pole at β . Lemma 9 expands all iterates in a uniformizer and uses $G_{f_\eta,\beta}(P_\eta) = 0$ to prove boundedness near β ; integrality makes the bound zero for almost every v . Lemma 8 controls the complement of all pole neighbourhoods. These three regions patch to the global local estimate (3).

From local comparison to the theorem

If $P(X) \subset H$, fibrality makes the restricted family constant after a coordinate change. Lemma 10 then reduces the formula to functoriality of ordinary Weil heights. Otherwise P_t meets H only finitely often. The divisor $D_0(f, P)$ is effective and

$$\deg D_0(f, P) = \widehat{h}_{f_\eta}(P_\eta). \quad (5)$$

Summing (3) over all places gives $\widehat{h}_{f_t}(P_t) = h_{X,D_0}(t) + O(1)$; normalizing (5) gives the degree-one divisor in the theorem. When the generic height is zero, the same local argument says that the specialized heights are bounded.

For an arbitrary divisor $B \in \text{Div}(X) \otimes \mathbb{Q}$, Corollary 2 uses comparison of equal-degree heights to obtain

$$\deg(B) \widehat{h}_{f_t}(P_t) = \widehat{h}_{f_\eta}(P_\eta) h_{X,B}(t) + O\left(h_{X,B}(t)^{1/2}\right),$$

and the error improves to $O(1)$ when X is rational.

Toy model, scope, and bottleneck

For $X = \mathbb{P}^1$, $f_t(z) = z^d + t$, and $P_t = 0$, the n th iterate has degree d^{n-1} in t , so $\widehat{h}_{f_\eta}(P_\eta) = 1/d$. The theorem reads $\widehat{h}_{f_t}(0) = d^{-1}h(t) + O(1)$ for a suitable representative of the standard height. Formula (2) is the higher-dimensional replacement for measuring that degree growth at the poles of the parameter.

The theorem does not cover arbitrary nonfibral higher-dimensional families; the paper observes that removing fibrality here would imply a bounded-error theorem for arbitrary families of projective endomorphisms. The technical bottleneck is not defining an escape rate, but proving that all its parameter dependence is captured by one $\mathbb{Q}$ -divisor while retaining an M_K -bounded error at every place, especially near poles with zero generic escape rate.