

Birational geometry of the moduli space of rank 2 parabolic vector bundles on a rational curve

Han-Bom Moon, Sang-Bum Yoo

arXiv:1409.6263

<http://arxiv.org/abs/1409.6263v2>

Mori's program stays inside the moduli problem

Work over an algebraically closed field of characteristic zero (the paper writes it as $\mathbb{C}$). Fix distinct points $p_1, \dots, p_n \in \mathbb{P}^1$, with $n \geq 5$. A rank-two parabolic bundle is $(E, \{V_i\}, \mathbf{a})$, where $V_i \subset E|_{p_i}$ is a line and $\mathbf{a} = (a_1, \dots, a_n) \in ((0, 1) \cap \mathbb{Q})^n$. Let $M(\mathbf{a})$ be the coarse moduli space of S-equivalence classes of degree-zero semistable objects. A weight is *effective* when its stable locus is nonempty, and *general* when it is effective and stability equals semistability.

For a maximal-Picard weight, write

$$M^+ = \mathrm{Bl}_{[\mathbf{p}]}((\mathbb{P}^1)^n //_{L\mathrm{SL}_2}), \quad L = \mathcal{O}(a_1, \dots, a_n), \quad (1)$$

and let E denote the exceptional divisor. The paper identifies the effective cone of $M(\mathbf{a})$ with that of M^+ .

Main theorem (Moon–Yoo, Theorems 6.2 and 7.3). *Assume first that $\mathbf{a}$ is general and $\rho(M(\mathbf{a})) = n + 1$. For every even subset $I \subset [n]$, $|I| = 2i$, set $e_j = \mathrm{pr}_j^* \mathcal{O}_{\mathbb{P}^1}(1)$, the j th coordinate class under the Picard-group identification, and*

$$D_I = \mathcal{O}\left(\sum_{j \in I} e_j\right) - (i - 1)E. \quad (2)$$

Then $\mathrm{Eff}(M(\mathbf{a}))$ is the polyhedral cone whose extremal rays are precisely the D_I ; hence it has 2^{n-1} rays. Without the Picard-rank hypothesis, for every general weight $\mathbf{a}$ and every $\mathbb{Q}$ -divisor $D \in \mathrm{int} \mathrm{Eff}(M(\mathbf{a}))$, the projective model

$$M(\mathbf{a})(D) = \mathrm{Proj} \bigoplus_{m \geq 0} H^0(M(\mathbf{a}), \mathcal{O}(\lfloor mD \rfloor))$$

is isomorphic to $M(\mathbf{b})$ for some parabolic weight $\mathbf{b}$.

Thus a large, non-simplicial effective cone is computed explicitly, and every interior chamber of its Mori decomposition retains a direct moduli interpretation.

Ingredient I: GIT produces the first blow-up

For an effective weight (equivalently here, an effective linearization) with $\sum a_i < 2$, semistability forces $E \simeq \mathcal{O}^{\oplus 2}$, and Proposition 3.5 gives

$$M(\mathbf{a}) \simeq (\mathbb{P}^1)^n //_{L\mathrm{SL}_2}. \quad (3)$$

The Hilbert–Mumford condition says that the total weight of points coinciding at any $y \in \mathbb{P}^1$ is at most half the total weight. The first fundamental theorem for SL_2 then expresses every invariant divisor through collision divisors $D_{\{i,j\}} = \{x_i = x_j\}$. With maximal stable locus the resulting cone is the cone over the hypersimplex $\Delta(2, n)$.

After a harmless generic perturbation, scale the weights without changing their ratios so that every crossing is simple. The first wall is $\sum a_i = 2$; Proposition 3.7 identifies its crossing with the blow-up in (1) at the configuration $[\mathbf{p}]$. Later crossings are flips or blow-downs. Consequently the maximal-Picard case has basis $e_1, \dots, e_n, E$ and reduces the general effective-cone calculation to a single-point blow-up of the GIT quotient.

Ingredient II: conformal blocks measure vanishing

Write a divisor as $\mathcal{O}(k_1, \dots, k_n) - tE$ and put $2N = \sum k_i$. For $t \geq 0$, Lemma 6.3 identifies its space of sections with the $\mathfrak{sl}_2$ conformal block

$$H^0(M^+, \mathcal{O}(k_1, \dots, k_n) - tE) \simeq \mathbb{V}_{N-t}(k_1, \dots, k_n), \quad (4)$$

because the exceptional multiplicity is exactly the order of vanishing of an invariant section at $\mathbf{p}$.

Nonzero conformal blocks are counted by boxed Catalan paths (double sequences). Before using Lemma 6.5, the proof permutes the labels so $k_1 \geq \dots \geq k_n$ and removes zero labels by propagation of vacua. The lemma then peels from a positive-shape path of height > 1 a nonempty even subset T : it lowers k_j by one for $j \in T$ and lowers the level and height by one. Multiplication of conformal blocks therefore splits an effective divisor into the level-one class D_T from (2) and a smaller effective class. Induction shows that $E = D_\emptyset$ and the level-one D_I generate the whole cone. Proposition 6.6 constructs n independent supporting functionals through each ray, proving extremality rather than mere generation.

Ingredient III: theta divisors identify every model

Lemma 7.1 makes $M(\mathbf{a})$ a Mori dream space, using a log-Fano weight and preservation under the wall-crossing flips. If the Picard rank is below $n + 1$, the proof first realizes $M(\mathbf{a})$ as a rational contraction of a maximal-Picard $M(\mathbf{a}')$ and embeds its effective cone into that of $M(\mathbf{a}')$; this reduces the model theorem to the maximal-Picard case. There write $D = \mathcal{O}(b_1, \dots, b_n) - tE$ and set $N = \frac{1}{2} \sum b_i$.

If $t > 0$, interiority implies $mD - E$ is effective for $m \gg 0$. The conformal-block vanishing rule then gives $b_i < N - t$ for every i . Pauly's generalized theta divisor for the weight $\mathbf{b}/(N - t)$ is ample and has the section space in (4), so its section ring yields

$$M(\mathbf{a})(D) \simeq M\left(\frac{\mathbf{b}}{N - t}\right). \quad (5)$$

If $t \leq 0$, the exceptional component does not change the model; the section ring contracts to the GIT quotient for $\mathcal{O}(\mathbf{b})$, which (3) identifies with $M(c\mathbf{b})$ for a sufficiently small $c > 0$. These are the two cases of Theorem 7.3.

The first nontrivial picture

For $n = 5$, the Picard rank is 6 but the theorem gives $2^4 = 16$ extremal rays:

$$E, \quad \mathcal{O}(e_i + e_j) \ (1 \leq i < j \leq 5), \quad \mathcal{O}\left(\sum_{j \in I} e_j\right) - E \ (|I| = 4).$$

The ten middle rays are the pair-collision invariants of the GIT quotient; the five four-subset rays record first-order vanishing at the blown-up configuration. The Catalan-path reduction says that every effective divisor class arising at higher level is numerically generated by these level-one classes (and E).

The hypotheses matter. Both results concern rank two and degree zero on $\mathbb{P}^1$ with general weights; the explicit ray formula also requires $n \geq 5$ and maximal Picard rank, while the model theorem holds for every general weight. Lower-Picard cones are obtained by pushing through flips and contractions. Boundary divisors are discussed separately in Remark 7.4 and may drop parabolic points or change degree. Extending the closure-under-Mori-program statement to arbitrary rank or a positive-genus base curve remains the paper's Question 1.3. The hard step is the combinatorial peeling lemma: it converts nonvanishing of a general conformal block into a terminating factorization by the exact classes that become the extremal rays.