Rational self-maps with a regular iterate on a semiabelian variety

Jason Bell, Dragos Ghioca, Zinovy Reichstein

arXiv:2208.06938

http://arxiv.org/abs/2208.06938v1

When does a regular iterate regularize the map?

A semiabelian variety is an algebraic group fitting into an exact sequence

$$1 \longrightarrow \mathbf{G}_m^N \longrightarrow G \longrightarrow A \longrightarrow 1$$

with A an abelian variety. Let $\Phi\colon G \dashrightarrow G$ be a dominant rational self-map. Although Φ^m can be a morphism while Φ is not, the paper shows that on G this failure must be visible on a nontrivial quotient dynamical system.

Main theorem (Bell–Ghioca–Reichstein, Theorem 1.1)**.** *Let K be algebraically closed and suppose that $\Phi^m\colon G \to G$ is regular for some $m \geq 1$. Assume that there is no nonconstant homomorphism of semiabelian varieties $\chi\colon G \to G_0$ such that*

$$\chi \circ \Phi^{mk} = \chi \qquad \text{for some } k \geq 1.$$

If $\operatorname{char} K = 0$*, then Φ is regular.*

If $\operatorname{char} K = p > 0$*, the same conclusion holds provided one also excludes a semiabelian variety G_1 over a finite subfield, a nonconstant group homomorphism of semiabelian varieties $\chi\colon G \to G_1$, and $k, r \geq 1$ for which*

$$\chi \circ \Phi^{mk} = F^r \circ \chi,$$

where F is the corresponding Frobenius endomorphism.

Thus regularity descends from an iterate unless a quotient is periodic; in positive characteristic, a Frobenius quotient is the additional genuine obstruction. The characteristic-zero statement is the dynamical core of the result.

Ingredient 1: regularity can be pulled through an isogeny

Every dominant regular self-map of G has the form $T_\alpha \circ \Psi_0$, with Ψ_0 a group endomorphism. It factors as a purely inseparable homomorphism followed by an étale morphism. Lemma 4.1 uses this factorization to prove the following pointwise cancellation statement: if $g = f \circ \Psi$ with $\Psi\colon G \to G$ dominant and regular, then g is regular at x exactly when f is regular at $\Psi(x)$.

The nontrivial implication is local. In completed local rings, let A be the image of the ring at $\Psi(x)$. The purely inseparable–étale factorization and smoothness of G give

$$\operatorname{Frac}(A) \cap \widehat{\mathcal{O}}_{G,x} = A.$$

Hence a rational germ of f whose pullback is regular was already regular. This is precisely the cancellation that fails for general smooth affine varieties.

Ingredient 2: persistent indeterminacy is a hypersurface

Assume for contradiction that Φ is not regular and put $\Psi = \Phi^m$. The indeterminacy locus W_n of Φ^n is pure of codimension one, and

$$W_{n+m} \subseteq W_n.$$

The descending chain $W_1 \supseteq W_{m+1} \supseteq W_{2m+1} \supseteq \cdots$ stabilizes. Applying the cancellation lemma to two consecutive stabilized iterates yields

$$W = \Psi^{-1}(W).$$

The finitely many irreducible components of W are permuted by Ψ. After replacing Ψ by an iterate, purity of inverse images lets one choose an irreducible codimension-one component that is itself totally invariant. This is Proposition 1.2: nonregularity has produced a geometric object rigid enough to classify.

Ingredient 3: an invariant hypersurface forces a quotient

Let W be the totally invariant hypersurface. Quotienting G by the translation stabilizer of W preserves total invariance and reduces to the case $\mathrm{Stab}(W) = 0$. If $\dim G = 1$, then W is a point. After translating it to the identity, Ψ is a group endomorphism with infinitesimal kernel. For $\mathbf{G}_m$ this means an exponent ± 1 in characteristic zero or $\pm p^r$ in characteristic p; for an elliptic curve, the separable part is an automorphism, while an inseparable part is a Frobenius power after passage to a finite-field form. An iterate therefore gives precisely the periodic or Frobenius quotient in Proposition 1.3.

Assume now $\dim G > 1$. The Pink–Rössler structure theorem writes

$$W = \gamma + h_1(X_1) + \cdots + h_\ell(X_\ell),$$

where $\gamma \in G$, each X_i is a closed invariant subvariety of a semiabelian variety A_i, and $h_i \colon A_i \to G$ is a homomorphism with $\Psi_0 h_i = h_i f_i$ for a pure-weight endomorphism f_i of A_i. Trivial stabilizer and codimension one make the product map from the A_i an isogeny onto G.

In characteristic zero every weight is zero. After an iterate, all f_i are identities, so Ψ_0 is the identity on the factor images and hence on G; invariance of W and its trivial stabilizer then kill the translation part. Undoing the stabilizer quotient supplies a nonconstant χ with $\chi\Psi^k = \chi$. In characteristic p, a positive-weight factor is defined over a finite field and carries a Frobenius power. On that factor $f_i - \mathrm{id}$ is dominant, so a translation conjugacy kills the corresponding component of the translation part of Ψ. Projection of an equivariant isogeny then satisfies $\chi\Psi^k = F^r\chi$, not merely the analogous relation for Ψ_0. These are exactly the alternatives excluded in the theorem.

Sharp model, hard point, and scope

On $\mathbf{G}_m$, the rational map

$$\Phi(x) = -\frac{x}{x+1}$$

is not regular but satisfies $\Phi^2 = \mathrm{id}$. The identity quotient is fixed by the regular iterate, so this example shows that the quotient hypothesis cannot simply be removed.

The hardest bridge is from a set-theoretically invariant hypersurface to a semiabelian quotient: it requires both the Pink–Rössler classification and the codimension-one/trivial-stabilizer argument making the factor map an isogeny. The result is special to semiabelian rigidity; it is false for general smooth affine varieties. It is vacuous when G is abelian because rational maps from a semiabelian variety to an abelian variety are already regular, while positive characteristic necessarily retains the Frobenius alternative.