

# Monodromy of rational curves on K3 surfaces of low genus

Sailun Zhan

arXiv:2004.08719

<http://arxiv.org/abs/2004.08719v3>

## The enumerative set has no hidden labels

Let  $(S, L)$  be a generic complex K3 surface of genus  $g$ , so  $L^2 = 2g - 2$ , and consider the rational curves in  $|L|$ . Their numbers for  $g = 1, 2, 3$  are 24, 324, 3200. As  $S$  travels around loops in the open moduli where these curves remain distinct, analytic continuation permutes them. The resulting image in the relevant symmetric group is the monodromy group.

**Main theorem** (Zhan, Theorems 1.1–1.3). *For the genus-1 elliptic K3 surfaces with section considered in the paper, the monodromy on the 24 nodal rational curves is  $S_{24}$ . For generic primitive K3 surfaces of genus 2 and 3, the monodromy groups on the 324 and 3200 rational curves are respectively*

$$S_{324} \quad \text{and} \quad S_{3200}. \quad (1)$$

Thus the geometric family imposes no invariant partition, parity constraint, or other label on these curves. Over a number field, Hilbert irreducibility then yields dense specializations in which an individual curve is defined over a non-Galois extension of degree 24, 324, or 3200.

## Genus one: two overlapping permutation patterns

An elliptic K3 surface with section has a Weierstrass model over  $\mathbf{P}^1$

$$y^2z = x^3 + Axz^2 + Bz^3, \quad A \in H^0(\mathcal{O}(8)), \quad B \in H^0(\mathcal{O}(12)). \quad (2)$$

The 24 rational curves are the nodal fibers over the zeros of  $\Delta = 4A^3 + 27B^2$ . Hence the problem is the monodromy of the degree-24 cover

$$Z = \{(A, B, t) : \Delta(t) = 0\} \longrightarrow V = H^0(\mathcal{O}(8)) \times H^0(\mathcal{O}(12)). \quad (3)$$

Two explicit factorizations of  $\Delta$  produce subgroups conjugate to the diagonal  $S_{12}$  acting on two blocks of size 12 and the diagonal  $S_8$  acting on three blocks of size 8. Their incompatible block structures make the combined action primitive (Lemma 2.3). The classification of primitive degree-24 groups, together with  $|M| > 12!$ , leaves  $A_{24}$  and  $S_{24}$ ; the  $S_8$  subgroup contains an odd permutation, so  $M = S_{24}$ .

## Genus two: bitangents of a sextic

A generic primitive genus-2 K3 surface is the double cover of  $\mathbf{P}^2$  branched over a smooth plane sextic  $C$ . A member of  $|L|$  is rational exactly when its image is one of the 324 bitangent lines of  $C$ : the two tangencies remove the ramification in the induced double cover of the line. Harris's theorem says that the bitangents of a generic plane curve of degree at least 5 have full symmetric monodromy. Applied at degree 6, this immediately gives  $S_{324}$ .

## Genus three: double transitivity

A generic primitive genus-3 K3 surface is a smooth quartic  $S \subset \mathbf{P}^3$ . Its rational hyperplane sections are plane quartics with three nodes, cut out by planes tangent to  $S$  at three noncollinear points. Let  $W$  parametrize quartic surfaces, let  $J_{\text{ord}} \rightarrow W$  record ordered tangent triples, and let  $J_{\text{un}} = J_{\text{ord}}/S_3$  record their tangent planes.

Before quotienting by the triple permutations, the fibers of the incidence space over the configuration of points are linear subspaces of codimension 9. Their irreducibility makes  $J_{\text{ord}}$  irreducible, hence the monodromy transitive. Repeating the incidence argument while fixing one tangent plane shows that its stabilizer is transitive on the remaining planes (Lemma 4.1). Thus  $M \subset S_{3200}$  is doubly transitive.

## The decisive degeneration supplies one transposition

A standard local-monodromy lemma says that a degree- $n$  cover has a simple transposition if some fiber has  $n - 1$  distinct points—one double point and  $n - 2$  simple points—and the covering space is locally irreducible at the double point. Zhan constructs exactly this collision for quartic K3 surfaces.

Choose an irreducible plane quartic  $g(x, y, z) = 0$  with two nodes and one cusp, and vary

$$S_f : \quad g(x, y, z) + wf(x, y, z, w) = 0, \quad (4)$$

where  $f$  is cubic. Dimension counts exclude reducible hyperplane sections for general  $f$ . Incidence spaces stratified by one, two, or three tangent points, together with semicontinuity of Hessian corank and Milnor number, show that every other rational tangent section has three nodes (Lemmas 4.4–4.5). Thus a smooth member of (4) has one rational tangent curve with two nodes and one cusp and otherwise only trinodal rational tangent curves (Proposition 4.6).

The Euler characteristic of the compactified Jacobian is 1 for a trinodal rational curve and 2 for a rational curve with two nodes and one cusp. Since the total Yau–Zaslow count is 3200, the special surface therefore has

$$3198 \cdot 1 + 1 \cdot 2 = 3200. \quad (5)$$

The  $J_{\text{un}}$ -fiber has 3199 distinct points: the cuspidal point counts with multiplicity two. The ordered incidence  $J_{\text{ord}}$  is smooth along it, and  $J_{\text{un}}$  is locally irreducible, so the local-monodromy lemma yields a transposition. Double transitivity conjugates this transposition to one swapping any prescribed pair; these transpositions generate  $S_{3200}$ .

## A toy branch point

The family of roots of  $u^2 - t$  over  $t \neq 0$  has two sheets. A loop around  $t = 0$  exchanges the roots, because the special fiber has one double root. This is the local picture behind the cuspidal quartic in (4): two trinodal curves coalesce and a small loop exchanges them. What the toy model omits is the global burden of proving that the degenerating K3 remains smooth, that no other rational sections collide, and that the incidence cover is locally irreducible.

## Proof roadmap and hardest point

The three genera use different geometry:

$$\begin{aligned} g = 1 & : \text{ explicit discriminants } \Rightarrow \text{ primitive large subgroup } \Rightarrow S_{24}, \\ g = 2 & : \text{ double plane } \Rightarrow \text{ sextic bitangents } \Rightarrow S_{324}, \\ g = 3 & : \text{ irreducible incidences } \Rightarrow \text{ double transitivity + one collision } \Rightarrow S_{3200}. \end{aligned} \quad (6)$$

The hard point is not the final group theory but the controlled collision in genus 3. The deformation (4) must land on the branch divisor simply, while avoiding every higher singularity and reducible section.

## A second check from the Gauss map

Section 5 verifies the special count without compactified Jacobians. For the Gauss map of a “good” quartic, let  $C_d$  be its double-cover curve and  $C_{\text{par}}$  its parabolic curve. Piene’s formula reads

$$b(\mu_0 - 2) = \rho + \delta_b + v_b'' + i, \quad (7)$$

where  $b = 480$  is the degree of  $C_d^*$ ,  $\mu_0 = 36$  the degree of the dual surface,  $\rho = 320$  the degree of  $C_d$ , and  $i = C_d \cdot C_{\text{par}} = 2560$ . A local analysis of the node–cusp collision shows that there are 1918 other node-and-cusp tangent curves and that the normalization ramification term is  $v_b'' = 1918 + 2 = 1920$ . Equation (7) then gives  $\delta_b = 11520$ .

If  $t$  is the number of trinodal tangent curves, their three nodes contribute  $3t$  to  $\delta_b$ ; the 1918 node–cusp curves contribute 1918, while the exceptional curve contributes 2 at its cusp and 3 at each of its two nodes. Thus

$$3t + 1918 + 2 + 2 \cdot 3 = 11520, \quad t = 3198, \quad (8)$$

recovering (5). This calculation exposes the local singularity bookkeeping hidden by the much shorter Euler-characteristic proof.

The theorem is generic and restricted to  $g \leq 3$  (with the stated elliptic-with-section convention for  $g = 1$ ). Full symmetric monodromy for higher-genus primitive K3 surfaces remains open; the argument does not claim that every individual K3 surface has 3200 distinct rational sections.