Some families of polynomials satisfying uniform boundedness for rational preperiodic points

Hector Pasten

Hector Pasten, arXiv:2408.13262v1 (2024), 2 pages
http://arxiv.org/abs/2408.13262v1

Uniform boundedness in a non-isotrivial family

The *UB property* for $\mathcal{F} \subseteq \mathbb{Q}(z)$ means that some $B = B(\mathcal{F})$ bounds the number of $\mathbb{Q}$-rational preperiodic points of every $f \in \mathcal{F}$ on $\mathbf{P}^1$. The Morton–Silverman conjecture predicts this for all degree-d rational maps; even the full family $z^2 + c$, $c \in \mathbb{Q}$, remains open.

Let $\mathrm{pole}(\psi)$ denote the pole divisor of $\psi \in \mathbb{Q}(t)$ on $\mathbf{P}^1$. Its support may be irreducible over $\mathbb{Q}$ while splitting into several conjugate geometric points.

Main theorem (Pasten, Theorem 1.1)**.** *Fix an integer $d \geq 2$. Suppose that the support of* $\mathrm{pole}(\psi)$ *is irreducible on $\mathbf{P}^1_{\mathbb{Q}}$ and contains at least two geometric points. Then*

$$\mathcal{F}_{d,\psi} = \{z^d + \psi(c) : c \in \mathbb{Q}\}$$

has the UB property.

The bound is uniform in c but may depend on (d, ψ). Since the pole hypothesis forces ψ to be nonconstant, these are explicit non-isotrivial families.

Ingredient 1: Chebotarev removes every parameter denominator

The pole hypothesis gives a homogeneous presentation

$$\psi(y/x) = \frac{G(x,y)}{A\,H(x,y)^n}, \qquad G, H \in \mathbb{Z}[x,y], \quad \deg G = n \deg H,$$

where $A \neq 0$, $n > 0$, and H is irreducible of degree at least two. Chebotarev supplies infinitely many primes ℓ for which $H \bmod \ell$ has no linear factor over $\mathbb{F}_\ell$; discard the finitely many with $\ell \mid A$.

Write any parameter as $c = b/a$ with $(a, b) = 1$. The pair $(a : b)$ is a point of $\mathbf{P}^1(\mathbb{F}_\ell)$, whereas a zero of the homogeneous binary form $H \bmod \ell$ at such a point would give a linear factor. Hence $\ell \nmid H(a, b)$ for *every* $c \in \mathbb{Q}$, so ℓ does not divide the denominator of $\psi(c)$. Since $z^d + \alpha$ has good reduction precisely at primes not dividing the denominator of α, one prime chosen this way works simultaneously for the entire family (Lemma 2.1). This reversal of quantifiers—first ℓ, then all c—is the paper's decisive construction.

Ingredient 2: two good primes bound all rational periods

Choose two distinct primes p, q from Lemma 2.1 and put $N = (p^2 - 1)(q^2 - 1)$. The cited Corollary B of Morton–Silverman implies that every rational periodic point of every $z^d + \psi(c)$ has exact period at most N. Crucially, N is independent of c; all parameter dependence was eliminated by the common-good-reduction step.

Ingredient 3: bounded periods control the whole tail

Pasten then invokes the following theorem of Doyle–Poonen, reproduced as Theorem 2.2 in the paper: for fixed $d \geq 2$ and $N \geq 1$, there is a bound $B(d, N)$ such that any $z^d + \alpha$, $\alpha \in \mathbb{Q}$, whose rational periodic points all have exact period at most N has at most $B(d, N)$ rational preperiodic points. Applying it with the common N above proves Theorem 1.1. In other words, good reduction bounds the possible cycles, while the Doyle–Poonen theorem uniformly bounds the rational tails feeding those cycles.

A concrete subfamily and the scope of the argument

For $d = 2$ and $\psi(t) = 2/(t^2 + 8)$, the pole support is the irreducible degree-two locus $t^2 + 8 = 0$. Moreover, if $c^2 + 8 = s^2$ with $c, s \in \mathbb{Q}$, then $z^2 + 2/s^2$ has the rational fixed points $z = (1 \pm c/s)/2$. The conic $c^2 + 8 = s^2$ has the point $(1, 3)$ and infinitely many rational points. Thus infinitely many members can have an affine rational preperiodic point: the theorem is not merely a construction in which all interesting rational dynamics disappears.

The result concerns the restricted parameter image $\psi(\mathbb{Q})$, not all constants $c \in \mathbb{Q}$, and it is formulated over $\mathbb{Q}$. It does not provide a numerical value of B: the deepest uniform tail bound is the imported Doyle–Poonen theorem, while the paper's new two-page argument is the Chebotarev mechanism producing parameter-independent good reduction.

Codex mathematical paper digest, 2026-10-18. Source paper should be consulted for full statements and proofs.