An upper bound for the height for regular affine automorphisms of A^n

Chong Gyu Lee

arXiv:0909.3107

http://arxiv.org/abs/0909.3107v1

A two-sided height inequality

Let $f : \mathbf{A}^n \to \mathbf{A}^n$ be a polynomial automorphism, let $\phi_0, \psi_0 : \mathbf{P}^n \dashrightarrow \mathbf{P}^n$ be the homogeneous extensions of f and f^{-1}, and put $d = \deg f$ and $d' = \deg f^{-1}$. The map is *regular* when $Z(\phi_0) \cap Z(\psi_0) = \varnothing$. Thus the forward and backward indeterminacy at the hyperplane $H = \mathbf{P}^n \setminus \mathbf{A}^n$ are separated. Write h for the logarithmic projective height.

Main theorem (Lee, Theorem 3.1)**.** *For a regular affine automorphism f of $\mathbf{A}^n$, there is a constant $C = C(f)$ such that every algebraic point $P \in \mathbf{A}^n$ satisfies*

$$\frac{1}{d}h(f(P)) + \frac{1}{d'}h(f^{-1}(P)) \geq \left(1 + \frac{1}{dd'}\right) h(P) + C.$$

Silverman's inequality for the jointly regular pair $\{\phi_0, \psi_0\}$ has coefficient 1 on $h(P)$. The extra $1/(dd')$ is the decisive gain: it proves Kawaguchi's conjectured estimate in all dimensions and supplies the geometric input used in his construction of canonical heights for regular affine automorphisms.

Ingredient 1: one exceptional divisor carries infinity

Resolve ϕ_0 by a sequence of smooth blowups $\pi_V : V \to \mathbf{P}^n$, and let $\phi = \phi_0 \circ \pi_V$ be the resulting morphism. If H_V is the strict transform of H and $E_1, \ldots, E_k$ are the exceptional divisors, then

$$\mathrm{Pic}(V) = \mathbf{Z}H_V \oplus \bigoplus_i \mathbf{Z}E_i.$$

Regularity gives $\phi_0(H \setminus Z(\phi_0)) \subset Z(\psi_0)$. Consequently $\phi_* H_V = 0$, while $\phi_* \phi^* H = H$. Since every coefficient of $\phi^* H$ is positive, Theorem 2.9 forces a unique *essential* exceptional divisor E with coefficient one and $\phi_* E = H$; every other exceptional divisor is contracted in codimension at least two. This turns the complicated boundary of the resolution into one distinguished channel through which infinity survives.

Ingredient 2: nefness controls the remaining coefficients

For a birational morphism $\rho : \widetilde{X} \to X$, Lemma 3.4 says that an effective nef divisor D satisfies $\rho^* \rho_* D \geq D$. This is a negativity-lemma calculation: the exceptional coefficients in $D - \rho^* \rho_* D$ are nonpositive. Applied to the resolution, it yields

$$\pi_V^* H = H_V + d'E + M, \qquad \phi^* H = dH_V + E + I, \qquad d'I \geq M,$$

where M and I are effective and avoid H_V and E. Lemma 3.6 is what makes the last inequality coefficientwise: in the span of the nonessential exceptional divisors, a class is effective exactly when all its coefficients are nonnegative.

Ingredient 3: combine the forward and backward resolutions

Resolve ψ_0 similarly. Because the two indeterminacy loci are disjoint, both resolutions are dominated by one blowup $\pi_U : U \to \mathbf{P}^n$. On U the forward and backward pullbacks have matching decompositions; with E, F the two essential divisors and M, I, N, J the nonessential effective parts, the inequalities are $d'I \geq M$ and $dJ \geq N$. Direct substitution shows that

$$D := \frac{1}{d}\phi_U^* H + \frac{1}{d'}\psi_U^* H - \left(1 + \frac{1}{dd'}\right)\pi_U^* H = \left(1 - \frac{1}{dd'}\right) H_U + \frac{dJ - N + d'I - M}{dd'}$$

is effective. The Weil height machine converts this divisor inequality, up to a bounded function, into the theorem.

Proof roadmap and a concrete model

The argument therefore has three steps: resolve each rational extension; use push–pull to identify the sole essential divisor and nefness to bound all other coefficients; then pass to the common resolution and read the height inequality from the effective divisor D.

The paper's three-dimensional Hénon-type example

$$f(x, y, z) = (y, z + y^2, x + z^2)$$

has $(d, d') = (2, 4)$. After resolving the two disjoint loci at infinity, the displayed divisor becomes an explicit nonnegative combination of the strict transform of H and the exceptional divisors. This is a useful toy model for the general bookkeeping, but the general proof cannot rely on explicit coordinates or a prescribed sequence of blowups.

The hardest point is not resolution itself but the uniform sign control on every nonessential exceptional coefficient. The theorem also depends essentially on regularity; it does not assert the strengthened bound for an arbitrary polynomial automorphism whose forward and backward indeterminacy loci meet.

Codex mathematical paper digest, 2026-10-24. Source paper should be consulted for full statements and proofs.