

S-integral preperiodic points for monomial semigroups over number fields

Marley Young

arXiv:2402.13713

<http://arxiv.org/abs/2402.13713v1>

A semigroup form of the Ih finiteness problem

Let K be a number field, let S be a finite set of places containing the archimedean ones, and let

$$\mathcal{G} = \langle f_1, \dots, f_s \rangle, \quad f_i(z) = a_i z^{d_i}, \quad a_i \in K^\times, \quad d_i \in \mathbb{Z}, \quad |d_i| \geq 2. \quad (1)$$

A point $\alpha \in \overline{K}$ is *preperiodic for $\mathcal{G}$* if two prefixes along one word agree: $f_{i_1 \dots i_n}(\alpha) = f_{i_1 \dots i_m}(\alpha)$ for some $n > m \geq 0$, where $f_{i_1 \dots i_n} = f_{i_n} \circ \dots \circ f_{i_1}$. This is weaker than asking that the whole semigroup orbit be finite. Relative S -integrality means that no conjugate of α meets a conjugate of β at a place outside S .

Main theorem (Young, Theorem 1.5). *Fix $D \geq 1$. There is an effectively computable uniform bound C such that, for every $\beta \in \overline{K}$ with $[K(\beta) : K] \leq D$ which is not preperiodic for $\mathcal{G}$, at most C points $\alpha \in \overline{K}$ are both preperiodic for $\mathcal{G}$ and S -integral relative to β .*

This proves the monomial-semigroup case of the paper's analogue of Ih's conjecture. Uniformity in the bounded-degree moving point β is the subtle feature: both the dynamical sequence witnessing preperiodicity and the local distances to β vary.

Ingredient I: heights and measures uniform in the word

A witness $i_1, \dots, i_n$ for α determines a sequence

$$\mathbf{f} = (f_{i_1}, \dots, f_{i_m}, \overline{f_{i_{m+1}}, \dots, f_{i_n}})$$

obtained from $\mathcal{G}$, for which α has finite forward orbit. Kawaguchi's canonical height $\widehat{h}_{\mathbf{f}}$ therefore vanishes at α . Proposition 2.1 gives

$$\widehat{h}_{T\mathbf{f}}(f_{i_1}(x)) = |d_{i_1}| \widehat{h}_{\mathbf{f}}(x), \quad |\widehat{h}_{\mathbf{f}}(x) - h(x)| \leq 2c(\mathcal{G}). \quad (2)$$

If β is not semigroup-preperiodic, Lemma 2.2 and Northcott's theorem give a positive lower bound for $\widehat{h}_{\mathbf{f}}(\beta)$, uniform over all such sequences (after the bounded-degree base change).

Proposition 3.3 associates to every $\mathbf{f}$ an adelic equilibrium measure $\rho_{\mathbf{f}} = \{\rho_{\mathbf{f},v}\}_v$ whose height is $\widehat{h}_{\mathbf{f}}$. Its local potentials have a common Hölder exponent and constants depending only on the finite set of generators. Thus the Favre–Rivera–Letelier quantitative equidistribution estimate (Theorem 3.1 and Proposition 3.2) is uniform in the word, rather than only for one fixed map.

Ingredient II: binomial structure controls local proximity

The equality of two monomial prefixes makes every nonzero preperiodic α a root of a binomial $X^N - a$. Proposition 4.1 sharpens this: $\alpha = \zeta\gamma$, where ζ is a root of unity and either $\gamma = 1$ or

$$\gamma^M = a_1^{m_1} \cdots a_s^{m_s} b, \quad |m_i| < M, \quad (3)$$

with b in a fixed finite set of bounded height. Capelli's criterion and Schinzel's binomial factorization results give

$$\frac{M}{2} \leq [K(\gamma) : K] \leq M, \quad \log(MQ) \ll_{[K:\mathbb{Q}]} \log[K(\alpha) : K] \quad (4)$$

when ζ has order Q . This is the essential degree lower bound: the exponents occurring in a linear form cannot grow invisibly while $[K(\alpha) : K]$ stays small.

Applying archimedean and non-archimedean lower bounds for linear forms in logarithms to (3), Theorem 5.1 obtains, at every place v ,

$$\min_{\sigma: K(\alpha)/K \hookrightarrow \overline{K}_v} \log |\sigma(\alpha) - \beta|_v > -C_v (h(\beta) + 1) \log[K(\alpha) : K]. \quad (5)$$

It prevents a few exceptionally close conjugates from overwhelming an equidistribution average.

The product-formula contradiction

Write the preperiodic relation as $g_2(g_1(\alpha)) = g_1(\alpha)$ with $g_1(z) = az^k$ and $g_2(z) = bz^\ell$. A direct calculation gives the exact height identity

$$\widehat{h}_{\mathbf{f}}(\beta) = \sum_{v \in M_K} \log \max\{|\alpha|_v, |\beta|_v\}. \quad (6)$$

For the Galois orbit of α , consider

$$\Gamma = \frac{1}{[K(\alpha) : K]} \sum_{v \in M_K} \sum_{\sigma: K(\alpha)/K \hookrightarrow \overline{K}_v} \log |\sigma(\alpha) - \beta|_v. \quad (7)$$

The product formula says $\Gamma = 0$. At $v \notin S$, relative S -integrality and the ultrametric inequality make the local average in (7) exactly $\log \max\{|\alpha|_v, |\beta|_v\}$. Only the finitely many places in S require estimates.

At a non-archimedean $v \in S$, Lemma 6.1 bounds the number of roots of unity that can lie in a fixed small ball, and (5) controls those exceptional conjugates. At an archimedean place, $\rho_{\mathbf{f},v}$ is normalized Haar measure on the circle $|z|_v = |\alpha|_v$; Jensen's formula gives

$$\int \log |z - \beta|_v d\rho_{\mathbf{f},v} = \log \max\{|\alpha|_v, |\beta|_v\}. \quad (8)$$

Quantitative equidistribution is applied to a logarithm truncated at $\varepsilon \asymp [K(\alpha) : K]^{-1/2}$, while (5) controls the removed singularity. The resulting error is $O(h(\beta)[K(\alpha) : K]^{-1/3})$.

Consequently, if $[K(\alpha) : K]$ were large, (7) would be arbitrarily close to the positive quantity (6), contradicting $\Gamma = 0$ and the uniform lower bound from (2). Hence $[K(\alpha) : K]$ is bounded. Proposition 4.1 also bounds $h(\alpha)$, so Northcott finishes the finiteness argument.

Power-map model and scope

For the single generator $f(z) = z^d$, the nonzero preperiodic points are roots of unity, (3) has $b = 1$, and (8) is the familiar unit-circle Jensen identity. The full proof replaces this rigid cyclotomic picture by a finite family of binomial coefficient classes and makes every height and equidistribution constant independent of the chosen semigroup word.

The result concerns monomials on $\mathbb{P}^1$ with possibly negative exponents and the paper's prefix-collision notion of semigroup preperiodicity. It does not prove the stated conjecture for arbitrary rational semigroups; the binomial structure and explicit equilibrium circles are used decisively.