On birational geometry of the space of parametrized rational curves in Grassmannians

Atsushi Ito

arXiv:1409.0219

http://arxiv.org/abs/1409.0219v2

The compactification and its birational models

Let k be algebraically closed, V an n-dimensional vector space, and fix $d \geq 0$, $0 \leq r \leq n-1$, and $s = n - r$. The Quot scheme

$$R = \mathrm{Quot}_{\mathbb{P}^1}(V \otimes \mathcal{O}_{\mathbb{P}^1}; \mathrm{rk} = r, \deg = d)$$

compactifies degree-d maps $\mathbb{P}^1 \to \mathrm{Gr}(V, r)$. Ito's goal is not only to find its effective and movable cones, but to construct every chamber as a modular compactification.

For $m \geq \lceil d/s \rceil$, condition $(\star_m)$ on a family $\mathcal{E} \to V \otimes \mathcal{O}_{\mathbb{P}^1 \times T}$ requires that $\mathcal{E}$ be locally free of rank s and fiber degree $-d$, that $R^1\pi_*(\mathcal{E}(m-1)) = 0$, and that the fiber map

$$H^0(\mathcal{E}_t(m)) \longrightarrow V \otimes H^0(\mathcal{O}_{\mathbb{P}^1}(m))$$

be injective.

Main theorem (Theorems 1.3, 1.5, 1.6 and Corollary 1.7)**.** *For every $m \geq \lceil d/s \rceil$ there is a smooth projective fine moduli space R_m for $(\star_m)$, with a natural birational map $R \dashrightarrow R_m$. If $d \geq 1$, $r \leq n-2$, and $\lfloor d/s \rfloor + 1 \leq m \leq d-1$, then R_m is a small $\mathbb{Q}$-factorial modification of R. Writing*

$$\alpha = c_1(\mathcal{B}_d) - c_1(\mathcal{B}_{d-1}), \qquad \beta = c_1(\mathcal{B}_{d-1}), \qquad \mathcal{B}_j = \pi_*\mathcal{B}(j),$$

for the universal quotient $\mathcal{B}$, its nef cone is

$$\mathrm{Nef}(R_m) = \mathbb{R}_{\geq 0}(-(d-m)\alpha + \beta) + \mathbb{R}_{\geq 0}(-(d-m-1)\alpha + \beta).$$

The base chamber is $\mathrm{Nef}(R) = \mathbb{R}_{\geq 0}\alpha + \mathbb{R}_{\geq 0}\beta$*. For $r = 0, 1$,*

$$\mathrm{Mov}(R) = \mathrm{Nef}(R) \ \cup \bigcup_{m=\lfloor d/s \rfloor + 1}^{d-1} \mathrm{Nef}(R_m).$$

For $2 \leq r \leq n-2$, the dual Quot scheme R' and the models $R'_{m'}$, where $\lfloor d/r \rfloor + 1 \leq m' \leq d-1$, supply the chambers on the other side of the ray α, with

$$\mathrm{Nef}(R') = \mathbb{R}_{\geq 0}\alpha + \mathbb{R}_{\geq 0}(2d\alpha - \beta), \qquad \mathrm{Nef}(R'_{m'}) = \mathbb{R}_{\geq 0}((d+m')\alpha - \beta) + \mathbb{R}_{\geq 0}((d+m'+1)\alpha - \beta).$$

In this range the complete union is

$$\mathrm{Mov}(R) = \mathrm{Nef}(R) \cup \mathrm{Nef}(R') \cup \bigcup_{m=\lfloor d/s \rfloor + 1}^{d-1} \mathrm{Nef}(R_m) \cup \bigcup_{m'=\lfloor d/r \rfloor + 1}^{d-1} \mathrm{Nef}(R'_{m'}).$$

Consequently R is a Mori dream space; moreover, it is log Fano.

The extreme rays are explicit. For $r = 0, 1$,

$$\mathrm{Mov}(R) = \mathbb{R}_{\geq 0}\alpha + \mathbb{R}_{\geq 0}(-(d - \lfloor d/s \rfloor - 1)\alpha + \beta),$$

whereas for $2 \leq r \leq n-2$ the first ray is replaced by $\mathbb{R}_{\geq 0}((d + \lfloor d/r \rfloor + 1)\alpha - \beta)$. The cases $d = 0$ or $r = n-1$ reduce respectively to a Grassmannian or a projective space.

Ingredient 1: a modular zero locus

Set $V_m = V \otimes H^0(\mathcal{O}_{\mathbb{P}^1}(m))$ and $G_m = \mathrm{Gr}((m+1)s - d, V_m)$. Multiplication by $H^0(\mathcal{O}(1))$ gives a bundle map on $G_{m-1} \times G_m$; its zero locus is R_m. The two tautological subbundles reconstruct a bundle $\mathcal{K} \to V \otimes \mathcal{O}$ satisfying $(\star_m)$. Conversely, cohomology and base change send any $(\star_m)$-family to the two Grassmannians, and Strømme's functorial resolution on $\mathbb{P}^1$ forces the map to factor through the zero locus. This proves the fine-moduli property. An affine presentation of such resolutions is a principal $\mathrm{GL}(s+d) \times \mathrm{GL}(d)$-bundle over R_m; its open parameter space is smooth and irreducible. A common nonempty open with $R_d \simeq R$ then gives the birational map.

Ingredient 2: determinantal wall crossings

The projections $R_m \to G_{m-1}$ and $R_m \to G_m$ are relative Grassmannians over degeneracy loci X^i_m. Determinantal dimension formulas show that these loci are irreducible and Cohen–Macaulay; the next stratum has codimension at least two, so Serre's criterion gives normality. The same calculation shows that, for the interior range of m, both projections are isomorphisms in codimension one. Thus adjacent models

$$R_{m+1} \longrightarrow X^0_m \longleftarrow R_m$$

are related by a small modification. This is the hardest technical point: the fiber ranks and exceptional-stratum dimensions must be exact to distinguish small, divisorial, and fiber-type boundary contractions.

Because the Picard rank is two, the pullbacks of the two Plücker classes span $\mathrm{Nef}(R_m)$. The recurrence $c_1(\mathcal{B}_m) = -(d-1-m)\alpha + \beta$ identifies successive chamber rays. As a toy model, the diagram above resembles the two small resolutions of a determinantal variety: crossing a shared nef ray changes the resolution without changing divisors. The full proof additionally needs the endpoint contractions and the moduli interpretation of every R_m.

Ingredient 3: the missing half and the corollary

For $2 \le r \le n-2$, dualizing on the locally free locus gives an SQM $R \dashrightarrow R'$. A Chern-class computation identifies its natural basis as $\alpha' = \alpha$, $\beta' = 2d\alpha - \beta$; applying the same construction to R' supplies the mirror sequence of chambers. The outer contractions locate the effective-cone edges, so the listed nef cones exhaust the movable cone. Finally, the anticanonical class lies in $\mathrm{Mov}(R) \cap \mathrm{Eff}(R)^\circ$. On a suitable smooth SQM it is nef and big, yielding the separate log Fano conclusion. The note suppresses the full formulas for every degeneracy stratum and endpoint fiber; these are essential in the paper but not needed to see the proof's architecture.