Torelli theorem for the product of moduli spaces of vector bundles over a curve

Indranil Biswas, Arijit Dey, Anubhab Pahari

Indranil Biswas, Arijit Dey, and Anubhab Pahari, arXiv:2609.10764v1 (2026), 13 pages
http://arxiv.org/abs/2609.10764v1

Can a product remember all of its curves?

For a smooth complex projective curve X, write $M_X(r,d)$ for the coarse moduli space of semistable vector bundles of rank r and degree d, and $M_X(r,L)$ for the fixed-determinant fiber. A single $M_X(r,L)$ determines X under the usual Torelli hypotheses; the paper asks whether taking products can obscure the individual factors.

Main theorem (Biswas–Dey–Pahari, Theorem 1.1/3.3)**.** *Let $X_1,\dots,X_m$ and $X'_1,\dots,X'_n$ be irreducible smooth projective complex curves of genus at least 4, and let $r_i, r'_j \geq 2$. Then*

$$\prod_{i=1}^{m} M_{X_i}(r_i,d_i) \simeq \prod_{j=1}^{n} M_{X'_j}(r'_j,d'_j)$$

if and only if $m=n$ and, after a permutation σ of the factors,

$$X_i \simeq X'_{\sigma(i)}, \qquad r_i = r'_{\sigma(i)}, \qquad d_i \equiv \pm d'_{\sigma(i)} \pmod{r_i}$$

for every i.

The statement is stronger than recovering only the product of the Jacobians: the abstract product variety recovers each curve and the discrete bundle data, modulo the familiar dualization ambiguity.

Ingredient 1: the determinant map isolates the rigid fibers

The determinant morphism

$$\det : M_X(r,d) \longrightarrow \mathrm{Pic}^d(X), \qquad E \longmapsto \bigwedge^r E,$$

is identified in Lemma 3.1 as the Albanese morphism. Its fiber over L is $M_X(r,L)$, whose Albanese is trivial. Consequently an isomorphism between the two products in the theorem induces an isomorphism of their Albanese varieties and a compatible square of determinant maps. Restricting that square over a chosen point gives

$$\prod_i M_{X_i}(r_i,L_i) \simeq \prod_j M_{X'_j}(r'_j,L'_j).$$

This passage removes the Jacobian directions and leaves factors with Picard number one and trivial Pic^0.

Ingredient 2: extremal nef rays recognize product factors

Theorem 2.6 supplies the new structural input. If normal projective varieties V_i, W_j satisfy $\rho(V_i) = \rho(W_j) = 1$ and $\mathrm{Pic}^0(V_i) = \mathrm{Pic}^0(W_j) = 0$, then

$$\prod_i V_i \simeq \prod_j W_j \quad \Longrightarrow \quad m = n \ \text{ and } \ V_i \simeq W_{\sigma(i)}.$$

Indeed, discreteness of the Picard groups splits N^1 of a product, and its nef cone becomes an orthant. Its extremal rays are exactly the pullbacks of the one-dimensional nef cones of the factors. A product isomorphism preserves these rays, so it determines a permutation.

Numerical proportionality on a matched ray is promoted to an isomorphism of line bundles after clearing the finite torsion in $\mathrm{Pic}^\tau / \mathrm{Pic}^0$. Choose very ample representatives on the two matched factors. The complete linear system of either pullback depends on only that coordinate, and its image is the embedded factor itself. The product isomorphism identifies the two images, hence identifies the factors. Turning a cone-theoretic permutation into an actual factor isomorphism is the proof's hardest step.

Ingredient 3: fixed-determinant Torelli finishes the match

The cited Picard theorem for $M_X(r, L)$ gives precisely the hypotheses of Theorem 2.6. Thus the fixed-determinant product isomorphism pairs the spaces $M_{X_i}(r_i, L_i)$ with $M_{X'_{\sigma(i)}}(r'_{\sigma(i)}, L'_{\sigma(i)})$. The established Torelli theorem for a single such moduli space then recovers

$$X_i \simeq X'_{\sigma(i)}, \quad r_i = r'_{\sigma(i)}, \quad \deg L_i \equiv \pm \deg L'_{\sigma(i)} \pmod{r_i},$$

which is the advertised classification. Conversely, these conditions give factorwise isomorphisms, so the criterion is genuinely if and only if.

Stacks, automorphisms, and scope

For fixed-determinant moduli stacks, an isomorphism descends through the good-moduli-space maps to an isomorphism of the coarse products; the same four conditions follow, and the cited single-factor stack theorem gives the converse (Theorem 3.4). If the distinct factor types are pairwise non-isomorphic and occur with multiplicities n_i, the nef-ray argument also yields

$$\mathrm{Aut}\left(\prod_i M_i^{n_i}\right) \simeq \prod_i \big(\mathrm{Aut}(M_i)^{n_i} \rtimes S_{n_i}\big),$$

and analogously for the stacks.

The theorem is over $\mathbb{C}$, assumes genus at least 4 and rank at least 2, and concerns semistable-bundle moduli spaces. The proof uses the special Picard and Torelli properties of fixed-determinant fibers; it does not assert a comparable unique-factorization theorem for arbitrary projective varieties.