

The equidistribution of small point for strongly regular pairs of polynomial maps

Chong Gyu Lee

arXiv:1203.1224

<http://arxiv.org/abs/1203.1224v2>

From affine dynamics to arithmetic measures

Let K be a number field and let $f: \mathbb{A}^n \rightarrow \mathbb{A}^n$ be a polynomial automorphism. Its extensions $f, f^{-1}: \mathbb{P}^n \dashrightarrow \mathbb{P}^n$ are generally rational rather than morphisms, so the usual construction of a canonical adelic metric for a polarized projective endomorphism does not apply. The paper replaces a single map by a pair. A pair $S = \{F, G\}$ is *strongly regular* when, in particular, $I(F) \cap I(G) = \emptyset$, both maps are algebraically stable and commute, their relevant height-growth constants are finite, and

$$\deg(F \circ G) < \min\{\deg F, \deg G\}.$$

The decisive case also has $\deg F = \deg G = d \geq 2$.

Main theorem (Lee, Theorem A). *Let $f: \mathbb{A}^n \rightarrow \mathbb{A}^n$ be a regular polynomial automorphism over K , meaning $I(f) \cap I(f^{-1}) = \emptyset$. For every place v of K there is an f -invariant probability measure $\mu_{f,v}$ on the Berkovich space $(\mathbb{P}_{\mathbb{C}_v}^n)^{\text{an}}$ such that, for any sequence of periodic points x_m meeting each proper subvariety only finitely often,*

$$\frac{1}{[K(x_m) : K]} \sum_{y \in \text{Gal}(\bar{K}/K)x_m} \delta_y \implies \mu_{f,v}.$$

Here the displayed denominator is the size of the Galois orbit (equivalently the residue degree for the associated closed point).

Thus one canonical measure governs algebraic periodic points simultaneously at archimedean and nonarchimedean places. At an archimedean place it agrees with the Green-current equilibrium measure identified by Dinh–Sibony.

Ingredient I: two escape rates cover affine space

For a polynomial map F define the local escape rate

$$G_{F,v}(x) = \lim_{m \rightarrow \infty} d^{-m} \log^+ \|F^m(x)\|_v.$$

Theorem 3.2 constructs v -adic open sets V_F, V_G with $V_F \cup V_G = \mathbb{A}^n$ on which respectively $G_{F,v}$ and $G_{G,v}$ equal $\log^+ \|x\|_v + O(1)$. The proof homogenizes F and G . Joint regularity gives, by the Nullstellensatz, identities expressing powers of every coordinate through the homogeneous components of F, G , while the degree drop in $F \circ G$ supplies a competing estimate. With carefully chosen constants these estimates rule out a point lying outside both regions and make each region forward invariant.

The same inequalities give uniform, not merely pointwise, convergence of the normalized iterates on their regions (Proposition 3.3); this is what is needed for continuity and semipositivity of the eventual metric. An elimination argument shows that the pair has good reduction away from finitely many places (Propositions 4.3–4.5), where the metric is exactly the standard one. The archimedean analogue is Theorem 5.1.

Ingredient II: a polarized sequence of morphisms

Because $I(F) \cap I(G) = \emptyset$ and the degrees agree, the map

$$\phi_m: \mathbb{P}^n \longrightarrow \mathbb{P}^{2n}, \quad x \longmapsto (F^m(x), G^m(x))$$

is a genuine morphism and $\phi_m^* \mathcal{O}_{\mathbb{P}^{2n}}(1) \simeq \mathcal{O}_{\mathbb{P}^n}(d^m)$. Pulling back the standard metrics and taking d^m -th roots produces metrics on $\mathcal{O}_{\mathbb{P}^n}(1)$. On $\mathbb{A}^n$ their potentials tend to $\max(G_{F,v}, G_{G,v}) - \log^+ \|x\|_v$.

Theorem 6.5 proves uniform convergence also along the hyperplane at infinity. There algebraic stability sends the appropriate strata between the two indeterminacy loci, giving telescoping Cauchy bounds. On the affine chart, the auxiliary pair $\Phi = (F, G)$ and $\Psi = (G, F)$ on $\mathbb{A}^{2n}$ packages the maximum of the two escape rates into the uniform-convergence statement from Ingredient I. The limit is therefore a semipositive adelic metric $\overline{L}_S$ on $\mathcal{O}(1)$. Yuan's equidistribution theorem now immediately yields Theorem B: every generic sequence small for $h_{\overline{L}_S}$ equidistributes to the Chambert–Loir measure $c_1(\overline{L}_S)_v^n$.

Ingredient III: periodic points are generic and small

For a regular automorphism, put $\ell_+ = \dim I(f) + 1$ and $\ell_- = \dim I(f^{-1}) + 1$. Proposition 7.2 gives

$$\ell_+ + \ell_- = n, \quad (\deg f)^{\ell_-} = (\deg f^{-1})^{\ell_+}.$$

Consequently $\{f^{\ell_-}, f^{-\ell_+}\}$ is a same-degree strongly regular pair (Lemma 7.3), so Ingredient II supplies its adelic metric.

Theorem C supplies the missing genericity: $\text{Per}(f)$ is Zariski dense. After reduction at a good finite place, every point over the algebraic closure of the finite residue field is periodic. These points lie in equalizers of the two iterates above. A finiteness theorem for equalizers of morphisms of unequal degree forces infinitely many horizontal components in the arithmetic model; their generic-fibre points are periodic and Zariski dense. One may therefore choose a generic periodic sequence.

Such points have $h_{\overline{L}_S} = 0$. Zhang's successive-minima inequality, combined with their Zariski density, gives $h_{\overline{L}_S}(\mathbb{P}^n) = 0$; hence the sequence is small. Applying Theorem B proves Theorem A, and applying the same limit to $f(x_m)$ proves f -invariance of $\mu_{f,v}$ (Corollaries 7.5–7.6).

A model: a plane Hénon map

For $f(x, y) = (y, p(y) - ax)$ with $a \neq 0$ and $\deg p = d \geq 2$, the extensions of f and f^{-1} have disjoint points of indeterminacy at infinity. Here $\ell_+ = \ell_- = 1$, so no degree-balancing iterates are needed. Forward and backward escape rates occupy complementary regions, and $\max(G_f, G_{f^{-1}})$ already displays the two-sided potential used in the general construction. Higher dimension adds unequal forward/backward degrees and more complicated infinity strata, exactly the difficulties handled by Proposition 7.2 and Theorem 6.5.

Scope and technical bottleneck

The theorem assumes regularity, algebraic coefficients, and a generic sequence; it does not say that every enumeration of periodic points equidistributes, nor does it give an effective rate. The deepest internal step is not the final appeal to Yuan but the construction of a globally semipositive adelic metric from rational dynamics: local uniform convergence must survive indeterminacy at infinity and be standard at almost every place. The paper establishes precisely those two facts before invoking the general equidistribution theorem.