# Canonical heights for Hénon maps

Patrick Ingram

arXiv:1111.3609

http://arxiv.org/abs/1111.3609v1

## Problem and setup

A Hénon map

$$\varphi(x,y) = (ay,\ x + f(y)), \qquad d = \deg f \geq 2,$$

is an affine-plane automorphism that does not extend to a projective-surface endomorphism. Its two escape rates define

$$\widehat{h}^{\pm}_{\varphi}(P) = \lim_{N\to\infty} d^{-N} h(\varphi^{\pm N}(P)), \qquad \widehat{h}_{\varphi} = \widehat{h}^{+}_{\varphi} + \widehat{h}^{-}_{\varphi}.$$

The paper studies their variation in families and periodic specialization of a generic section.

**Main theorem** (Theorems 1.1–1.3)**.** *Let $C$ be a smooth projective curve over a number field $K$, put $F = K(C)$, and let $P \in \mathbf{A}^2(F)$. For a Hénon map over $F$ there are $\mathbf{Q}$-divisors $D^+, D^-$ on $C$ such that*

$$\widehat{h}^{\pm}_{\varphi_t}(P_t) = h_{D^{\pm}}(t) + O(1).$$

*If moreover $\varphi = (y, x + f(y))$ is non-isotrivial, then either $P$ is periodic over $F$, or*

$$\{t \in C(\overline{K}) : P_t \text{ is periodic for } \varphi_t\}$$

*has bounded height. The zero-height input is that, for such a non-isotrivial map over a function field, $\widehat{h}_{\varphi}(P) = 0$ holds exactly at periodic points.*

Thus an affine dynamical height is controlled by a divisor height on the base. For $K$-rational parameters, Northcott turns bounded height into finiteness.

## Proof mechanism

**1. Local escape regions.** At each place, invariant regions $B^{\pm}_v$ make the degree-$d$ term dominate. On $B^+_v$, $\widehat{\lambda}^+_{v,\varphi}(x,y) = \log|y|_v$ up to a uniform archimedean error, and it vanishes exactly when no forward iterate enters $B^+_v$. The backward statement is symmetric.

**2. Turn local escape into divisors on $C$.** For a place $\beta$ of $F$, set

$$D^{\pm}(\varphi, P) = \sum_{\beta \in C(\overline{K})} \widehat{\lambda}^{\pm}_{\beta,\varphi}(P)[\beta].$$

The functional equation makes every nonzero coefficient rational. Lemma 5.2 compares specialized canonical and Weil local heights near $\mathrm{Supp}(D^{\pm})$; Lemmas 5.3–5.4 bound both at zero-coefficient poles and elsewhere. Summation over places gives the global $O(1)$ formulas.

**3. Rule out nonperiodic points of height zero.** For $d \geq 3$, the $v$-adic filled Julia set lies in unit disks around root pairs. At each genuinely bad place, a large subset of any zero-height set clusters in three consecutive $y$-coordinates. Iterating this selection over all bad places produces two rational points whose nonzero differences violate the product formula. With no such place, a root translation makes $f$ constant-field defined, the isotrivial alternative.

For $d = 2$, Lemma 2.2 conjugates to $(y, x + y^2 + b)$, adding only finitely many bad places. Theorem 1.4 then gives $B$ and $\varepsilon > 0$ such that every point is either periodic of period at most $B$ or has $\widehat{h}_{\varphi}(P) \geq \varepsilon \max\{h(b), 1\} > 0$. so height zero again forces periodicity.

**4. Specialize.** For nonperiodic $P$, rigidity gives $\widehat{h}_{\varphi}(P) > 0$, so $D = D^+ + D^-$ is ample and $\widehat{h}_{\varphi_t}(P_t) = h_D(t) + O(1)$. Periodic fibers have height zero, hence bounded parameter height.

## Toy model, difficulty, and scope

Locally, $y = u^{-m}\cdot$(a unit) models why $[\beta]$ receives escape coefficient $m$ and specialization grows like $m \log|u(t)|_v^{-1}$; uniformity at every place is the real difficulty. The hardest step is the two-dimensional product-formula packing. Zero-height rigidity requires $a = 1$ and non-isotriviality, and bounded height over $\overline{K}$ is not finiteness without a degree bound. The period conjecture is separate.