

Reductions Modulo Primes of Systems of Polynomial Equations and Algebraic Dynamical Systems

Carlos D’Andrea, Alina Ostafe, Igor E. Shparlinski, Martín Sombra

arXiv:1505.05814v2

<http://arxiv.org/abs/1505.05814v2>

An effective good-reduction theorem

Put $\mathbf{X} = (X_1, \dots, X_m)$. For $F \in \mathbf{Z}[\mathbf{X}]$ let $h(F)$ be the logarithm of its largest absolute coefficient. The paper’s basic question is unusually rigid: if an integral system has exactly T distinct complex zeros, when does reduction modulo p retain exactly T distinct geometric zeros?

Main theorem (D’Andrea–Ostafe–Shparlinski–Sombra, Theorems 2.1 and 4.3). *Let $F_1, \dots, F_s \in \mathbf{Z}[\mathbf{X}]$ have a common zero set in $\mathbf{C}^m$ consisting of T distinct points, and set $d = \max_i \deg F_i$ and $h = \max_i h(F_i)$. There is $A \geq 1$ such that*

$$\log A \leq (11m + 4)d^{3m+1}h + (55m + 99)\log((2m + 5)s)d^{3m+2}, \quad (1)$$

and, for every prime $p \nmid A$, the reduced system has exactly T distinct zeros in $\overline{\mathbf{F}}_p^m$.

Consequently, if $\mathbf{F} = (F_1, \dots, F_m)$ has degree at most $d \geq 2$ and height at most h , with $m \geq 2$, and has only finitely many k -periodic points over $\mathbf{C}$, then some $A_k \geq 1$ satisfies

$$\log A_k \ll_{d,h,m} d^{k(3m+2)},$$

and $\mathbf{F} \bmod p$ has at most d^{km} k -periodic points over $\overline{\mathbf{F}}_p$ whenever $p \nmid A_k$.

Here “ k -periodic” means $\mathbf{F}^{(k)}(x) = x$; k need not be the least period. The first statement preserves the exact geometric count, not merely an upper bound or a count of $\mathbf{F}_p$ -rational points.

Ingredient I: flat closure fixes the total degree

Let $V \subset \mathbf{A}_{\mathbf{Q}}^m$ be the zero-dimensional reduced variety cut out by the F_i , and take its closure $\overline{V} \subset \mathbf{P}_{\mathbf{Z}}^m$. The authors prove that $\overline{V}$ is flat, reduced, of pure relative dimension zero, and has no irreducible component at infinity. Invariance of the Hilbert polynomial in this projective flat family therefore gives

$$\deg \overline{V}_p = T \quad (2)$$

for every fibre. Equation (2) alone is insufficient: the fibre can acquire multiplicity or points at infinity, while the equations $F_i \bmod p$ can have additional affine zeros.

Ingredient II: one eliminant controls all three defects

For $\mathbf{U} = (U_0, \dots, U_m)$ and $L^{\text{aff}} = U_0 + U_1X_1 + \dots + U_mX_m$, the eliminant of V factors as

$$E_V(\mathbf{U}) = \lambda \prod_{\xi \in V(\overline{\mathbf{Q}})} (U_0 + \xi_1U_1 + \dots + \xi_mU_m). \quad (3)$$

Arithmetic Bézout bounds the heights of the points and hence $h(E_V)$. An effective arithmetic Nullstellensatz then supplies $\alpha, N \geq 1$ with

$$\alpha E_V^N \in (F_1, \dots, F_s, L^{\text{aff}}), \quad (4)$$

while a resultant supplies β for which $E_V \bmod p$ is squarefree and has degree T in U_0 whenever $p \nmid \beta$.

Take $A = \alpha\beta$. The β condition makes the degree- T fibre in (2) reduced and affine. The α condition says that every affine zero ξ of all $F_i \bmod p$ contributes the distinct linear divisor $L^{\text{aff}}(\mathbf{U}, \xi)$ of $E_V \bmod p$. Thus there are at most T such zeros, while the flat fibre already supplies T of them. This proves the exact equality and explains why both the eliminant and the Nullstellensatz certificate are indispensable.

Ingredient III: encode dynamics by polynomial equations

The degree and height estimates for composition give

$$\deg F_i^{(k)} \leq d^k, \quad h(F_i^{(k)}) \leq h \frac{d^k - 1}{d - 1} + d(d + 1) \frac{d^{k-1} - 1}{d - 1} \log(m + 1). \quad (5)$$

The k -periodic locus is exactly

$$V_k = Z(F_1^{(k)} - X_1, \dots, F_m^{(k)} - X_m). \quad (6)$$

Its assumed finiteness lets Theorem 2.1 transfer its complex cardinality to good reductions; Bézout and (5) give $\#V_k(\mathbf{C}) \leq d^{km}$ and the stated bound for A_k . Rational maps require an extra variable and an equation excluding every denominator encountered before time k ; this is why the rational-map bound in Theorem 4.2 is weaker.

Toy model and the orbit-intersection mechanism

For $F_i = X_i^d$, equation (6) becomes $X_i^{d^k} - X_i = 0$. If $p \nmid d^k - 1$, it has exactly d^{km} geometric solutions, so the periodic point estimate is sharp. This model removes all height growth and shows that the exponent km comes from the degree of the fixed-point equations, not from a loss in the reduction argument.

The same reduction theorem also drives Theorem 5.3. If the iterates of a rational system R *generically escape* a variety W —for every $r \geq 1$, the iterate $R^{(r)}$ is well-defined and only finitely many complex z satisfy $z, R^{(r)}(z) \in W$ —then, for $N \geq \exp(c/\varepsilon)$ and outside primes dividing an integer B with $\log B \leq \exp(c/\varepsilon)$, any nonrepeating length- N orbit spends at most an ε proportion of its time in W . Too many visits give, by a repeated-gap lemma, many distinct pairs separated by one $r = O(1/\varepsilon)$. Bézout bounds the corresponding finite complex incidence set, and Theorem 2.1 transfers that bound modulo p , contradicting the repeated-gap count. Theorem 6.2 replaces generic escape by a uniform complex-orbit bound L and needs only $N \geq 2L/\varepsilon + 1$.

Hard point and scope

The difficult step is quantitative simultaneity: the same explicitly bounded integer must prevent nonreduced fibres, escape to infinity, and extraneous solutions. The bounds are effective but very large and not asserted to be optimal. The zero-dimensional and finiteness hypotheses are essential. Orbit-frequency results additionally require their stated escape or uniform Mordell–Lang hypothesis and sufficiently long nonrepeating orbits; they do not give an unconditional uniform dynamical Mordell–Lang theorem.