Calabi-Yau Threefolds Fibred by Mirror Quartic K3 Surfaces

Charles F. Doran, Andrew Harder, Andrey Y. Novoseltsev, Alan Thompson

arXiv:1501.04019

http://arxiv.org/abs/1501.04019v2

A one-dimensional modular description

Set

$$M_2 = H \oplus E_8 \oplus E_8 \oplus \langle -4 \rangle.$$

An M_2-polarized K3 family carries this rank-19 lattice as a primitive constant subsystem of the fibrewise Néron–Severi groups, containing an ample class. Its generalized functional invariant is the classifying map $g : U \to \mathcal{M}_{M_2}$, where $\mathcal{M}_{M_2} \simeq \Gamma_0(2)^+ \backslash \mathfrak{H}$ compactifies to $\mathbb{P}^1$. A fundamental family is the minimal resolution of

$$\lambda w^4 + xyz(x + y + z - w) = 0 \subset \mathbb{P}^3,$$

whose special parameters are $\lambda = 0, 1/256, \infty$.

Main theorem (Theorem 2.3 and Corollary 2.8)**.** *Let $\mathcal{X}_U \to U$ be a non-isotrivial M_2-polarized family over a quasi-projective curve, with general Néron–Severi lattice isometric to M_2. Then $\mathcal{X}_U$ is uniquely determined, up to isomorphism, by g; in particular it is the pullback of the fundamental family.*

Moreover, every Calabi–Yau threefold fibred by K3 surfaces whose smooth part has this polarization and general Néron–Severi lattice is birational to a threefold obtained from such a pullback. Conversely, the smooth threefolds produced by the paper's numerical and resolution conditions are Calabi–Yau.

Here *Calabi–Yau* means smooth projective, with trivial canonical bundle and $H^1(\mathcal{O}) = 0$. The theorem is a classification up to birational equivalence, not an assertion that every ramified pullback is smooth.

The explicit ramification criterion

Let $g : \mathbb{P}^1 \to \mathcal{M}_{M_2} \simeq \mathbb{P}^1$ have degree n. Write $[x_1, \ldots, x_k]$, $[y_1, \ldots, y_l]$, and $[z_1, \ldots, z_m]$ for its ramification profiles above $0, \infty, 1/256$, and let $r = \sum_{g(p) \notin \{0, 1/256, \infty\}} (e_p - 1)$. If $\overline{\mathcal{X}}_g$ is the normalization of the pullback, then Proposition 2.4 says

$$\omega_{\overline{\mathcal{X}}_g} \simeq \mathcal{O} \quad \Longleftrightarrow \quad k + l + m - n - r = 2$$

together with exactly one of

$$l = 2, \quad y_1, y_2 \in \{1, 2, 4\}; \qquad \text{or} \qquad l = 1, \quad y_1 = 8.$$

Under these conditions Proposition 2.5 constructs a projective birational model $\mathcal{X}_g$ with trivial canonical sheaf and at worst $\mathbb{Q}$-factorial terminal singularities. Its possible remaining singularities lie over $g^{-1}(1/256)$; if g is unramified there, then $\mathcal{X}_g$ is smooth. Kollár torsion-freeness gives $H^1(\mathcal{X}_g, \mathcal{O}) = 0$, completing the Calabi–Yau implication.

Ingredient 1: why the classifying map is enough

On simply connected pieces of U, Ehresmann triviality supplies compatible markings, and global Torelli identifies two families with the same g. Different global gluings would therefore differ by a fibrewise automorphism fixing M_2. On a general fibre, its action on the rank-3 orthogonal complement $M_2^{\perp}$ is irreducible. Nikulin's constraints force its order to be 1 or 2; order 2 would act as -1 on $M_2^{\perp}$, but cannot act trivially on the discriminant group $A_{M_2^{\perp}} \simeq \mathbb{Z}/4\mathbb{Z}$. Thus the gluing automorphism is trivial. This is precisely why the analogous statement fails for the nearby lattice M_1, whose fibres admit a polarization-preserving antisymplectic involution.

Ingredient 2: adjunction and crepant local models

Riemann–Hurwitz and adjunction first produce $k + l + m - n - r - 2 = 0$. Normalization above the multiplicity-four fibre at ∞ contributes $\sum_i (y_i/\gcd(y_i, 4) - 1)$, leaving only the two alternatives displayed above. The resolution is then checked fibre by fibre. Above 0, base change creates curves of cA_{x_i-1} singularities resolved torically. Above ∞, the cases $y_i = 1, 2, 4, 8$ require separate crepant subdivisions. Above $1/256$, ramification order z_i turns an isolated A_1 point into an isolated cA_{z_i-1} terminal singularity. Tracking projectivity, crepancy, and factoriality through these local models is the hardest technical part of the classification.

Ingredient 3: Hodge numbers and deformations

For the principal case $l = 2$, Proposition 3.5 counts invariant divisor classes and components of resolved singular fibres:

$$h^{1,1}(\mathcal{X}_g) = 20 + \sum_{i=1}^{k} (2x_i^2 + 1) + c_1 + c_2,$$

where $c_j = 30, 10, 0$ for $y_j = 1, 2, 4$, respectively. The transcendental rank-3 local system and its monodromy give Corollary 3.9,

$$h^{2,1}(\mathcal{X}_g) = k + \frac{m_{\mathrm{odd}} - n}{2},$$

where m_{odd} counts odd z_i. If g is unramified above $1/256$, this simplifies to $h^{2,1} = k = r$.

These formulas also explain deformation theory. Under $l = 2$ and no ramification above $1/256$, every small deformation of $\mathcal{X}_g$ comes from deforming g while preserving the profiles above 0 and ∞ (Proposition 4.1). Proposition 4.3 splits arbitrary other monodromy cycles into transpositions, showing that g can be deformed to have only simple ramification away from the three special values.

As a concrete check, the mirror quintic corresponds to

$$g(s : t) = \frac{As^5}{t(s - t)^4}, \quad (k, l, m, n, r) = (1, 2, 5, 5, 1),$$

and the formulas recover $(h^{1,1}, h^{2,1}) = (101, 1)$. The example is a useful toy model for the cover construction; the full proof additionally needs the Torelli rigidity and every singular-fibre resolution above.