# Singularities of pluri-fundamental divisors on Gorenstein Fano varieties of coindex 4

Jinhyung Park

arXiv:2206.00881

http://arxiv.org/abs/2206.00881v1

## Pluri-fundamental divisors in coindex four

Work over **C**. Let $X$ be a Gorenstein canonical Fano variety of dimension $n \geq 4$ and coindex four. Its fundamental divisor $H$ is the ample Cartier divisor determined by

$$-K_X \sim (n-3)H.$$

The question is whether a general $Y_m \in |mH|$ inherits canonical or terminal singularities from $X$.

**Main theorem** (Park, Theorem 1.1)**.** *With $X$ and $H$ as above:*

(1) *If $h^0(X,H) \geq n-2$, then a general $Y_m \in |mH|$ has canonical singularities for every $m \geq 1$.*
(2) *If, in addition, $X$ is terminal, then a general $Y_m$ is terminal for every $m \geq 1$, except for the three cases $(n,m) = (4,1), (4,2), (4,3)$, for which this assertion is not made.*
(3) *If $X$ is smooth, then $h^0(X,H) \geq n-2$, and a general $Y_m$ is terminal for every $m \geq 1$ except possibly $(n,m) = (4,2)$.*

The uniformity in every multiple $m$ is the point: the proof controls all exceptional divisors over the base locus by one numerical inequality.

## Ingredient 1: cut the base locus down to dimension two

Take a log resolution $f_m \colon X_m \to X$ of $\operatorname{Bs}|mH|$ and write

$$K_{X_m} = f_m^* K_X + \sum_i a_{m,i} E_{m,i}, \qquad |f_m^* mH| = |M_m| + \sum_i r_{m,i} E_{m,i}.$$

All $a_{m,i}, r_{m,i}$ are nonnegative integers. Repeatedly cutting by a general member of $|H|$ gives a "ladder" of purely log terminal pairs, ending in a canonical Calabi–Yau threefold. The assumption $h^0(X,H) \geq n-2$ keeps the ladder moving and shows

$$\dim \operatorname{Bs}|mH| \leq 2 \qquad (m \geq 1).$$

This low-dimensional bound is what makes effective nonvanishing available in the next step.

## Ingredient 2: the decisive discrepancy estimate

Proposition 2.3 proves, for every $m \geq 1$,

$$a_{m,i} \geq \frac{m+n-3}{m}\, r_{m,i} - 1. \tag{$*$}$$

If $(*)$ failed, let $c_0$ be the first coefficient at which the pair becomes log canonical and set $\Delta = (c_0/k)(D_1 + \cdots + D_k)$ for $k > c_0$ and general $D_j \in |mH|$. A minimal log-canonical center $W$ of $(X, \Delta)$ lies in the base locus, hence $\dim W \leq 2$. Subadjunction produces a klt pair $(W, \Gamma)$, while

$$mH - (K_X + \Delta) \sim_{\mathbf{Q}} (m + n - 3 - c_0 m)H$$

is ample. Low-dimensional effective nonvanishing gives a nonzero section of $mH|_W$, and extension from the lc center makes $H^0(X, mH) \to H^0(W, mH|_W)$ surjective. But $W$ lies in the base locus, so this restriction map is zero—the contradiction proving $(*)$.

For a general $Y_m$, the discrepancy along $E_{m,i}$ is $a_{m,i} - r_{m,i}$. If $r_{m,i} \geq 1$, $(*)$ gives

$$a_{m,i} - r_{m,i} \geq \frac{n-3}{m} r_{m,i} - 1 > -1.$$

Thus $(X, Y_m)$ is purely log terminal, and adjunction makes the Gorenstein divisor $Y_m$ canonical.

## Ingredient 3: upgrading canonical to terminal

For $m \geq 4$, $|mH|$ is base point free. Assume therefore $n \geq 5$ and $m \leq 3$. When $m = 1$, $(*)$ makes every discrepancy of $Y_1$ strictly positive. If $m = 2$ or 3 and $Y_m$ were not terminal, the only borderline possibility is an exceptional divisor with $a_{m,i} = r_{m,i} = 1$. Its center must lie in $\operatorname{Sing} X$ and in $\operatorname{Bs}|mH| \subseteq \operatorname{Bs}|H|$. On a common refinement it then has multiplicity at least one for $|H|$, contradicting the already established terminality of a general $Y_1$.

When $X$ is smooth, only the fourfold cases $m = 1, 3$ remain beyond this argument. The case $m = 1$ is Heuberger's anticanonical theorem. For $m = 3$, Park first improves the base-locus bound to one dimension and proves that an isolated singularity of multiplicity at most two is terminal. If $H^4 \geq 2$, restriction to a general anticanonical Calabi–Yau threefold and Kawamata freeness rule out the required multiplicity-three point. In the borderline case $H^4 = 1$, Fujita's classification and a genus-four canonical curve cut out by $H, H, 2H$ control the common base locus of $|2H|$ and $|3H|$; a length-six intersection then excludes the same multiplicity.

## A sharp fourfold and the hard point

Let $Z = X_{2,6} \subset \mathbf{P}(1,1,1,1,2,3)$ be cut out by a general quadric in the four weight-one variables $x_0, \ldots, x_3$ and a general sextic. It is a terminal Gorenstein Fano threefold with a point $p$ at which a general fundamental divisor is singular. For $X = Z \times \mathbf{P}^1$, the fourfold is terminal and Gorenstein, but a general $Y \in |-K_X|$ is singular along $\{p\} \times \mathbf{P}^1$; it is canonical, not terminal. Hence the $(4,1)$ exception in the terminal singular case is real.

The conceptual hard point is $(*)$: it turns a high-dimensional base-locus problem into nonvanishing on a center of dimension at most two. The smooth fourfold $m = 3$ endpoint then needs a separate, much finer multiplicity and canonical-curve argument. The paper leaves the smooth $(4,2)$ case open, and the hypothesis $h^0(X, H) \geq n - 2$ is not known in general for singular coindex-four Fano varieties.