

Projectivity of the moduli space of vector bundles on a curve

Jarod Alper, Pieter Belmans, Daniel Bragg, Jason Liang, Tuomas Tajakka

arXiv:2206.06195

<http://arxiv.org/abs/2206.06195v1>

From a nonseparated stack to a projective variety

Let k be algebraically closed of characteristic zero and let X/k be a smooth, projective, connected curve of genus $g \geq 2$. Write $\mathcal{M}_X^{\text{ss}}(r, d)$ for the stack of slope-semistable vector bundles of rank $r > 0$ and degree d , and $\mathcal{M}_X^{\text{ss}}(r, L)$ for the fixed-determinant stack, where $\deg L = d$. Jordan–Hölder theory identifies the closed points with polystable bundles and makes their isomorphism classes the S -equivalence classes of semistable bundles.

Main theorem (Alper–Belmans–Bragg–Liang–Tajakka, Theorem 5.1). *The good moduli spaces*

$$\mathcal{M}_X^{\text{ss}}(r, d) \longrightarrow M_X^{\text{ss}}(r, d), \quad \mathcal{M}_X^{\text{ss}}(r, L) \longrightarrow M_X^{\text{ss}}(r, L)$$

are projective varieties, of respective dimensions

$$r^2(g-1)+1 \quad \text{and} \quad (r^2-1)(g-1). \tag{1}$$

Their k -points correspond to S -equivalence classes.

The projectivity is classical, but the paper gives a modern construction that separates two tasks: good-moduli-space machinery first produces a proper algebraic space, and determinantal line bundles then prove it is a projective scheme.

The proper good moduli space

Semistability is open, bounded, and irreducible, so the two stacks above are irreducible finite-type algebraic stacks with affine diagonal (Theorem 2.17). Langton’s semistable reduction theorem gives universal closedness. The two valuative test-stack criteria for extending filtrations and pairs of degenerations give Θ -reductivity (Proposition 3.8) and S -completeness (Proposition 3.10). The existence criterion for good moduli spaces then yields separated algebraic spaces; universal closedness makes them proper (Theorem 3.12).

This step also explains the quotient relation. A semistable bundle E specializes to its associated graded $\text{gr } E$, while the closed points are exactly the polystable bundles (Lemma 3.11). Thus two points of the stack have the same image precisely when their graded bundles are isomorphic. The stable locus is nonempty, and the tangent calculation there gives (1).

A line bundle whose sections detect cohomology

Let $p : X \times \mathcal{M}_X(r, d) \rightarrow \mathcal{M}_X(r, d)$ and $q : X \times \mathcal{M}_X(r, d) \rightarrow X$ be the projections, and let $\mathcal{E}_{\text{univ}}$ be the universal bundle. For a vector bundle V on X , define

$$\mathcal{L}_V = (\det Rp_*(q^*V \otimes \mathcal{E}_{\text{univ}}))^\vee. \quad (2)$$

If

$$d \operatorname{rk} V + r \deg V + r \operatorname{rk} V(1 - g) = 0, \quad (3)$$

the perfect complex in (2) has virtual rank zero and its differential has a determinant section s_V . Proposition 4.7 identifies its value:

$$s_V([E]) \neq 0 \iff H^0(X, E \otimes V) = H^1(X, E \otimes V) = 0. \quad (4)$$

The bundle initially lives on the stack. At a closed point $E = \bigoplus_j E_j^{\oplus m_j}$, its stabilizer acts on the fiber by $\prod_j \det(g_j)^{\chi(X, E_j \otimes V)}$. Every E_j has slope d/r , so (3) makes each exponent zero. The descent criterion for good moduli spaces therefore gives a line bundle L_V on $M_X^{\text{ss}}(r, d)$ (Proposition 4.9). On a fixed-determinant fiber, L_V depends only on $\operatorname{rk} V$ and $\deg V$, although the section s_V still depends on the particular bundle V . That distinction provides many sections of one line bundle.

Semiampleness and separation

Put $h = \gcd(r, d)$, $r_1 = r/h$, and $d_1 = d/h$. For $m \geq 1$ choose

$$\operatorname{rk} V = mr_1, \quad \deg V = mr_1(g - 1) - md_1, \quad (5)$$

which is exactly condition (3). Proposition 5.2 says that, for a stable E and sufficiently large m , a general stable V as in (5) makes $E \otimes V$ cohomology-free. Intersecting these nonempty open conditions over the stable summands of a polystable E gives a section nonvanishing at $[E]$. Hence a power of L_V is basepoint-free.

The technical heart is Lemma 5.8. A hypothetical map $V^\vee \rightarrow E$ with lower-slope image F gives an extension $0 \rightarrow G \rightarrow V^\vee \rightarrow F \rightarrow 0$. Boundedness and Riemann–Roch show that the parameter space for all such extensions has dimension strictly smaller than the moduli of stable V once m is large: the coefficient of m in the difference is $kr_1(\mu(F) - \mu(E)) < 0$. Thus a general V avoids every bad image. The variants with a one-unit degree perturbation in Lemma 5.9 also control the dimensions and surjectivity of maps to finitely many stable summands.

For separation (Proposition 5.4), suppose a polystable E has a stable summand E_0 absent from E' . Starting with the perturbed V , an elementary transformation at a point $x \in X$ chooses a hyperplane so that $E_0^\vee \rightarrow V$ factors through the transformed bundle F , while no map from the dual of a stable summand of E' does. A Grassmannian dimension count and Bertini–Kleiman produce such a hyperplane. Therefore

$$H^0(X, F \otimes E_0) \neq 0, \quad H^0(X, F \otimes E') = 0, \quad (6)$$

so the corresponding determinant section separates the two points.

The basepoint-free power defines a quasi-finite map from the fixed-determinant proper algebraic space to projective space: (6) separates different sets of stable summands, and only finitely many multiplicity patterns remain for a fixed set. Proper plus quasi-finite is finite; pulling back $\mathcal{O}(1)$ is ample and proves projectivity. Finally the determinant map to the projective Jacobian Pic_X^d and fiberwise ampleness give the degree- d statement. For $r = 1$, this reduces to the familiar toy model $M_X^{\text{ss}}(1, d) = \operatorname{Pic}_X^d$, while a fixed determinant is a single point. The argument is specifically for curves in characteristic zero; the cited good-moduli-space existence theorem is the source of that restriction.