

On the realizability of arithmetic degrees of morphisms

Brett Nasserden

arXiv:2212.01909

<http://arxiv.org/abs/2212.01909v1>

Which spectral values occur arithmetically?

Let $X/\overline{\mathbb{Q}}$ be a normal projective variety, let $f : X \rightarrow X$ be a surjective endomorphism, and choose an ample height h_H . The arithmetic degree of $P \in X(\overline{\mathbb{Q}})$ is

$$\alpha_f(P) = \lim_{n \rightarrow \infty} \max\{1, h_H(f^n(P))\}^{1/n}. \quad (1)$$

It is independent of H . Every value in (1) is either 1 or the modulus of an eigenvalue of $f^* : N^1(X)_{\mathbb{R}} \rightarrow N^1(X)_{\mathbb{R}}$. Nasserden calls an eigenvalue μ with $|\mu| > 1$ a *potential arithmetic degree*, and calls it realizable if $\alpha_f(P) = |\mu|$ for some algebraic point P . The converse to the spectral restriction is false in general: the paper constructs abelian varieties with a middle eigenvalue that no point realizes.

Main theorem (Nasserden, Theorem 5.21). *Let X be a projective $\mathbb{Q}$ -factorial toric variety over $\overline{\mathbb{Q}}$, and let $f : X \rightarrow X$ be any surjective endomorphism; f need not preserve the torus action. Then every eigenvalue μ of f^* on $N^1(X)_{\mathbb{R}}$ with $|\mu| > 1$ is realizable: there is $P \in X(\overline{\mathbb{Q}})$ such that*

$$\alpha_f(P) = |\mu|. \quad (2)$$

Thus the Neron–Severi spectrum gives exactly all nontrivial candidate growth rates for toric endomorphisms. This is stronger than realizing only the largest eigenvalue, which follows from general results.

The canonical-height engine

After replacing f by an iterate, Proposition 3.2 allows one to work with a positive eigenvalue $\lambda = |\mu|^m$ and later take m th roots. The iterate may be chosen so that f^* fixes every ray of the rational polyhedral nef cone. Let D_λ be a nef eigendivisor and choose a facet of $\text{Nef}(X)$ not containing it. Toric Mori theory associates an extremal contraction

$$\begin{array}{ccc} X & \xrightarrow{f} & X \\ \phi \downarrow & & \downarrow \phi \\ Y & \xrightarrow{g} & Y \end{array} \quad (3)$$

after a further iterate. The chosen facet is $\phi^* \text{Nef}(Y)$, so D_λ supplies the spectral direction lost under contraction.

Propositions 5.4 and 5.5 provide the decisive height calculation. Choose an ample H on Y so that $A = D_\lambda + \phi^* H$ is ample. If Q is fixed by g and the fiber $F = \phi^{-1}(Q)$ is positive-dimensional, then $\phi^* H|_F = 0$. The canonical height $\widehat{h}_{D_\lambda}$ cannot vanish identically on F : otherwise the ample height $\widehat{h}_{D_\lambda} + h_H \circ \phi$ would be constant there. Hence some $P \in F$ has $\widehat{h}_{D_\lambda}(P) \neq 0$, and

$$h_A(f^n(P)) = \lambda^n \widehat{h}_{D_\lambda}(P) + h_H(Q). \quad (4)$$

Taking n th roots in (4) realizes λ . For a fibering contraction, one uses a fixed point on the base. For a birational contraction, one takes Q in the image of the exceptional locus, so its fiber is again large enough for the same nonvanishing argument.

Why a suitable fixed fiber always exists

Lemma 5.20 proves that every surjective endomorphism of a projective $\mathbb{Q}$ -factorial toric variety has a preperiodic point. Its induction on dimension follows the toric minimal model program. If a K_X -negative birational extremal contraction exists, the exceptional locus is an invariant torus-orbit closure after iteration; it is a lower-dimensional $\mathbb{Q}$ -factorial toric variety. If X is a Mori fiber space $\phi : X \rightarrow Y$, induction first gives a preperiodic point of g on Y and then applies again to the invariant toric fiber. The remaining Picard-number one case is Lemma 5.19: maps of degree greater than one have dense preperiodic points, while an automorphism is reduced through invariant singular strata to projective space, where an eigenvector gives a fixed point. Iteration turns each preperiodic point into the fixed point required in (4).

The proof of Theorem 5.21 now has two branches. If ϕ is birational, equivariance preserves the exceptional locus and its image; Lemma 5.20 and Proposition 5.5 realize the missing eigenvalue. If ϕ is fibering, the same lemma on Y and Proposition 5.4 realize it on a fiber. Every nef-cone facet of a projective toric variety is contractible in this manner, which is the special toric feature closing the argument.

A product model and the scope

For $X = \mathbb{P}^1 \times \mathbb{P}^1$ and $f = (g_a, g_b)$ with $\deg g_a = a > 1$ and $\deg g_b = b > 1$, the two rulings are eigendirections of f^* with eigenvalues a and b . To realize a , take a point P_1 with $\alpha_{g_a}(P_1) = a$ and a preperiodic point P_2 for g_b ; then

$$\alpha_f(P_1, P_2) = \max\{a, 1\} = a.$$

This is the transparent version of the general proof: freeze the unwanted directions on an invariant fiber and let one canonical height grow.

The theorem concerns eigenvalues of modulus greater than one and existence of some algebraic realizing point; it does not assert that the realizing orbit is Zariski dense or classify the height-zero locus. The paper separately proves the small-arithmetic-nondensity conjecture for equivariant toric maps and develops partial extensions to varieties admitting int-amplified maps. Those statements are not needed for the non-equivariant toric theorem above.