

On the Arithmetic Dynamics of Monomial Maps

Jan-Li Lin

arXiv:1704.02661

<http://arxiv.org/abs/1704.02661v1>

Problem and setup

Let $X = X(\Delta)$ be a projective toric variety over $\overline{\mathbf{Q}}$ with dense torus $T \simeq \mathbf{G}_m^n$. An integral matrix $A \in \text{Mat}_n(\mathbf{Z})$ with $\det A \neq 0$ defines a dominant monomial map $\varphi_A : T \rightarrow T$, hence an equivariant rational self-map $\varphi_A : X \dashrightarrow X$. Write X_φ for the points whose entire forward orbit avoids indeterminacy. For an ample Weil height h_X and $h_X^+ = \max\{h_X, 1\}$, the arithmetic degree is

$$\alpha_\varphi(P) = \lim_{k \rightarrow \infty} h_X^+(\varphi^k(P))^{1/k},$$

provided the limit exists. The issue is not the dense torus, where the answer was known, but the boundary: an orbit can move among different torus strata before becoming periodic.

Main theorem (Theorem A). *Factor the characteristic polynomial of A into distinct monic irreducible factors $f_1, \dots, f_s \in \mathbf{Z}[t]$, and put $\rho(f) = \max\{|\lambda| : f(\lambda) = 0\}$. Then $\alpha_\varphi(P)$ exists for every $P \in X_\varphi(\overline{\mathbf{Q}})$, and*

$$\{\alpha_\varphi(P) : P \in X_\varphi(\overline{\mathbf{Q}})\} = \{1, \rho(f_1), \dots, \rho(f_s)\}.$$

Thus the boundary creates no new growth rates. In particular the first three standard arithmetic-degree conjectures—existence, algebraic integrality, and finiteness of the set of values—hold for these maps on arbitrary projective toric compactifications.

The geometric reduction

The orbit–cone correspondence writes X as finitely many tori O_σ . The linear map A sends a cone σ into a cone τ precisely when φ_A is defined on all of O_σ and maps it monomially to O_τ ; otherwise the whole stratum is indeterminate. Deleting cones that eventually reach indeterminacy leaves a subfan $\tilde{\Delta}_A$ with $X_\varphi = X(\tilde{\Delta}_A)$. Its finite directed graph forces every surviving stratum to enter a periodic cycle. Removing a finite initial segment does not change an arithmetic degree, so one may take

$$P \in O_0 \longrightarrow O_1 \longrightarrow \cdots \longrightarrow O_{p-1} \longrightarrow O_0.$$

On each O_j , the return map φ^p is a monomial self-map of a torus. Heights induced by any two projective toric compactifications are comparable up to positive constants and $O(1)$ (Lemma 3.1), so the known torus calculation applies without changing exponential growth.

Three proof ingredients

1. **Residue subsequences exist.** For each $0 \leq j < p$, the sequence along times $kp + j$ has limit $\alpha_{\varphi^p}(\varphi^j(P))^{1/p}$. A priori the full sequence could therefore have p different limits.
2. **Semiconjugacy identifies them.** Adjacent return maps form a commuting square of dominant monomial maps. For a torus point, its arithmetic degree equals the dynamical degree on the identity component of the smallest algebraic subgroup containing its orbit. The vertical monomial map carries that component onto the corresponding component below; on character lattices the two return matrices are conjugate over $\mathbf{Q}$. Their spectral radii, and hence consecutive residue limits, agree. This closes the gap between convergence on each stratum and convergence of the original orbit.
3. **The return-map spectrum descends from A .** Let N be the lattice of one-parameter subgroups of the dense torus and $N_\sigma = N \cap \text{span}_{\mathbf{R}}(\sigma)$. If $O_0 = O_\sigma$, then $A^p N_\sigma \subset N_\sigma$, and the return map on O_σ is represented on rational cocharacters by the induced operator

$$\overline{A^p} : (N/N_\sigma)_{\mathbf{Q}} \longrightarrow (N/N_\sigma)_{\mathbf{Q}}.$$

Let G_0 be the identity component of the smallest algebraic subgroup containing the φ^p -orbit of P , with rational cocharacter space $W \subset (N/N_\sigma)_{\mathbf{Q}}$. The torus result gives $\alpha_{\varphi^p}(P) = \rho(\overline{A^p}|_W)$. This restriction is a rational invariant subquotient of A^p , so each of its eigenvalues has the form λ^p for an eigenvalue λ of A . If λ^p realizes the spectral radius, rationality also puts every Galois conjugate $(\lambda')^p$ in the same characteristic polynomial. Hence the largest modulus is $\rho(f_j)^p$, where f_j is the irreducible factor containing λ . Thus $\alpha_\varphi(P) = \alpha_{\varphi^p}(P)^{1/p} = \rho(f_j)$. All listed values already occur on the dense torus, proving equality rather than mere containment.

A model to keep in mind

For $A = \text{diag}(2, 3)$ on $(\mathbf{P}^1)^2$, $\varphi(x, y) = (x^2, y^3)$ and the only arithmetic degrees are 1, 2, 3. A non-torsion y -coordinate gives rate 3; if y is torsion but x is not, the rate is 2; torsion in both coordinates gives 1. Boundary strata such as $x = 0$ merely discard a coordinate and inherit the appropriate smaller monomial map. The general proof replaces this fixed coordinate picture by a periodic cycle of toric strata and the invariant-subspace lemma.

Scope and hardest point

The same orbit graph also yields dynamical Mordell–Lang on arbitrary toric varieties, while a quotient by the connected subgroup governing preperiodic points yields the paper’s dynamical Manin–Mumford criterion. These are separate theorems, not consequences of the height statement alone. The delicate step for Theorem A is precisely that a boundary stratum may be periodic only after p steps: one must both synchronize the p height subsequences and recover their spectral radius from A , rather than silently treating the stratum as A -invariant.