Singularities of Cox Rings of Fano Varieties

Morgan V. Brown

arXiv:1109.6368

http://arxiv.org/abs/1109.6368v2

Problem and setup

Choose Weil divisors $D_1, \dots, D_r$ forming a basis of the torsion-free part of $\mathrm{Cl}(X)$. The corresponding total coordinate ring is

$$\mathrm{Cox}(X; D_1, \dots, D_r) = \bigoplus_{a \in \mathbf{Z}^r} H^0\left(X, \mathcal{O}_X\left(\sum_i a_i D_i\right)\right).$$

Although X may be smooth, this multigraded affine ring can be singular. The paper relates its singularities to the birational positivity of X.

Main theorem (Theorem 1.2)**.** *Let (X, Δ) be a $\mathbf{Q}$-factorial log Fano pair over $\mathbf{C}$. Then $\mathrm{Cox}(X; D_1, \dots, D_r)$ is normal with log terminal singularities, hence Cohen–Macaulay. If X is a smooth complete Fano variety, then $\mathrm{Cox}(X)$ is Gorenstein with canonical singularities.*

For the second assertion, smooth Fano varieties have finitely generated free class group; a cited computation makes the Cox ring's canonical module free of rank one. Thus it is Gorenstein, and log terminal singularities are canonical in the Gorenstein case. The content of the paper is the first assertion, proved by induction on the Picard rank.

Proof mechanism

1. The rank-one cone. When $\rho(X) = 1$, the Cox ring is a section ring of an ample generator L. Theorem 2.1 resolves its affine cone through the total space of L and a resolution of X. If $-K_X = mL$ in the reduced log Fano calculation, the exceptional zero section has discrepancy $m - 1 > -1$; the remaining discrepancies come from X. This is the induction base (up to the finite cover needed to pass from a Picard generator to a class-group generator).

2. Lower the Picard rank geometrically. Log Fano varieties are Mori Dream Spaces. For $\rho(X) > 1$, choose a Cartier divisor L with neither L nor $-L$ effective and transverse to the Mori chamber walls. In

$$Y = \mathbf{P}_X(\mathcal{O}_X \oplus \mathcal{O}_X(L)), \qquad Y_0 = Y \setminus (E_0 \cup E_\infty),$$

Y_0 is the associated $\mathbf{G}_m$-bundle. Moving through Mori chambers contracts the two boundary sections and produces a projective small compactification X' of Y_0 with $\rho(X') = \rho(X) - 1$.

Theorem 4.1 proves that (X', Δ') is again $\mathbf{Q}$-factorial log Fano. Chamber transversality gives $\mathbf{Q}$-factoriality, while the strict transform of $-\pi^*(K_X + \Delta)$ is ample. For small $\varepsilon > 0$ the boundary pair

$$(Y,\ \pi^*\Delta + (1 - \varepsilon)(E_0 + E_\infty))$$

is klt by an explicit discrepancy comparison on a lifted log resolution. The flips and divisorial contractions from Y to X' are log-MMP steps, so klt singularities survive.

3. Relate the two Cox rings. If $L = mD_r$, Theorem 3.3 identifies the Cox ring of X' with the $\mathbf{Z}/m$-invariant subring of $\mathrm{Cox}(X; D_1, \dots, D_r)$. Thus the original Cox ring is recovered from the lower-rank one by a cyclic cover. Proposition 6.1 first proves normality directly using divisorial valuations on Laurent monomials.

4. Cross the cover without worsening singularities. Lemma 6.2 embeds both Cox rings into finite abelian covers of the same Picard-graded section ring and proves that these covers are étale in codimension two. Log terminal singularities are invariant under such a finite cover. The induction hypothesis for X' therefore transfers to X, completing Theorem 1.2.

Toy model, difficulty, and scope

For $X = \mathbf{P}^1 \times \mathbf{P}^1$ and $L = (1, -1)$, the two boundary divisors of $\mathbf{P}_X(\mathcal{O} \oplus \mathcal{O}(L))$ contract to lines and the small compactification is $\mathbf{P}^3$. Both Cox rings are polynomial rings in four variables; here $m = 1$, so the cyclic-cover step is invisible. This isolates the general proof's geometry while removing its singular and covering complications.

The delicate point is not finite generation alone but preserving klt discrepancies while reducing rank and then returning through a cover. The theorem does not say that every finitely generated Cox ring is Cohen–Macaulay; the introduction gives hyperkähler counterexamples. The related log Calabi–Yau statement (Theorem 1.3) assumes that X is a Mori Dream Space and yields log canonical, rather than log terminal, singularities.

Codex mathematical paper digest, 2026-10-28. Source paper should be consulted for full statements and proofs.