

Moduli spaces of rational curves on Fano threefolds

Roya Beheshti, Brian Lehmann, Eric Riedl, Sho Tanimoto

arXiv:2010.06795

<http://arxiv.org/abs/2010.06795v3>

A finiteness theorem across all degrees

Work over an algebraically closed field of characteristic 0. For a smooth Fano threefold X , let $\text{Rat}(X)$ be the union of those components of $\overline{M}_{0,0}(X)$ whose general stable map has irreducible domain. The degree of a curve C is its anticanonical degree $-K_X \cdot C$.

Main theorem (Beheshti–Lehmann–Riedl–Tanimoto, Theorems 1.1 and 7.8). *For every smooth Fano threefold X , there is a polynomial $P(d)$ such that the number of components of $\text{Rat}(X)$ parametrizing curves of anticanonical degree at most d is at most $P(d)$.*

The polynomial depends on X and is not asserted to be optimal. The result proves the polynomial-growth bound suggested by Batyrev’s heuristic and, more structurally, reduces the study of high-degree free curves to finitely many low degrees.

The proof separates components whose curves dominate X from those sweeping only a proper subvariety. The two halves require rather different geometry.

Non-dominant families live on controlled surfaces

For a big and nef divisor L on a smooth projective variety Y , the Fujita invariant is

$$a(Y, L) = \inf\{t \in \mathbb{R} : K_Y + tL \text{ is pseudo-effective}\}. \quad (1)$$

Here $a(X, -K_X) = 1$. Theorem 4.2, imported from earlier work, places every non-dominant component inside the proper closed union of subvarieties $Y \subset X$ with $a(Y, -K_X) > 1$. The paper’s Theorem 4.1 then classifies the surface components: each is either swept out by $-K_X$ -lines or is an exceptional divisor of one of Mori’s contraction types $E1$ – $E5$. Consequently only finitely many surfaces Y_i need be counted.

On a resolution S_i of a contractible Y_i , the finite list of polarized surfaces has an irreducible rational-curve space in every numerical class. A line-swept S_i is ruled (Lemma 4.5): over a base of positive genus its only rational curves are fibers, while over $\mathbb{P}^1$ it is a Hirzebruch surface, hence toric, with at most one family in each numerical class. Lattice counting against the big and nef divisor $-K_X|_{S_i}$ is therefore polynomial (indeed at most linear at each fixed degree). This controls every non-dominant component.

The dominant engine: Movable Bend-and-Break

The original technical input is stronger than a mere count.

Main theorem (Movable Bend-and-Break, Theorem 1.4). *Let $M \subset \text{Rat}(X)$ generically parametrize free curves. If the general curve has anticanonical degree at least 6, then M contains a stable map from a two-component curve such that both restrictions are free curves on X .*

Thus a high-degree component meets a boundary stratum obtained by gluing two lower-degree free families. Iterating this decomposition makes low-degree families into generators. The final proof of Theorem 7.8 invokes this result together with the combinatorial component bound of [LT19b, Theorem 5.13] to obtain a polynomial bound for all dominant families.

How the breaking family is forced

Suppose first that a general free map is birational onto its image and is an immersion, with normal bundle

$$N_{f/X} \simeq \mathcal{O}(a) \oplus \mathcal{O}(b), \quad a \leq b, \quad -K_X \cdot C = a + b + 2. \quad (2)$$

Lemma 6.4 shows that passing through r general points costs $2r$ conditions, while meeting s general members of a basepoint-free family of curves costs s . The authors choose r, s so the incidence locus is one-dimensional. Mori’s Bend-and-Break then supplies a reducible limit. Degree bookkeeping in Proposition 6.5 shows that a limit with at least two free components has only four possible shapes:

- (i) exactly two free components;
 - (ii) two free components joined by an $E5$ -line;
 - (iii) free components plus two $E5$ -lines;
 - (iv) free components plus an exceptional conic.
- (3)

The incidence conditions leave too little degree or freedom for any other non-free component. If the restricted tangent bundle has only one positive summand, Propositions 6.1–6.2 instead identify the maps as multiple covers of a free anticanonical conic and break the cover directly.

Why the exceptional $E5$ line is the hard point

When X has no $E5$ contraction, the authors contract the disjoint relevant $E1$, $E3$, and $E4$ divisors and choose the basepoint-free incidence curves as pullbacks from the contraction. They avoid every possible non-free piece in (3), leaving only (i) (Theorem 6.7).

An $E5$ divisor is different: its lines move in dimension greater than one and can connect two otherwise disjoint free curves. The obstruction is real. On $\mathbb{P}_{\mathbb{P}^2}(\mathcal{O} \oplus \mathcal{O}(2))$, the lines in the minimal family of moving $\mathbb{P}^2$ sections form a degree-5 curve family that breaks into two bundle fibers joined by a line in the rigid $E5$ divisor, not into two free curves (Example 6.8).

There are exactly six smooth Fano threefold types admitting an $E5$ contraction. For the five with a unique $E5$ divisor, Proposition 6.10 uses induction: if a connector $C_1 \cup \ell \cup C_2$ occurs and one C_i already breaks, transversality and smoothness of the stable-map point let one regroup and smooth the chain into the desired two free pieces. Only degrees 6–9 remain, and Proposition 6.11 checks them from the classification. The unique threefold with two $E5$ divisors is handled separately in Claim 8.14 using six generators of its nef monoid, low-degree uniqueness, and explicit treatment of the remaining connector classes. This is why the uniform theorem starts at degree 6.

Toy model and proof roadmap

For a conic in $\mathbb{P}^3$, a pencil can specialize it to two intersecting lines. Both lines are free, and smoothing their node returns to the original component. Movable Bend-and-Break is this picture with two extra burdens: one must impose enough incidences to force breaking inside an arbitrary component, and then exclude the non-free exceptional pieces in (3).

The argument for the polynomial bound can be summarized as

$$\begin{array}{lll} \text{non-dominant families} & \longrightarrow & \text{finitely many ruled or contractible surfaces} \longrightarrow \text{polynomial surface count,} \\ \text{dominant free families} & \longrightarrow & \text{two-free-component boundary} \longrightarrow \text{low-degree generation and polynomial count.} \end{array} \tag{4}$$

Alongside this, Theorem 1.3 says that when $-K_X$ is very ample, a dominant universal-family evaluation with non-irreducible general fiber comes only from conic images or curves lying in fibers of a del Pezzo fibration.

Scope

The theorem concerns a fixed smooth Fano threefold and the subspace $\text{Rat}(X)$; it does not count components whose general stable map already has reducible domain. Neither very ampleness of $-K_X$ nor a Picard-rank hypothesis is needed for Theorem 7.8 or Movable Bend-and-Break, although very ampleness is used for some finer Manin-component statements. The paper establishes an upper bound, not the conjectural exact asymptotic or the optimal exponent, and its dominant-family count explicitly depends on the earlier result [LT19b, Theorem 5.13].