

Dynamical modular curves for quadratic polynomial maps

John R. Doyle

arXiv:1707.02257

<http://arxiv.org/abs/1707.02257v1>

From orbit portraits to modular curves

Write $f_c(z) = z^2 + c$. The finite preperiodic points of f_c form a directed graph (the fixed point at infinity is omitted). Doyle calls a finite graph G *admissible* when every vertex has out-degree one and in-degree zero or two, and when, for every $N \geq 2$, it has at most

$$R(N) = \frac{1}{N} \sum_{d|N} \mu(N/d) 2^d$$

cycles of length N . These are exactly the evident unramified constraints coming from a quadratic polynomial. “Strongly admissible” additionally requires zero or two fixed points.

Choose a minimal set of vertices $P_1, \dots, P_r$ generating G . A generator whose orbit is new is imposed by a generalized dynatomic equation $\Phi_{M_i, N_i}(x_i, t) = 0$; a generator whose orbit attaches to the preceding graph is imposed by an equation $f_t^{\kappa_i}(x_i) + f_t^{\lambda_i}(x_{j_i}) = 0$. After removing loci where two intended components collide or the intended exact portrait becomes smaller, the resulting open curve $U_1(G)$ parametrizes f_c with marked points generating a graph isomorphic or nearly isomorphic to G ; exact isomorphism holds generically. Its affine closure is $Y_1(G)$, and $X_1(G)$ is the normalization of the projective closure.

Main theorem (Doyle, Theorem 1.7). *Let K be a field of characteristic zero. For every admissible graph G , the dynamical modular curve $X_1(G)$ is irreducible over K .*

Thus the graph conditions do not secretly split the parameter space into different algebraic families. As an arithmetic consequence (Theorem 4.1), every strongly admissible G actually occurs as the full graph $G(f_c, K)$ for some number field K and some $c \in K$.

Ingredient I: one eventual period and the generic Galois action

Bousch’s theorem says that every $\Phi_{M,N}(x, t)$ is irreducible over $\mathbf{C}$, and that its splitting-field Galois group consists of all permutations of the roots commuting with f_t . First suppose every cycle of G has the same length N . Graphs generated only by periodic points are irreducible because the Galois group freely permutes the N -cycles and rotates each cycle.

For the induction step, remove a pair $\{P, -P\}$ of maximal preperiod $M \geq 2$, obtaining H . Choose $k \geq 0$ and a generator P_1 of H so that $f(P) = -f^k(P_1)$. Over $\mathbf{C}(X_1(H))$, restoring the pair means solving one quadratic equation

$$h(z) = f_t(z) + f_t^k(x_1) = 0. \quad (1)$$

Its roots $\pm y$ have portrait (M, N) and a common image. Bousch’s Galois description supplies an automorphism that transposes y and $-y$ while fixing every marked point already in H . Hence neither root lies in $\mathbf{C}(X_1(H))$, so (1) stays irreducible. This proves irreducibility for all graphs with a single eventual period.

Ingredient II: different periods have independent branching

Let $G_{M,N}$ denote the full graph containing every point of eventual period N and preperiod at most M . Replacing an arbitrary G by its normal closure reduces the theorem to

$$G = \bigsqcup_{i=1}^s G_{M_i, N_i}, \quad N_i \neq N_j \ (i \neq j),$$

and the modular curve is birational to the following fiber product:

$$X_1(G) \dashrightarrow X_1(G_{M_1, N_1}) \times_{\mathbf{P}^1} \cdots \times_{\mathbf{P}^1} X_1(G_{M_s, N_s}). \quad (2)$$

This birational description is enough for irreducibility. Irreducible factors alone do not make (2) irreducible. The decisive point is that the covers in (2) have pairwise disjoint branch loci over the affine t -line. Indeed, if α is a multiple root of $\Phi_{M,N}(x, c)$, the chain rule forces either a multiple root of Φ_N (a parameter at the root of a period- N hyperbolic component) or $f_c^k(\alpha) = 0$ for some $0 \leq k \leq M - 2$. In the second case the orbit of α hits the critical point, so 0 is preperiodic; after the first alternative is excluded, its eventual period is exactly N . A fixed c cannot satisfy either alternative for two different eventual periods. The only common branch point is therefore infinity.

A Riemann–Hurwitz argument now says that two such Galois closures are linearly disjoint over $\mathbf{C}(t)$: a nontrivial intersection would be a cover of $\mathbf{P}^1$ ramified at at least two places, including a finite place common to both covers. Induction makes (2) irreducible. The Lefschetz principle then gives the assertion over every characteristic-zero field.

Proof roadmap and a toy model

The proof has a clean two-level architecture:

1. Bousch’s maximal generic Galois action prevents a same-period graph from splitting when successive pairs of preimages are adjoined.
2. Distinct eventual periods have disjoint affine ramification, so their already-irreducible parameter curves remain irreducible after fiber product.

For a toy model, mark two distinct 3-cycles. The naive equations $\Phi_3(x_1, t) = \Phi_3(x_2, t) = 0$ also contain the diagonal loci $x_1 = f_t^j(x_2)$ for $j = 0, 1, 2$; those points mark the same cycle twice. Removing them leaves the single component that genuinely parametrizes two disjoint cycles. This is exactly why $U_1(G)$ is defined by both equations and noncollision conditions.

The full generic preperiodic Galois group

Let L be the extension of $\mathbf{C}(t)$ generated by every preperiodic point of f_t , and let $\mathcal{G}$ be the direct limit of all admissible preperiodic graphs. Doyle’s stronger Theorem 5.2 says

$$\mathrm{Gal}(L/\mathbf{C}(t)) \simeq \mathrm{Aut}(\mathcal{G}). \quad (3)$$

For each finite normal graph H , the preceding argument identifies $\mathbf{C}(X_1(H))/\mathbf{C}(t)$ as Galois with group $\mathrm{Aut}(H)$. Passing from the directed union of these fields to the inverse limits of their Galois and graph-automorphism groups gives (3). If $G_f = G(f, \bar{K})$ is the full preperiodic graph over an algebraic closure, this motivates the representation $\rho_{f,K} : \mathrm{Gal}(\bar{K}/K) \rightarrow \mathrm{Aut}(G_f)$. Theorem 5.7 proves that its image has finite index when K is a finitely generated extension of $\mathbf{C}$ and $f \in K[z]$ is a non-isotrivial quadratic polynomial.

Hardest point and scope

The subtle step is not the irreducibility of one dynatomic polynomial, but showing that structures with different eventual periods do not acquire a common subcover; the branch-locus lemma is what converts a dynamical multiple-root analysis into linear disjointness. At boundary points of $X_1(G)$ the intended exact portrait can become smaller, and at postcritically finite parameters even a point of $U_1(G)$ may mark a graph only nearly isomorphic to G ; exact isomorphism holds away from finitely many such fibers. The realization theorem needs strong admissibility, and the paper gives examples showing that the analogous assertion can fail for nearly or weakly admissible graphs.