

Fano varieties in Mori fibre spaces

Giulio Codogni, Andrea Fanelli, Roberto Svaldi, Luca Tasin

arXiv:1406.7634

<http://arxiv.org/abs/1406.7634v2>

Which Fano varieties can be fibres?

A Mori fibre space (MFS) is a dominant projective morphism $f: X \rightarrow Y$ with connected fibres and $\dim Y < \dim X$, where X is $\mathbb{Q}$ -factorial with klt singularities, $\rho(X/Y) = 1$, and $-K_X$ is f -ample. A terminal $\mathbb{Q}$ -factorial Fano variety F is *fibre-like* if it occurs as a sufficiently general fibre in the precise open family of Definition 2.14.

Write $N^1(F)_{\mathbb{Q}}$ for divisor classes modulo numerical equivalence. The finite group $\text{Mon}(F)$ is the maximal integral linear symmetry group preserving the canonical line, the intersection form, the nef cone, and the contraction type and deformation data attached to its faces. Let $\text{HMon}(F)$ be the image of $\text{Aut}(F)$ on $N^1(F)$.

Main theorem (Codogni–Fanelli–Svaldi–Tasin, Theorem 1.2). *Let F be a terminal $\mathbb{Q}$ -factorial Fano variety.*

1. *If $N^1(F)_{\mathbb{Q}}^{\text{Aut}(F)} = \mathbb{Q}[K_F]$, then F is fibre-like; it occurs in an isotrivial MFS over a curve.*
2. *If $\dim N^1(F)_{\mathbb{Q}}^{\text{Mon}(F)} > 1$, then F is not fibre-like.*
3. *If $H^1(F, T_F) = 0$, so F is rigid, then F is fibre-like if and only if $N^1(F)_{\mathbb{Q}}^{\text{Aut}(F)} = \mathbb{Q}[K_F]$.*

Thus relative Picard rank one does not force the geometric fibre itself to have Picard rank one. It says instead that monodromy must collapse the fibre’s divisor theory to the canonical line.

Ingredient I: monodromy remembers the MMP

For a suitable open $U \subset Y$, the divisor classes of the fibres form a local system with finite monodromy. Theorem 2.5 extends the relevant part of Deligne’s invariant-cycle principle to the singular setting: under the paper’s $\mathbb{Q}$ -factoriality, rational-singularity, and relative Fano hypotheses, restriction is surjective,

$$N^1(X)_{\mathbb{Q}} \longrightarrow N^1(F_t)_{\mathbb{Q}}^{\pi_1(U,t)}. \quad (1)$$

After shrinking U , Theorem 2.6 shows that monodromy preserves the nef cone. The numerical case analysis in Theorem 2.7 reads top self-intersections, exceptional divisors, and image dimensions on each face to preserve whether the corresponding MMP step is divisorial, flipping, or fibre type. Separately, after a finite étale cover, a relative Cartier divisor and a flat relative Proj put the targets of monodromy-related contractions in one deformation family. Theorem 2.12 packages these facts as factorization through $\text{Mon}(F_t)$; for an isotrivial family it factors through $\text{HMon}(F_t)$.

Ingredient II: relative rank one forces one invariant

Suppose $F = F_t$ is a fibre of an MFS. Surjectivity in (1), connectedness of the fibres, and $\rho(X/Y) = 1$ give the exact sequence

$$0 \longrightarrow N^1(Y)_{\mathbb{Q}} \xrightarrow{f^*} N^1(X)_{\mathbb{Q}} \longrightarrow N^1(F)_{\mathbb{Q}}^{\pi_1(U,t)} \longrightarrow 0. \quad (2)$$

Consequently the last term is one-dimensional. Since the actual monodromy factors through $\text{Mon}(F)$ and K_F is fixed,

$$N^1(F)_{\mathbb{Q}}^{\text{Mon}(F)} = \mathbb{Q}[K_F]. \quad (3)$$

This proves the necessary criterion. Its force is that it also applies to nonisotrivial Mori fibre spaces.

Ingredient III: manufacture the sufficient MFS

Assume the automorphism-invariant subspace is exactly $\mathbb{Q}[K_F]$. The image $\text{HMon}(F)$ is finite because it permutes the faces of the rational polyhedral nef cone. Choose generators $[f_1], \dots, [f_g]$ of this image and representatives $f_i \in \text{Aut}(F)$, and set $G = \langle f_1, \dots, f_g \rangle$. For a genus- g projective curve C , map the standard generators of $\pi_1(C)$ by $a_i \mapsto f_i$ and $b_i \mapsto f_i^{-1}$; the surface-group relation is satisfied and the image is G . Using this actual action on F , form the diagonal quotient

$$X = (F \times \tilde{C})/\pi_1(C). \quad (4)$$

It is an isotrivial family over C with fibre F . The action of G on $N^1(F)$ has image $\text{HMon}(F)$, so their invariant subspaces agree. An equivariant anticanonical embedding places (4) inside the corresponding projective space bundle, proving projectivity. The terminal and $\mathbb{Q}$ -factorial properties descend in the form needed by the construction. Finally, restriction to a fibre fits into

$$0 \longrightarrow N^1(C)_{\mathbb{Q}} \longrightarrow N^1(X)_{\mathbb{Q}} \longrightarrow N^1(F)_{\mathbb{Q}}^{\text{HMon}(F)} \longrightarrow 0. \quad (5)$$

The seesaw principle supplies exactness on the right. The two outer spaces are one-dimensional, so $\rho(X) = 2$ and $\rho(X/C) = 1$; together with relative anticanonical ampleness, (4) is the required MFS.

Why rigidity closes the gap

In a general nonisotrivial family, geometric monodromy may lie in the larger group $\text{Mon}(F)$ rather than come from automorphisms of one fixed fibre. If $H^1(F, T_F) = 0$, local deformations of F are trivial, so the family is isotrivial and its monodromy factors through $\text{HMon}(F)$. The necessary argument then gives the automorphism-invariant condition, while the quotient construction gives the converse. This is Theorem 3.5.

A rank-two model and the boundary of the result

For $F = \mathbb{P}^1 \times \mathbb{P}^1$, write H_1, H_2 for the two ruling classes. The involution exchanging the factors fixes only $\mathbb{Q}(H_1 + H_2) = \mathbb{Q}[-K_F]$, so the sufficient criterion builds an isotrivial MFS with this fibre. In contrast, on $\text{Bl}_p \mathbb{P}^2$ the two extremal contractions have different MMP types (one divisorial, one fibre type); a permitted monodromy cannot exchange them, leaving a two-dimensional invariant space. The necessary criterion rules it out. This matches Theorem 4.1, which says that the only smooth del Pezzo surfaces that are not fibre-like are the blow-ups of $\mathbb{P}^2$ in one or two points.

The sufficient condition is not necessary without rigidity: cubic surfaces supply fibre-like examples not detected by it. The paper also classifies smooth fibre-like Fano threefold deformation types of Picard rank greater than one and develops toric and homogeneous applications, but these use further case-specific geometry. The conceptual bottleneck is Theorem 2.12: turning topological monodromy into an action constrained by the nef cone and every relevant MMP contraction, then combining its invariants with the global relative Picard-rank calculation (2).